

Light-Cone Dynamics of Particles and Fields

Thomas Heinzl¹
Universität Regensburg
Institut für Theoretische Physik
93040 Regensburg
Germany

¹present address: Friedrich-Schiller-Universität Jena, Theoretisch-Physikalisches Institut, Max-Wien-Platz 1, D-07743 Jena, e-mail: heinzl@tpi.uni-jena.de

Abstract

We review the foundations as well as a number of important applications of light-cone dynamics. Beginning with the peculiarities of relativistic particle dynamics we discuss the choice of a time parameter as the gauge fixing within reparametrization invariant dynamical systems. Including Poincaré invariance, we are naturally led to Dirac's forms of relativistic dynamics. Among these, the front form is our main focus as it is the basis for light-cone dynamics. We explain the peculiar features of the light-cone formulation such as boost and Galilei invariance or separation of relative and center-of-mass motion. Combining light-cone dynamics and field quantization leads to the introduction of light-cone quantum field theory. We show how the positivity of the kinematical longitudinal momentum implies the triviality of the light-cone vacuum. We point out that its special features make the light-cone formulation a unique framework to deal with bound states as few-body systems based on quantum field theory. In a first application, we analyze spontaneous symmetry breaking for scalar field theory in 1+1 dimensions. The importance of modes with vanishing longitudinal momentum is elucidated. For fermionic field theories, we suggest to reconstruct vacuum properties, like chiral condensates, from the particle spectrum. The latter can be obtained by solving the light-cone Schrödinger equation as we explicitly demonstrate for the 't Hooft and Schwinger models. Finally, we make contact with phenomenology by calculating the pion wave function within the Nambu and Jona-Lasinio model. We are thus able to predict a number of observables like the pion charge and core radius, the r.m.s. transverse momentum and the pion distribution amplitude. The latter turns out being not very different from the asymptotic one.

Contents

1	Introduction: Strong Interactions	7
1.1	Hadrons	7
1.2	Quantum Chromodynamics	9
1.3	Chiral Symmetry in QCD	12
1.4	Outline	14
I	Foundations	17
2	Relativistic Particle Dynamics	19
2.1	The Free Relativistic Particle	20
2.2	Dirac's Forms of Relativistic Dynamics	27
3	The Front Form	33
3.1	Generalities	33
3.2	Lorentz Boosts	36
3.3	Galilei Subgroups and Consequences	37
4	Light-Cone Quantization of Fields	41
4.1	Construction of the Poincaré Generators	41
4.2	Schwinger's (Quantum) Action Principle	43
4.3	Quantization as an Initial/Boundary-Value Problem	45
4.3.1	The Cauchy Problem	47
4.3.2	The Characteristic Initial-Value Problem	48
4.3.3	The Hamiltonian Formulation	48
4.3.4	The Initial-Boundary-Value Problem	50
4.3.5	Fermions	52
4.4	First-Order Systems	55
5	The Light-Cone Vacuum	58
5.1	Basics	58
5.2	Trivial versus Nontrivial Vacua	60
5.2.1	Constant Source	60
5.2.2	Mass Shift: Bosons	65

5.2.3	Mass Shift: Fermions	67
6	Light-Cone Wave Functions	70
6.1	Definition	70
6.2	Properties	73
II	Applications	77
7	Zero Modes and the Light-Cone Vacuum	79
7.1	The Constrained Zero Mode	80
7.2	Perturbative Solution	82
7.3	Mean-Field Ansatz	84
7.4	The Tamm-Dancoff Approximation	91
7.4.1	General Remarks	91
7.4.2	One-Particle Tamm-Dancoff Approximation	95
7.4.3	Two-Particle Tamm-Dancoff Approximation	97
7.5	Discussion	99
8	't Hooft and Schwinger Model	101
8.1	Prelude: Fermion Condensates	101
8.2	State of the Art	103
8.3	't Hooft's Ansatz	107
8.3.1	Exact Endpoint Behavior	109
8.3.2	Variational Approach	111
8.4	Extension of 't Hooft's Ansatz	113
8.4.1	Purely Variational Approach	114
8.4.2	Variational Approach using the exact β	118
8.5	Comparison With Numerical Results	121
8.6	Discussion	125
9	The Pion Wave Function in the NJL Model	134
9.1	The Condensate	135
9.2	The Light-Cone Bound State Equation	138
9.3	Observables	141
9.4	The Pion Distribution Amplitude	147
	Conclusions	149
	Acknowledgments	152
A	Light-Cone Spinors	153
	Bibliography	156

Chapter 1

Introduction: Strong Interactions

1.1 Hadrons

The word ‘hadron’ originates from Greek where $\alpha\delta\rho\sigma$ means ‘big’ or ‘strong’. It refers to all particles participating in strong interactions. One can therefore say that hadrons are the physical degrees of freedom in strong interaction physics. Everything we know empirically in this area has been obtained by studying hadronic processes and properties. A (very) incomplete list of hadrons is provided by Table 1.1.

There are more than a hundred additional hadrons which, together with their properties can be found in the reviews of the particle data group [1]. One distinguishes between baryons having baryon number $B = 1$ and half-integer spin (fermions), and mesons with $B = 0$ and integer spin (bosons). This great variety of hadrons has been classified in the sixties within the quark model of Gell-Mann and Zweig [2, 3] which states that hadrons are grouped into multiplets of the group $SU(3)_F$, where the ‘ F ’ stands for the quantum number ‘flavor’. (Anti-)Quarks are postulated as the realizations of the fundamental (anti-)triplet of $SU(3)_F$. This implies that quarks come in three flavors, ‘up’ u , ‘down’ d and ‘strange’ s (the anti-quarks being $\bar{u}$, $\bar{d}$ and $\bar{s}$), which combine to give the quantum numbers of the hadrons. The quarks were postulated to be fermions carrying charges which are multiples of a third of the electron charge. Nowadays one knows a total number of $N_F = 6$ quarks with the additional heavy flavors ‘charm’ c , ‘bottom’ b and ‘top’ t . In the quark scheme, mesons consist of a quark and an antiquark, (anti-)baryons of three (anti-)quarks. Thus, for example, one finds the following octet of pseudo-scalar mesons with spin-parity $J^\pi = 0^-$,

$$\begin{aligned} \pi^0 &= \sqrt{1/2}(u\bar{u} - d\bar{d}) & K^0 &= d\bar{s} \\ \pi^- &= d\bar{u} & K^+ &= u\bar{s} \\ \pi^+ &= u\bar{d} & K^- &= s\bar{u} \\ \eta &= \sqrt{1/3}(u\bar{u} + d\bar{d} + s\bar{s}) & \bar{K}^0 &= s\bar{d} \end{aligned} \tag{1.1}$$

This octet structure is reminiscent in the name ‘the eightfold way’, the title of a

Table 1.1: *Masses and quantum numbers of a few hadrons. The upper part consists of baryons ($B = 1$), the lower part of mesons ($B = 0$). Masses are given in MeV.*

particle		mass	isospin	spin	B
proton	p	938	1/2	1/2	1
neutron	n	940	1/2	1/2	1
delta	Δ	1236	3/2	3/2	1
lambda	Λ	1116	0	1/2	1
pion	π	138	1	0	0
kaon	K	495	1/2	0	0
rho	ρ	770	1	1	0

famous book by Gell-Mann and Ne'eman [4].

It was soon realized thereafter, that the quark-flavor model is in conflict with the Pauli principle. The baryon resonance $\Delta^{++} = uuu$, with all quark spins pointing upwards, cannot have an anti-symmetric wave function as it should as a three-fermion state. The loophole is to introduce an additional quantum number called color, giving rise to another symmetry group, $SU(3)_C$. Upon postulating that hadrons should always appear as color singlets the Pauli principle is satisfied.

Using flavor and color symmetry, the wave function of mesons and baryons can thus be written as

$$|M\rangle = \frac{1}{\sqrt{N_C}} M_{ij} \delta_{ab} |q_i^a \bar{q}_j^b\rangle, \quad (1.2)$$

$$|B\rangle = \frac{1}{2\sqrt{N_C}} B_{ijk} \epsilon_{abc} |q_i^a q_j^b q_k^c\rangle. \quad (1.3)$$

Here, $N_C = 3$ is the number of colors (indices a, b, c), M_{ij} and B_{ijk} are tensors in flavor space. It is obvious that the baryon state is anti-symmetric in color which solves the Δ^{++} -puzzle.

If we include spin degrees of freedom and the spatial part of the hadron wave function, the total hadronic state can be written symbolically as

$$|\text{hadron}\rangle = |\text{space}\rangle \otimes |\text{spin}\rangle \otimes |\text{flavor}\rangle \otimes |\text{color}\rangle, \quad (1.4)$$

where the space, spin and flavor part is always symmetric, the color part guaranteeing overall anti-symmetry. The non-spatial part, $|\text{spin}\rangle \otimes |\text{flavor}\rangle \otimes |\text{color}\rangle$, is governed by the symmetry group $SU(2)_S \otimes SU(3)_F \otimes SU(3)_C$. For the positively charged pion π^+ one finds, for instance,

$$|\pi^+\rangle = \frac{1}{\sqrt{2N_C}} (|\bar{d}_{a\uparrow} u_{a\downarrow}\rangle - |\bar{d}_{a\downarrow} u_{a\uparrow}\rangle), \quad (1.5)$$

where the spatial part has been suppressed. The latter will, however, become relevant in later chapters.

It should be pointed out, that the flavor symmetry is not realized in nature as an exact symmetry, because the states in the $SU(3)_F$ multiplets have different masses. The color symmetry, on the other hand, is unbroken.

If one calculates the magnetic moments ($\mu_i = e_i/2M_i$) of the hadrons $i = p, n, \Lambda$ by adding those of their quark constituents, one can determine the *constituent masses* of the quarks

$$M_u = M_d = 340 \text{ MeV} , \quad M_s = 510 \text{ MeV} . \quad (1.6)$$

The constituent quark masses thus roughly sum up to the total bound state (hadron) mass.

At this point, the question arises whether quarks can really be ‘seen’ experimentally. The answer is both ‘yes’ and ‘no’. On the one hand, quarks have never been observed as free particles outside a hadron, a property called *quark confinement*. On the other hand, the deep-inelastic scattering (DIS) experiments beginning in the late sixties have revealed a rich substructure of hadrons, in particular the nucleon (Nobel prize 1990 for Friedman, Kendall, and Taylor). This substructure consists of quasi-free, point-like constituents (‘partons’) which carry the quantum number of quarks.

This large variety of facts about hadrons has to be explained by the dynamical theory of strong interactions which is the subject of the next section.

1.2 Quantum Chromodynamics

Twenty-five years after its inception [5, 6, 7], Quantum Chromodynamics (QCD) is generally accepted as *the* theory of strong interactions. It describes the interaction of matter (quarks and antiquarks) with intermediate gauge bosons named gluons. Like the other theories of the fundamental interactions, QCD is a gauge theory; its symmetry group, $SU(3)$ is associated with three ‘color’ charges mediating the strong interaction. As the gauge group $SU(3)$ is non-Abelian, the (eight) gluons themselves carry color charge and therefore can interact with each other¹. Nevertheless, in a somewhat condensed notation, the QCD Hamiltonian appears very similar to the one of Quantum Electrodynamics (QED). The former is

$$\mathcal{H}_{\text{QCD}} = \frac{1}{2}(\mathbf{E}^2 + \mathbf{B}^2) + \psi^\dagger(\boldsymbol{\alpha} \cdot \mathbf{D} + \beta\mathcal{M})\psi , \quad (1.7)$$

where $\mathbf{D}$ denotes the covariant derivative (in the fundamental representation),

$$\mathbf{D} = i\nabla - g\mathbf{A} \quad (1.8)$$

¹This explains the name ‘gluon’ which pictorially expresses the idea that these particles act like glue when they bind among themselves (‘glueballs’) or force quarks to bind into hadrons.

and $\mathcal{M} = \text{diag}(m_u, m_d, m_s, \dots)$ the quark-flavor mass matrix. If its mass eigenvalues were all equal, $SU(N_F)$ would be an exact symmetry of QCD. However, as already stated, the only exact (local) symmetry is its gauge invariance under $SU(3)_C$.

Note that, in contrast to the Abelian gauge theory QED, the chromo-electric and -magnetic fields $\mathbf{E}$ and $\mathbf{B}$ as well as the gauge potentials $\mathbf{A}$ are non-commuting (color) matrices so that there are implicit summations over color indices in (1.7). The self-interaction of the gluon fields $\mathbf{A}$ becomes evident upon expanding the chromo-magnetic fields $\mathbf{B}$ in terms of the $\mathbf{A}$, which, however, will not be done here as it can be found in any textbook on QCD.

Like its older ‘relative’ QED, QCD is a renormalizable relativistic quantum field theory. Any infinities arising from the point-like (local) nature of the interaction can therefore be consistently absorbed into a redefinition of the physical parameters (masses, couplings). As a result, the strong coupling parameter α_s is not a constant but is running with the typical momentum scale of the physical process under consideration. The microscopic reason for this are vacuum polarization effects: quarks screen and therefore weaken the color charge (analogous to QED), whereas the self-interacting gluons *anti*-screen the color charge which is the dominating effect. Unlike in QED, therefore, the running coupling $\alpha_s(Q)$ of QCD is weak for high momentum transfer Q (small distances). This is the realm of ‘asymptotic freedom’ where perturbative methods work. For small momentum transfer Q (large distances), the coupling is large, perturbation theory breaks down, and one has to utilize nonperturbative methods. Typical and well-established values for α_s are [1]

$$\alpha_s(M_Z) = \alpha_s(91.2\text{GeV}) = 0.118 \pm 0.006, \quad (1.9)$$

$$\alpha_s(M_\tau) = \alpha_s(1.78\text{GeV}) = 0.33 \pm 0.03. \quad (1.10)$$

Thus, the nonperturbative domain is characterized by a maximum momentum scale of approximately 1 GeV. In some loose sense one can therefore speak of two relevant phases of QCD, the weak coupling phase or perturbative QCD, and the strong coupling phase (nonperturbative QCD).

Let us return to our first subject, the hadrons. In principle, it is quite clear, what a hadron is in QCD: it is an eigenstate of the QCD Hamiltonian,

$$H_{QCD}|\text{hadron}\rangle = M_H|\text{hadron}\rangle, \quad (1.11)$$

where M_H denotes the hadron mass². The question, of course, is, whether the ‘QCD Schrödinger equation’ (1.11) can be solved. If we consider a typical hadronic scale like, say, the nucleon radius, $R_N \simeq 1$ fm, the associated energy

² In writing down (1.11) we neglect any possible ‘fine splitting’ due to electromagnetic or weak interactions.

scale is $Q \simeq \hbar c/R_N \simeq 200$ MeV. This number tells us that we are in the low-energy regime which implies that the binding of quarks into hadrons is a non-perturbative phenomenon. In other words, a perturbative solution of the ‘QCD Schrödinger equation’ will make no sense in general.

There are two main routes out of this dilemma. Firstly, one can try to perform brute-force ab-initio calculations which involve sophisticated computer simulations on the largest machines available. Technically, one makes use of a space time discretization leading to a lattice formulation of gauge theories. This program has been developed over more than twenty years now, and has become the subject of entire textbooks [8].

Secondly, one can rely on a reputable tradition of physics, namely model building. There is an abundance of hadron models on the market, the most popular one being the constituent quark model and variants thereof. In this model, one starts with a non-relativistic phenomenological Hamiltonian of the form

$$H = H_0 + V_{\text{conf}} , \quad (1.12)$$

with an ad-hoc confining potential V_{conf} which typically is proportional to the inter-quark distance r (or r^2). The Hamiltonian describes the dynamics of two or three constituent quarks with masses $M_q \gtrsim 300$ MeV like in (1.6) which are treated as parameters. The main virtue of the model consists in its rather accurate reproduction of the hadron masses (‘spectroscopy’). An exhaustive compendium of results can be found in the recent reviews [9, 10, 11].

However, as any model, also the constituent quark model has its shortcomings. First of all, it works where it is not supposed to work. A hadron size of $R_H \simeq 1$ fm leads to an average constituent momentum of $\langle p \rangle \simeq 200$ MeV which is of the order of the constituent mass M_q . Therefore, hadrons are *relativistic* bound states and a non-relativistic treatment is not appropriate. Furthermore, the relation of the model with QCD is rather unclear. In QCD, as in any relativistic quantum field theory, one expects a bound state like e.g. the pion to be of the form

$$|\pi\rangle \sim \psi_2|q\bar{q}\rangle + \psi_3|q\bar{q}g\rangle + \psi_4|q\bar{q}q\bar{q}\rangle + \dots . \quad (1.13)$$

This means that hadrons are states containing an infinite number of quarks and gluons, which is consistent with the results of the DIS experiments where, with growing resolution Q^2 , an increasing number of partons is observed. This confirms that there are non-vanishing amplitudes $\psi_2, \psi_3 \dots$ to find two quarks, two quarks and a gluon ..., in general to find an arbitrary number of quarks and gluons in a hadron.

Another flaw of the model is that the constituent masses M_q do not coincide with the masses in the QCD Hamiltonian (1.7). This is qualitatively explained by the idea that the constituent quarks are *effective* quarks (quasi-particles) with effective masses originating from ‘medium effects’ due to the QCD vacuum. A

Table 1.2: *The presently observed quark flavors. Q/e is the electric charge in units of the electron charge. The (scale dependent!) quark masses are given for a scale of 1 GeV.*

flavor		Q/e	mass	
down	d	$-1/3$	10	MeV
up	u	$+2/3$	5	MeV
strange	s	$-1/3$	150	MeV
charm	c	$+2/3$	1.5	GeV
bottom	b	$-1/3$	5.1	GeV
top	t	$+2/3$	180	GeV

quantitative derivation of this mass generation within QCD is to a large extent still missing.

A more fundamental drawback of the constituent quark model is the following: it does not take into account all symmetries of QCD. This is particularly true for one prominent symmetry which is believed to govern most of the physics of light hadrons, namely chiral symmetry. This is our next topic.

1.3 Chiral Symmetry in QCD

If we have a look at Table 1.2, which provides a list of all quark flavors, we realize that there are large differences in the quark masses as they appear in the QCD Hamiltonian. In particular, there is a hierarchy,

$$\underbrace{m_u, m_d}_{\text{light quarks}} \ll m_s \ll \underbrace{m_c, m_b, m_t}_{\text{heavy quarks}} . \quad (1.14)$$

As the masses of heavy and light quarks are separated by the same scale as the perturbative and nonperturbative regime, one expects different physics associated with those two kinds of quarks. This expectation turns out to be true. The physics of heavy quarks is governed by a symmetry called ‘heavy quark symmetry’ leading to a very successful ‘heavy quark effective theory’ [12]. The physics of light quarks, on the other hand, is governed by chiral symmetry which we are now going to explain.

Let us write the QCD Hamiltonian in the following way,

$$H_{\text{QCD}} = H_\chi + \bar{\psi} \mathcal{M} \psi , \quad (1.15)$$

where we recall the mass matrix for the light flavors, $\mathcal{M} = \text{diag}(m_u, m_d, m_s)$. To a good approximation, one can set $\mathcal{M} = 0$. In this case, the QCD Hamiltonian H_χ

is invariant under the symmetry group $SU(3)_R \otimes SU(3)_L$, the chiral flavor group. Under the action of this group, the left and right handed quarks independently undergo a chiral rotation. Due to Noether's theorem, there are sixteen conserved quantities, eight vector charges and, more important for us, eight pseudo-scalars, the chiral charges Q_5^a satisfying

$$[Q_5^a, H_\chi] = 0 . \quad (1.16)$$

This states both that the chiral charges are conserved, and that H_χ is chirally invariant. Under parity, $Q_5^a \rightarrow -Q_5^a$. Now, if $|A\rangle$ is an eigenstate of H_χ , so is $Q_5^a|A\rangle$ with the same eigenvalue. Thus, one expects (nearly) degenerate parity doublets in nature, which, however, do not exist empirically. The only explanation for this phenomenon is that chiral symmetry is spontaneously broken. In contradistinction to the Hamiltonian, the QCD ground state (the vacuum) is *not* chirally invariant,

$$Q_5^a|0\rangle \neq 0 . \quad (1.17)$$

For this reason, there must exist a non-vanishing vacuum expectation value, the quark condensate,

$$\langle \bar{\psi}\psi \rangle = \langle \bar{\psi}_R\psi_L + \bar{\psi}_L\psi_R \rangle . \quad (1.18)$$

This condensate is not invariant (it mixes left and right) and therefore serves as an order parameter of the symmetry breaking. Note that in QCD the quark condensate is a renormalization scale dependent quantity. A recent estimate can be found in [13], with a numerical value,

$$\langle 0|\bar{\psi}\psi|0\rangle(1\text{ GeV}) \simeq (-229\text{ MeV})^3 . \quad (1.19)$$

In terms of the full quark propagator,

$$S(p) = \frac{\not{p} + M(p)}{p^2 - M^2(p)} , \quad (1.20)$$

where we have allowed for a momentum dependent (or 'running') mass $M(p)$, the quark condensate is given by

$$\langle 0|\bar{\psi}\psi|0\rangle = -i \int \frac{d^4p}{(2\pi)^4} \text{tr} S(p) = -4iN_C \int \frac{d^4p}{(2\pi)^4} \frac{M(p)}{p^2 - M^2(p)} . \quad (1.21)$$

We thus see that the involved Dirac trace yields a non-vanishing condensate only if the effective quark mass $M(p)$ is non-zero. This links the existence of a quark condensate to the mechanism of dynamical mass generation. In this way we have found another argument that the bare quarks appearing in the QCD Hamiltonian (1.15) indeed acquire constituent masses. Of course we are still lacking a microscopic mechanism for that.

Table 1.3: *Masses of the pseudo-scalar octet mesons (in MeV).*

meson	π^0	$\pi^\pm$	$K^0, \bar{K}^0$	$K^\pm$	η
mass	135	140	500	494	549

Goldstone's theorem [14] now states that for any symmetry generator which does not leave the vacuum invariant, there must exist a massless boson with the quantum number of this generator. This results in the prediction that in massless QCD one should have an octet of massless pseudo-scalar mesons. In reality one finds what is listed in Table 1.3.

The non-vanishing masses of these mesons are interpreted as stemming from the non-vanishing quark masses in the QCD Hamiltonian which break chiral symmetry explicitly. They can be consistently viewed as perturbations within an effective field theory dubbed 'chiral perturbation theory', which has become one of the standard tools to make quantitative predictions for low-energy strong-interaction physics [15].

It should be stressed that chiral symmetry has nothing to say about the mechanism of confinement which presumably is a totally different story. This is also reflected within a QCD based derivation of chiral symmetry breaking, the instanton model [16]. This model explains many facts of low-energy hadronic physics but is known not to yield confinement. It is therefore possible that confinement is not particularly relevant for the understanding of hadron structure [16].

1.4 Outline

This report is divided into two main parts, devoted to the foundations and applications of light-cone dynamics, respectively. In the first part, we begin with some general remarks on relativistic dynamics (Chapter 2). As a paradigm example we discuss the free relativistic particle which is the prototype of a reparametrization invariant system. We show that the choice of the time parameter is not unique as this corresponds to a gauge fixing, the purpose of which is to get rid of the reparametrization redundancies. By considering the stability subgroups of the Poincaré group, we find that there are essentially three reasonable choices of 'time' for a relativistic system, corresponding to Dirac's instant, point and front form, respectively.

The latter choice is the basis of light-cone dynamics which we analyze in detail in Chapter 3. We find some unique features like a maximal number of kinematical Poincaré generators, boost invariance of relative coordinates and light-cone wave

functions and separation of center-of-mass and relative motion. Many of these features are reminiscent of what is known from non-relativistic physics.

Chapter 4 is devoted to light-cone *field* quantization. We show how the Poincaré generators are defined in this case and utilize Schwinger's quantum action principle to derive the canonical commutators. The relation between equal-time commutators, the field equations and their solutions for different initial and/or boundary conditions is clarified. It is found that light-cone field theories, being of first order in the velocity, generally are constrained systems which require a special treatment. We use a method of phase space reduction due to Faddeev and Jackiw to take this into account. This leads us naturally to the construction of Fock space.

In Chapter 5 we discuss what might be viewed as the most spectacular feature of light-cone quantum field theory, the triviality of the vacuum. We show how this property comes about and illustrate the difference to standard instant-form quantization for some very simple model field theories.

We conclude the presentation of the foundations with Chapter 6, where we introduce the notion of light-cone wave functions. Their special properties make them useful tools for investigating hadronic physics. We show how light-cone wave functions of hadrons can be obtained by solving the light-cone Schrödinger equation.

The first application we will study in Part II is the phenomenon of spontaneous symmetry breaking for scalar field theory (Chapter 7), in a setting which can be viewed as a simplified variant of the Higgs mechanism. We analyze how a non-vanishing vacuum expectation value can arise in a trivial vacuum. We find that the vacuum properties are carried by the field mode carrying vanishing longitudinal momentum. This mode satisfies a complicated constraint equation which we solve both perturbatively and nonperturbatively.

As the methods of Chapter 7 do not work for fermionic fields, we have to consider alternatives which is the subject of Chapter 8. We use the particle spectrum of model field theories in 1+1 dimensions to reconstruct their vacuum properties. The spectrum is obtained by solving the light-cone Schrödinger equation. This is done here with unprecedented accuracy, which is made possible through the use of computer algebraic methods yielding analytic solutions even in high orders.

In Chapter 8 we finally make contact with phenomenology. We calculate the light-cone wave function of the pion within the Nambu and Jona-Lasinio model. This model is known to provide a good description of spontaneous chiral symmetry breaking, as it is governed by the same symmetry group as low-energy QCD. With the pion wave function at hand we predict a number of observables like the pion charge and core radius, the electromagnetic form factor and the r.m.s. transverse momentum. We conclude with a calculation of the pion distribution amplitude which is important for the study of exclusive processes in QCD.

Part I

Foundations

Chapter 2

Relativistic Particle Dynamics

The nature of elementary particles calls for a synthesis of relativity and quantum mechanics. The necessity of a quantum treatment is quite evident in view of the microscopic scales involved which are several orders of magnitude smaller than in atomic physics. These very scales, however, also require a relativistic formulation. As we have seen already, a typical hadronic scale of 1 fm, for instance, corresponds to momenta of the order of $p \sim \hbar c / 1 \text{ fm} \simeq 200 \text{ MeV}$. For particles with masses of the order of 1 GeV and below this implies sizable velocities of $0.2 c$ and larger.

It turns out that the task of combining the principles of quantum mechanics and relativity is not a straightforward one. One can neither simply extend ordinary quantum mechanics to include relativistic physics nor quantize relativistic mechanics using the ordinary correspondence rules. Nevertheless, Dirac and others have succeeded in formulating what is called “relativistic quantum mechanics”, which has become a subject of text books since [17, 18]. It should, however, be pointed out that this formulation, which is based on the concept of single-particle wave-functions and equations, is not really consistent. It does not correctly account for relativistic causality (retardation effects etc.) and the existence of anti-particles. As a result, one has to struggle with issues like the Klein paradox [17, 18], the definition of position operators [19] and the like.

The solution to these problems is provided by quantum *field* theory, with an inherently correct description of anti-particles that entails relativistic causality. In contrast to single-particle wave mechanics, quantum field theory is a (relativistic) many body formulation that necessarily involves (anti-)particle creation and annihilation. Physical particle states are typically a superposition of an infinite number of “bare” states, as any particle has a finite probability to emit or absorb other particles at any moment of time. With the advent of QCD, however, a conceptual difficulty concerning this many-particle picture has appeared and already been discussed in the introduction. At low energy or momentum transfer, hadrons, the bound states of QCD, are reasonably described as bound states of two or three constituent quarks and thus as *few-body* systems. It is rather unclear how such a constituent picture can arise in a quantum field theory such as QCD.

The physically motivated desire to describe hadrons as bound states of a small, fixed number of constituents is our rationale to go back and reanalyze the relation between Hamiltonian quantum mechanics and relativistic quantum field theory.

Quite generally, bound states are obtained by solving the Schrödinger equation,

$$i\hbar \frac{\partial}{\partial \tau} |\psi(\tau)\rangle = H |\psi(\tau)\rangle , \quad (2.1)$$

for *stationary states*,

$$|\psi(\tau)\rangle = e^{-iE\tau} |\psi(\tau = 0)\rangle . \quad (2.2)$$

This leads to the bound-state equation

$$H |\psi(\tau = 0)\rangle = E |\psi(\tau = 0)\rangle , \quad (2.3)$$

where E is the bound state energy. We would like to make this Hamiltonian formalism consistent with the requirements of relativity. It is, however, obvious from the outset that this procedure is not manifestly covariant as it singles out a time τ (and an energy E , respectively). Furthermore, it is not even clear what the time τ really is as it does not have an invariant meaning.

2.1 The Free Relativistic Particle

To see what is involved it is sufficient to consider the classical dynamics of a free relativistic particle. We want to find the associated canonical formulation as a basis for subsequent quantization. We will proceed by analogy with the treatment of classical free strings which is described in a number of textbooks [20, 21]. Accordingly, the relativistic point particle may be viewed as an infinitely short string.

We recall that the action for a relativistic particle is essentially given by the arc length of its trajectory

$$S = -ms_{12} \equiv -m \int_1^2 ds . \quad (2.4)$$

This action¹ is a Lorentz scalar as

$$ds = \sqrt{g_{\mu\nu} x^\mu x^\nu} \quad (2.5)$$

is the (infinitesimal) invariant distance. We can rewrite the action (2.4) as

$$S = -m \int_1^2 ds \sqrt{\dot{x}_\mu \dot{x}^\mu} \equiv \int_1^2 ds L(s) , \quad (2.6)$$

¹We work in natural units, $\hbar = c = 1$.

in order to introduce a Lagrangian $L(s)$ and the four velocity $\dot{x}^\mu \equiv dx^\mu/ds$. The latter obeys

$$\dot{x}^2 \equiv \dot{x}_\mu \dot{x}^\mu = 1 , \quad (2.7)$$

as the arc length provides a natural parametrization. Thus, $\dot{x}^\mu$ is a time-like vector, and we assume in addition that it points into the future, $\dot{x}^0 > 0$. In this way we guarantee relativistic causality ensuring that a real particle passing through a point A will always propagate into the future light cone based at A .

We proceed with the canonical formalism by calculating the canonical momenta as

$$p^\mu = -\frac{\partial L}{\partial \dot{x}_\mu} = m \dot{x}^\mu . \quad (2.8)$$

These are not independent, as can be seen by calculating the square using (2.7),

$$p^2 = m^2 \dot{x}^2 = m^2 , \quad (2.9)$$

which, of course, is the usual mass-shell constraint. This constraint indicates that the Lagrangian $L(s)$ defined in (2.6) is *singular*, which means that its Hessian $W^{\mu\nu}$ with respect to the velocities,

$$W^{\mu\nu} = \frac{\partial^2 L}{\partial \dot{x}_\mu \partial \dot{x}_\nu} = -\frac{m}{\sqrt{\dot{x}^2}} \left(g^{\mu\nu} - \frac{\dot{x}^\mu \dot{x}^\nu}{\dot{x}^2} \right) = -m (g^{\mu\nu} - \dot{x}^\mu \dot{x}^\nu) , \quad (2.10)$$

is degenerate. It has a zero mode given by the velocity itself,

$$W^{\mu\nu} \dot{x}_\nu = 0 . \quad (2.11)$$

The Lagrangian being singular implies that the velocities cannot be uniquely expressed in terms of the canonical momenta. This, however, is not obvious from (2.8), as we can easily solve for the velocities,

$$\dot{x}^\mu = p^\mu / m . \quad (2.12)$$

But if one now calculates the canonical Hamiltonian,

$$H_c = -p_\mu \dot{x}^\mu - L = -m \dot{x}^2 + m \dot{x}^2 = 0 , \quad (2.13)$$

one finds that it is vanishing! It therefore seems that we do not have a generator for the time evolution of our dynamical system. In the following, we will analyze the reasons for this peculiar finding.

First of all we note that the Lagrangian is homogeneous of first degree in the velocity,

$$L(\alpha \dot{x}^\mu) = \alpha L(\dot{x}^\mu) . \quad (2.14)$$

Thus, under a reparametrization of the world-line,

$$s \mapsto s' , \quad x^\mu(s) \mapsto x^\mu(s'(s)) , \quad (2.15)$$

where the mapping $s \mapsto s'$ is one-to-one with $ds'/ds > 0$ (orientation conserving), the Lagrangian changes according to

$$L(dx^\mu/ds) = L\left((dx^\mu/ds')(ds'/ds)\right) = (ds'/ds)L(dx^\mu/ds') . \quad (2.16)$$

This is sufficient to guarantee that the action is invariant under (2.15), that is, *reparametrization invariant*,

$$S = \int_{s_1}^{s_2} ds L(dx^\mu/ds) = \int_{s'_1}^{s'_2} ds' \frac{ds}{ds'} \frac{ds'}{ds} L(dx^\mu/ds') \equiv S' , \quad (2.17)$$

if the endpoints remain unchanged, $s_{1,2} = s'_{1,2}$. On the other hand, L is homogeneous of the first degree if and only if Euler's formula holds, namely

$$L = \frac{\partial L}{\partial \dot{x}^\mu} \dot{x}^\mu = -p_\mu \dot{x}^\mu . \quad (2.18)$$

This is exactly the statement (2.13), the vanishing of the Hamiltonian. Furthermore, if we differentiate (2.18) with respect to $\dot{x}^\mu$, we recover (2.11) expressing the singular nature of the Lagrangian. Summarizing, we have found the general result [22, 23] that if a Lagrangian is homogeneous of degree one in the velocities, the action is reparametrization invariant, and the Hamiltonian vanishes. In this case, the momenta are homogeneous of degree zero, which renders the Lagrangian singular.

In order to properly incorporate the reparametrization invariance we chose an arbitrary parametrization, $\tau \mapsto x^\mu(\tau)$, so that the infinitesimal arc length becomes

$$ds^2 = \left(\frac{ds}{d\tau}\right)^2 d\tau^2 \equiv h(\tau) d\tau^2 , \quad (2.19)$$

introducing a world-line metric (or einbein) $h(\tau)$. In what follows we let the over-dot denote the derivative with respect to τ , e.g. $\dot{f} = df/d\tau$. From (2.19) we get that

$$\dot{x}^2 = \left(\frac{ds}{d\tau}\right)^2 = h(\tau) , \quad (2.20)$$

so that, according to (2.7), $h(\tau) = 1$ corresponds to $d\tau = ds$. Therefore, the world-line metric h provides a 'length' scale for the four-velocity. We write the action as

$$S = -m \int_1^2 d\tau \sqrt{\dot{x}^2} = -m \int_1^2 d\tau \sqrt{h(\tau)} \equiv \int_1^2 d\tau L(\tau) . \quad (2.21)$$

The canonical momenta are

$$\pi^\mu \equiv -\frac{\partial L}{\partial \dot{x}_\mu} = m \frac{\dot{x}^\mu}{\sqrt{\dot{x}^2}} = m \frac{\dot{x}^\mu}{\sqrt{h(\tau)}} = \frac{m}{\sqrt{h(\tau)}} \frac{dx^\mu}{ds} \frac{ds}{d\tau} = m \frac{dx^\mu}{ds} \equiv p^\mu . \quad (2.22)$$

We note that they are independent of the world-line metric h , so that, upon squaring (2.22) one again finds the mass shell constraint $\pi^2 = p^2 = m^2$. However, if we want to express the velocities in terms of the momenta, we get

$$\dot{x}^\mu = \sqrt{h} p^\mu / m . \quad (2.23)$$

Therefore, it is exactly the world-line metric h that expresses the arbitrariness involved when solving for the velocities. Unlike for the momenta, the scale for the velocities is not fixed, unless one chooses a particular world-line metric as in (2.7).

By the same arguments as before, the canonical Hamiltonian still vanishes. The Lagrangian one-form thus coincides with the canonical one-form,

$$L(\tau)d\tau = -p_\mu dx^\mu . \quad (2.24)$$

From the latter we read off the non-vanishing Poisson brackets,

$$\{x^\mu, p^\nu\} = -g^{\mu\nu} , \quad (2.25)$$

which will be needed in a moment. To proceed we utilize the Dirac-Bergmann algorithm for singular systems [24, 25, 26, 27] (for monographs, see [22, 23, 28, 29]) by adding the constraint

$$\theta \equiv p^2 - m^2 \quad (2.26)$$

to the (vanishing) canonical Hamiltonian via a Lagrange multiplier λ . This yields the primary Hamiltonian

$$H_p = \lambda(p^2 - m^2) . \quad (2.27)$$

The constraint (2.26) is the only one arising and has vanishing Poisson bracket with itself whereupon it is called *first class*. Technically, this implies that demanding consistency, i.e. conservation of the constraint in time τ , does not lead to an equation determining the Lagrange multiplier. Physically, this means that the time development of the system is not completely determined. Such a behavior is typical for systems with a gauge or redundancy symmetry, an example of which is the reparametrization invariance discussed above. The redundancy consists in the fact that a single trajectory (world-line) can be described by an infinite number of different parametrizations. The physical objects, the trajectories, are therefore equivalence classes obtained by identifying all reparametrizations. An infinitesimal reparametrization is given by

$$\tau'(\tau) = \tau + \delta\tau(\tau) , \quad (2.28)$$

$$\delta x^\mu(\tau) = x^\mu(\tau + \delta\tau) - x^\mu(\tau) = \dot{x}^\mu \delta\tau . \quad (2.29)$$

In systems with redundancy symmetries, these symmetries are generated by the first class constraints. A familiar example is provided by gauge fields (like in electromagnetism), where the first class constraints are represented by Gauß's

law, the generator of (time-independent) gauge transformations. For the case at hand, we find using (2.25),

$$\delta x^\mu = \{x^\mu, \delta G\} \equiv \{x^\mu, \theta \delta \epsilon\} = -2p^\mu \delta \epsilon = -2m \frac{\dot{x}^\mu}{\sqrt{h}} \delta \epsilon. \quad (2.30)$$

Identifying $\delta \tau \equiv (2m/\sqrt{h})\delta \epsilon$ to account for the different dimensionalities, (2.29) and (2.30) coincide. Thus, the reparametrization (2.29) is indeed generated by the first-class constraint θ of (2.26). The Hamiltonian equations of motion become

$$\dot{x}^\mu = \{x^\mu, H_p\} = -2\lambda p^\mu, \quad (2.31)$$

$$\dot{p}^\mu = \{p^\mu, H_p\} = 0. \quad (2.32)$$

It is important to note at this point that the equation for $\dot{x}^\mu$ contains the Lagrange multiplier λ which is undetermined so far. This makes the whole dynamics of the system undetermined. If we compare (2.31) with (2.23), we find that the arbitrariness is again encoded in the world-line metric as the multiplier is

$$\lambda = -\sqrt{h}/2m. \quad (2.33)$$

To overcome the problem of determining λ or h , one has to *fix a gauge* by choosing an auxiliary condition,

$$\chi(x; \tau) = 0. \quad (2.34)$$

Demanding consistency, i.e. conservation of the gauge fixing condition in time,

$$\dot{\chi} = \frac{\partial \chi}{\partial \tau} + \{\chi, H_p\} = \frac{\partial \chi}{\partial \tau} + \lambda \{\chi, \theta\} = \frac{\partial \chi}{\partial \tau} - 2\lambda p^\mu \frac{\partial \chi}{\partial x^\mu} = 0, \quad (2.35)$$

we obtain the desired equation determining the Lagrange multiplier λ ,

$$\lambda = -\{\chi, \theta\}^{-1} \frac{\partial \chi}{\partial \tau} = \frac{1}{2(p \cdot \partial)_\chi} \frac{\partial \chi}{\partial \tau}. \quad (2.36)$$

We see that it is crucial for the determination of λ that the gauge fixing χ depends *explicitly* on the time parameter τ and on at least one of the x^μ . This is necessary for rendering the Poisson bracket $\{\chi, \theta\}$ nonvanishing. The analogue of this condition in the gauge theory context is well known: it corresponds to having a nonvanishing Faddeev-Popov matrix. For the relativistic particle, one can say that the gauge condition amounts to defining a time parameter τ in terms of the coordinates x^μ . This suggests that χ will in general be of the form

$$\chi(x; \tau) = \tau - F(x), \quad (2.37)$$

stating that τ is some function F of the coordinates, $\tau = F(x)$. In this case, (2.36) becomes

$$\lambda = -\frac{1}{2(p \cdot \partial)F(x)}. \quad (2.38)$$

The quantity $(p \cdot \partial)F(x)$ is nothing but the projection of the four-momentum p^μ onto the normal $N^\mu \equiv \partial^\mu F$ of the hypersurface $\tau = \text{const.}$ The knowledge of the Lagrange multiplier (or the einbein h) makes the dynamics uniquely determined, as (2.31) leads to

$$\dot{x}^\mu = -2\lambda p^\mu = \frac{p^\mu}{N \cdot p} . \quad (2.39)$$

Squaring this expression or using (2.33), the world-line metric is found to be

$$h(\tau) = 4m^2 \lambda^2 = \frac{m^2}{(N \cdot p)^2} = \frac{p^2}{(N \cdot p)^2} . \quad (2.40)$$

It thus measures the normal component of the momentum (squared) compared to $p^2 = m^2$. The most common choice for τ is the Galileian time t of, say, the observer rest frame,

$$\chi = x^0 - \tau \equiv t - \tau = 0 . \quad (2.41)$$

This gauge choice corresponds to a particular choice for the world-line metric h , given by

$$ds^2 = dt^2 - dx^i dx^i = (1 - v^2) dt^2 = \frac{1}{\gamma^2} dt^2 \equiv h(t) dt^2 , \quad (2.42)$$

with $v^i = dx^i/dt$. In this case, the Lagrange multiplier as derived from (2.36) becomes

$$\lambda = -\frac{1}{N \cdot p} = -\frac{1}{2p^0} . \quad (2.43)$$

From (2.31) and (2.39) we get

$$\dot{x}^\mu = -2\lambda p^\mu = p^\mu / p^0 , \quad (2.44)$$

This yields the four-velocity squared

$$\dot{x}^2 = \frac{p^2}{p_0^2} = \frac{m^2}{p_0^2} = \frac{1}{\gamma^2} \equiv h(t) , \quad (2.45)$$

in accordance with the result (2.42) for the world-line metric. The Hamiltonian generating the t -evolution is found by solving the mass-shell constraint θ for $p^0 = N \cdot p$,

$$H \equiv p^0 = \sqrt{\mathbf{p}^2 + m^2} . \quad (2.46)$$

It generates the equations of motion

$$\dot{x}^i = \{x^i, H\} = \frac{p^i}{p^0} , \quad (2.47)$$

$$\dot{p}^i = \{p^i, H\} = 0 , \quad (2.48)$$

where the first one is consistent with (2.39) so that the second one coincides with the Euler-Lagrange equation following from (2.21). The Poisson brackets above are interpreted as being taken at *equal time* t .

Altogether, the gauge-fixed dynamics is well-determined as can be seen from the (unique) solution

$$x^i(t) = \frac{p^i}{p^0}t + x^i(0) = v^i t + x^i(0) , \quad (2.49)$$

which corresponds to linear motion with constant velocity.

We have thus seen that it is the choice of time which makes the dynamics unambiguous. This problem and its solution are actually typical for reparametrization invariant systems, the most important example probably being general relativity. There, the problem of choosing a natural time variable is the main obstacle for developing a quantum theory of gravity (for a thorough discussion, see [30]).

In the remainder of this section we will perform a general analysis of the possible choices of gauge fixings, that is, of different time parameters τ . The choice of a particular time variable corresponds to a foliation of Minkowski space into hypersurfaces of equal time, $\tau = \text{const}$, which in general are three-dimensional objects, and the time direction ‘orthogonal’ to them. The time development thus continuously evolves the hypersurface $\Sigma_0 : \tau = \tau_0$ into $\Sigma_1 : \tau = \tau_1 > \tau_0$. Put differently, initial conditions provided on Σ_0 together with the dynamical equations (being differential equations in τ) determine the state of the dynamical system on Σ_1 .

The gauge fixing (2.37) expresses the time parameter in terms of the coordinates x^μ . These are orthogonal coordinates which are essentially tailored to fit the common choice of time, $\tau = t$. On the other hand, we have explicitly seen within this particular example, that the gauge (or time) choice corresponds to an associated choice of the world-line metric. It is the latter, on which we now want to concentrate. Consider some arbitrary coordinates $\xi^\alpha = \xi^\alpha(x)$, which may be curvilinear. We imagine that the ξ^i , $i = 1, 2, 3$, parametrize the three-dimensional hypersurface Σ , so that the remaining one, ξ^0 , represents the time variable, i.e. $\tau = \xi^0$. Then we have

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = g_{\mu\nu} \frac{\partial x^\mu}{\partial \xi^\alpha} \frac{\partial x^\nu}{\partial \xi^\beta} d\xi^\alpha d\xi^\beta \equiv h_{\alpha\beta}(\xi) d\xi^\alpha d\xi^\beta . \quad (2.50)$$

Introducing a vierbein $e^\mu_\alpha(\xi)$ the metric $h_{\alpha\beta}(\xi)$ is alternatively given by

$$h_{\alpha\beta}(\xi) = g_{\mu\nu} e^\mu_\alpha(\xi) e^\nu_\beta(\xi) . \quad (2.51)$$

This transformation is well known from general relativity, where it corresponds to the transformation from a local inertial frame described by the flat metric $g_{\mu\nu}$ to a non-inertial frame with coordinate dependent metric $h_{\alpha\beta}(\xi)$. For our purposes we write this metric in a (3+1)-notation as follows,

$$h_{\alpha\beta} = \begin{pmatrix} h_{00} & h_{0i} \\ h_{i0} & h_{ij} \end{pmatrix} . \quad (2.52)$$

Of particular interest is the component h_{00} , which explicitly reads

$$h_{00} = g_{\mu\nu} \frac{\partial x^\mu}{\partial \xi^0} \frac{\partial x^\nu}{\partial \xi^0} = g_{\mu\nu} e^\mu{}_0 e^\nu{}_0 \equiv n^2, \quad (2.53)$$

where we have defined the unit vector in ξ^0 -direction

$$n^\mu = \frac{\partial x^\mu}{\partial \xi^0} = e^\mu{}_0. \quad (2.54)$$

It is related to the normal vector N^μ via

$$n \cdot N = e^\mu{}_0 e_\mu{}^0 = \frac{\partial \xi^0}{\partial x^\mu} \frac{\partial x^\mu}{\partial \xi^0} = 1. \quad (2.55)$$

The h_{ij} are the metric components associated with the hypersurface. The invariant distance element thus becomes (setting $h_{0i} \equiv h_i$),

$$\begin{aligned} ds^2 &= h_{00} d\xi^0 d\xi^0 + 2h_{0i} d\xi^0 d\xi^i + h_{ij} d\xi^i d\xi^j, \\ &= \left(n^2 + 2h_i \frac{d\xi^i}{d\tau} + h_{ij} \frac{d\xi^i}{d\tau} \frac{d\xi^j}{d\tau} \right) d\tau^2, \end{aligned} \quad (2.56)$$

where, in the second step, we have used that $\xi^0 = \tau$. The quantities $w^i \equiv d\xi^i/d\tau$ are the velocities expressed in the new coordinates, so that the world-line metric can be written as

$$h(\tau) = n^2 + 2h_i w^i + h_{ij} w^i w^j. \quad (2.57)$$

For $\tau \equiv \xi^0 \equiv t$ we find $n = N = (1, \mathbf{0})$, $N^2 = 1$, $h_i = 0$, $w^i = v^i$ and $h_{ij} = -\delta_{ij}$ leading to the world-line metric $h(t) = 1 - v^2$ of (2.42, 2.45). The dynamics of the ξ^i is easily found as follows,

$$\dot{\xi}^i = \partial_\mu \xi^i \dot{x}^\mu = \partial_\mu \xi^i \frac{p^\mu}{N \cdot p} = \frac{p^\mu e_\mu{}^i}{N \cdot p} \equiv \frac{\pi^i}{N \cdot p}, \quad (2.58)$$

where we have introduced the momenta π^i canonically conjugate to ξ^i , satisfying

$$\{\xi^i, \pi^j\} = h^{ij}(\xi). \quad (2.59)$$

2.2 Dirac's Forms of Relativistic Dynamics

At this point one needs a satisfactory criterion in order to decide which choices for the time variable (or the world-line metric h) are sensible. To this end it is not sufficient to consider only the τ -development and the associated generator of time translations (i.e. the Hamiltonian). Instead, one has to refer to the full Poincaré group to be able to guarantee full relativistic invariance. The generators

of the Poincaré group are the four momenta P^μ and the six operators $M^{\mu\nu}$ which combine the angular momenta and boosts according to

$$L^i = \frac{1}{2}\epsilon^{ijk}M^{jk}, \quad (2.60)$$

$$K^i = M^{0i}, \quad (2.61)$$

with $i, j, k = 1, 2, 3$. These generators are elements of the Poincaré algebra which is defined by the Poisson bracket relations,

$$\begin{aligned} \{P^\mu, P^\nu\} &= 0, \\ \{M^{\mu\nu}, P^\rho\} &= g^{\nu\rho}P^\mu - g^{\mu\rho}P^\nu, \\ \{M^{\mu\nu}, M^{\rho\sigma}\} &= g^{\mu\sigma}M^{\nu\rho} - g^{\mu\rho}M^{\nu\sigma} - g^{\nu\sigma}M^{\mu\rho} + g^{\nu\rho}M^{\mu\sigma}. \end{aligned} \quad (2.62)$$

It is well known that the momenta P^μ generate space-time translations and the $M^{\mu\nu}$ rotations and Lorentz boosts, cf. (2.60, 2.61). In the following we will only consider proper and orthochronous Lorentz transformations, i.e. we exclude space and time reflections.

Any Poincaré invariant dynamical theory describing e.g. the interaction of particles should provide a particular realization of the Poincaré algebra. For this purpose, the Poincaré generators are constructed out of the fundamental dynamical variables like positions, momenta, spins etc. An elementary realization of (2.62) is given as follows. Choose the space-time point x^μ and its conjugate momentum p^μ as canonical variables, i.e. adopt (2.25),

$$\{x^\mu, p^\nu\} = -g^{\mu\nu}. \quad (2.63)$$

The Poincaré generators are then found to be

$$P^\mu = p^\mu, \quad M^{\mu\nu} = x^\mu p^\nu - x^\nu p^\mu \quad (2.64)$$

as is easily confirmed by checking (2.62) using (2.63). An infinitesimal Poincaré transformation is thus generated by

$$\delta G = -\frac{1}{2}\delta\omega_{\mu\nu}M^{\mu\nu} + \delta a_\mu P^\mu, \quad (2.65)$$

in the following way,

$$\delta x^\mu = \{x^\mu, \delta G\} = \delta\omega^{\mu\nu}x_\nu + \delta a^\mu, \quad \delta\omega^{\mu\nu} = -\delta\omega^{\nu\mu}. \quad (2.66)$$

The action of the Poincaré group on some function $F(x)$ is thus given by

$$\delta F = \{F, \delta G\} = \partial^\mu F \delta a_\mu - \frac{1}{2}(x^\mu \partial^\nu - x^\nu \partial^\mu)F \delta\omega_{\mu\nu}. \quad (2.67)$$

Though the realization (2.64) is covariant, it has several shortcomings. It does not describe any interaction; for several particles the generators are simply the

Table 2.1: All possible choices of hypersurfaces Σ : $\tau = \text{const}$ with transitive action of the stability group G_Σ . d denotes the dimension of G_Σ , that is, the number of kinematical Poincaré generators, $\mathbf{x}_\perp \equiv (x^1, x^2)$.

name	Σ	τ	d
instant	$x^0 = 0$	t	6
light front	$x^0 + x^3 = 0$	$t + x^3/c = 0$	7
hyperboloid	$x_0^2 - \mathbf{x}^2 = a^2 > 0, x^0 > 0$	$(t^2 - \mathbf{x}^2/c^2 - a^2/c^2)^{1/2}$	6
hyperboloid	$x_0^2 - \mathbf{x}_\perp^2 = a^2 > 0, x^0 > 0$	$(t^2 - \mathbf{x}_\perp^2/c^2 - a^2/c^2)^{1/2}$	4
hyperboloid	$x_0^2 - x_1^2 = a^2 > 0, x^0 > 0$	$(t^2 - x_1^2/c^2 - a^2/c^2)^{1/2}$	4

sum of the single particle generators. This point, however, is of minor importance to us, and will only be touched upon at the end of the next chapter. The solution of the problem, as already mentioned in the introduction to this chapter, is the framework of local quantum field theory. More importantly, the representation (2.64) does not take into account the mass-shell constraint, $p^2 = m^2$, which we already know to guarantee relativistic causality as it generates the dynamics.

To remedy the situation we proceed as before by choosing a ‘time’ variable τ , i.e. a foliation of space time into essentially space-like hypersurfaces Σ with time-like or light-like normals. Σ should be chosen in such a way that it intersects all possible world-lines once and only once. Apart from this necessary consistency with causality this foliation appears quite arbitrary. However, given a particular foliation one can ask the question which of the Poincaré generators will leave the hypersurface Σ invariant. The set of all such generators defines a subgroup of the Poincaré group called the stability group G_Σ of Σ . The associated generators are called *kinematical*, the others *dynamical*. The latter map Σ onto another hypersurface Σ' and thus involve the development in τ . One thus expects that the dynamical generators will depend on the Hamiltonian (and, therefore, the interaction) which, by definition, is a dynamical quantity.

It is clear, however, that the stability group corresponding to a particular foliation will be empty if the associated hypersurface looks very irregular and thus does not have a high degree of symmetry. One therefore demands in addition that the stability group acts transitively on Σ : any two points on Σ can be connected by a transformation from G_Σ . With this additional requirement there are exactly five inequivalent classes of hypersurfaces [31] which are listed in Table 2.1.

The first three choices have been found by Dirac in his seminal 1949 paper [32] on ‘forms of relativistic dynamics’. He called the associated forms the ‘instant’, ‘front’ and ‘point’ forms, respectively. These are the most important choices as the other two forms have a rather small stability group and thus are not very

useful. We have only listed them for the sake of completeness.

It is important to note that for all forms one has $\lim_{c \rightarrow \infty} \tau = t$, which means that in the non-relativistic case there is only one possible foliation leading to the absolute Galileian time t . This is consistent with the fact that there is no limiting velocity in this case implying that particle trajectories can have arbitrary slope (tangent vector). Therefore, the hypersurface $\Sigma_{nr} : t = \text{const}$ is the only one intersecting all possible world-lines.

To decide which of the Poincaré generators are kinematical we use the general formula (2.67) describing their action. Imagine that Σ is given in the form $\Sigma : \tau = F(x)$ as in Table 2.1. If P^μ or $M^{\mu\nu}$ are kinematical for some μ or ν , then, for these particular superscripts, we must have

$$\partial^\mu F = 0, \quad (x^\mu \partial^\nu - x^\nu \partial^\mu) F = 0, \quad (2.68)$$

saying that the gradient and the ‘angular derivative’ of F have to vanish. Let us now discuss the different forms in more detail along these lines.

The Instant Form

The choice of Galileian time $\tau = t$ is of course the most common one also in the relativistic case, and we have discussed it briefly in the preceding subsection. To complete this discussion, we construct the associated representation of the Poincaré generators on $\Sigma : t = 0$. The idea is to explicitly saturate the constraint $p^2 = m^2$ by eliminating the variable conjugate to $\tau = t$, i.e. $p^0 = N \cdot p = \sqrt{\mathbf{p}^2 + m^2}$, and setting $x^0 = 0$ in (2.64). Alternatively, one can follow Dirac [32] and add the constraint to *any* of the Poincaré generators,

$$P^\mu = p^\mu + \lambda^\mu (p^2 - m^2), \quad (2.69)$$

$$M^{\mu\nu} = x^\mu p^\nu - x^\nu p^\mu + \lambda^{\mu\nu} (p^2 - m^2). \quad (2.70)$$

Now we demand that the right-hand-sides should be independent of p^0 . Differentiating the right-hand-side of (2.69) with respect to p^0 one finds

$$1 + 2\lambda^0 p^0 = 0, \quad (2.71)$$

which is exactly the consistency condition (2.35) for the case at hand. Thus λ^0 is determined as $\lambda^0 = -1/2p^0$ which coincides with (2.43) upon identifying $\lambda^0 \equiv \lambda$. The multipliers λ^i turn out to be zero. The same procedure applied to the $M^{\mu\nu}$ of (2.70) yields as the nonvanishing Lagrange multipliers the λ^{0i} with

$$\lambda^{0i} = -\frac{x^i}{2p^0}. \quad (2.72)$$

Altogether we obtain the following (3+1)-representation of the Poincaré generators,

$$\begin{aligned} P^i &= p^i, & M^{ij} &= x^i p^j - x^j p^i, \\ P^0 &= \omega_p, & M^{i0} &= x^i \omega_p, & \omega_p &\equiv (p^i p^i + m^2)^{1/2}. \end{aligned} \quad (2.73)$$

This result is as expected: Compared to (2.64), p^0 has been replaced by ω_p , and x^0 has been set to zero. It should, however, be pointed out that for non-Cartesian coordinates the construction of the Poincaré generators is less straightforward, as will be seen in a moment.

As a general rule we note that only the generators containing p^0 in the representation (2.69, 2.70) lead to non-vanishing Lagrange multipliers. These are just the dynamical generators which are, for the instant form, given by the ordinary Hamiltonian P^0 and the boost generators M^{i0} . In agreement with (2.68), one has

$$\partial^i F = (x^i \partial^j - x^j \partial^i) F = 0 , \quad (2.74)$$

so that Σ is both translationally and rotationally invariant confirming that the dimension of its stability group is six (cf. Table 2.1). On the other hand,

$$\partial^0 F = 1 \neq 0 , \quad (2.75)$$

$$(x^0 \partial^i - x^i \partial^0) F = -x^i \neq 0 , \quad (2.76)$$

from which we read off that, apart from the Hamiltonian, also the boosts are dynamical, i.e., Σ is not boost invariant. The latter fact is, of course, well known because the boosts mix space and time. Under a boost along the $\mathbf{n}$ -direction with velocity $\mathbf{v}$ ($\mathbf{n} = \mathbf{v}/v$), t transforms as

$$t \rightarrow t' = t \cosh \omega + (\mathbf{n} \cdot \mathbf{x}) \sinh \omega , \quad (2.77)$$

where ω is the rapidity, defined through $\tanh \omega = v$. From (2.77) it is evident that the hypersurface $\Sigma : t = 0$ is not boost invariant.

In obtaining the representation (2.73), we make the Poincaré algebra compatible with the instant-form constraint $x^0 = 0$. An elementary calculation, using $\{x^i, p^j\} = \delta^{ij}$, indeed shows that the generators (2.73) really obey the bracket relations (2.62). We have already seen that the Hamiltonian $P^0 = \omega_p$ generates the correct dynamics.

The Point Form

As might have been expected, the situation is somewhat more complicated for the point form. The hypersurfaces of equal $\tau = (x^2 - a^2)^{1/2}$ are hyperboloids implying that curvilinear coordinates are involved. For the following discussion we chose $a = 0$ for simplicity. Note, however, that in this case the hypersurface $\Sigma : \tau = 0$ is the light-cone which is not transitive and has a smaller stability group of dimension four. We therefore assume $\tau > 0$ in what follows.

It is natural to introduce hyperbolic coordinates defined via

$$\begin{aligned} x^0 &= \tau \cosh \omega , \\ x^1 &= \tau \sinh \omega \sin \theta \cos \phi , \\ x^2 &= \tau \sinh \omega \sin \theta \sin \phi , \\ x^3 &= \tau \sinh \omega \cos \theta . \end{aligned} \quad (2.78)$$

The coordinates can thus be written in the compact form

$$x^\mu = \tau n^\mu = \tau (\cosh \omega, \mathbf{e}_r \sinh \omega) , \quad (2.79)$$

where $\mathbf{e}_r$ is the radial unit vector in $\mathbb{R}^3$. The new coordinates are a straightforward generalization of the (three-dimensional) polar coordinates. Using the general formula (2.50), they lead to a world-line metric

$$h(\tau) = 1 - \tau^2 \left(\frac{d\omega}{d\tau} \right)^2 - \tau^2 \sinh^2 \omega \left(\frac{d\theta}{d\tau} \right)^2 - \tau^2 \sinh^2 \omega \sin^2 \theta \left(\frac{d\phi}{d\tau} \right)^2 . \quad (2.80)$$

The normal vector on Σ is $N^\mu = n^\mu = x^\mu / \tau$ which can be interpreted as a four-dimensional radial vector. The Lagrange multiplier is therefore given by

$$\lambda = -\frac{1}{2N \cdot p} = -\frac{\tau}{2p \cdot x} , \quad (2.81)$$

so that the equation of motion for the x^μ becomes

$$\dot{x}^\mu = \frac{\tau}{p \cdot x} p^\mu . \quad (2.82)$$

The Hamiltonian, that is, the generator of τ -development, is

$$H = N \cdot p = x \cdot p / \tau . \quad (2.83)$$

Obviously, we cannot set τ equal to zero. Note that this Hamiltonian does *not* belong to the Poincaré generators as it is essentially the dilatation operator generating scale transformations [33]. Its explicit τ -dependence indicates that the dynamical system is not scale-invariant which is obvious as we consider a massive particle.

Let us analyze the Poincaré generators. To decide which of them are dynamical, we calculate with $\tau = F(x) = \sqrt{x^2}$,

$$\partial^\mu F = N^\mu = \frac{x^\mu}{\tau} \neq 0 , \quad (2.84)$$

$$(x^\mu \partial^\nu - x^\nu \partial^\mu) F = \frac{1}{\tau} (x^\mu x^\nu - x^\nu x^\mu) F = 0 . \quad (2.85)$$

Not unexpectedly, all momenta are dynamical, but Lorentz transformations are kinematical. This is obvious from the fact that τ is a Lorentz scalar rendering Σ boost and rotation invariant. The stability group G_Σ thus has again dimension six.

Let us finally mention that the point form, though already introduced by Dirac, has only rarely been used in physical applications [34, 33, 35, 36]. The reason is, of course, the curvilinear nature of Σ which particularly hampers the quantization.

We are left with the front form which is at the basis of the present work. We reserve the next chapter for its investigation.

Chapter 3

The Front Form

3.1 Generalities

For an arbitrary four-vector a we perform the following transformation to *light-cone* coordinates,

$$(a^0, a^1, a^2, a^3) \mapsto (a^+, a^1, a^2, a^-) , \quad (3.1)$$

where we have defined

$$a^+ = a^0 + a^3 , \quad a^- = a^0 - a^3 . \quad (3.2)$$

We also introduce the transverse vector part of a as

$$\mathbf{a}_\perp = (a^1, a^2) . \quad (3.3)$$

The metric tensor becomes

$$g_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & 1/2 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 1/2 & 0 & 0 & 0 \end{pmatrix} \quad (3.4)$$

The entries $1/2$ imply a slightly unusual scalar product


$$a \cdot b = g_{\mu\nu} a^\mu b^\nu = \frac{1}{2} a^+ b^- + \frac{1}{2} a^- b^+ - a^i b^i , \quad i = 1, 2 . \quad (3.5)$$

According to Table 2.1, the front form is defined by choosing the hypersurface $\Sigma : x^+ = 0$, which is a plane tangent to the light-cone. It can equivalently be viewed as the wave front of a plane light wave traveling towards the positive z -direction. Therefore, Σ is also called a *light-front*. It corresponds to a gauge fixing

$$\chi(x, \tau) = \tau - N \cdot x , \quad N = (1, 0, 0, -1) , \quad N^2 = 0 , \quad (3.6)$$

where N has been written in ordinary coordinates. We see that $N^+ = N^0 + N^3 = 0$ which implies that the normal N to the hypersurface lies *within* the hypersurface

Figure 3.1: The hypersurface $\Sigma : x^+ = 0$ defining the front form. It is a null-plane tangential to the light-cone, $x^2 = 0$.



[37, 38]. As N is a light-like or null vector, Σ is often referred to as a *null-plane* [37, 39]. We have depicted the front-form hypersurface Σ together with the light-cone in Fig. 3.1.

The unit vector in x^+ -direction is another null-vector,

$$n^\mu = \frac{\partial x^\mu}{\partial x^+} = \frac{1}{2}(1, 0, 0, 1), \quad (3.7)$$

so that $n \cdot N = 1$ as it should. Given the scalar product (3.5), we infer the invariant distance element

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = dx^+ dx^- - dx^i dx^i = \left(\frac{dx^-}{dx^+} - \frac{dx^i}{dx^+} \frac{dx^i}{dx^+} \right) dx^+ dx^+ \quad (3.8)$$

from which the metric h can be read off as

$$h(x^+) = \dot{x}^- - \dot{x}^i \dot{x}^i \equiv v^- - \mathbf{v}_\perp^2. \quad (3.9)$$

Within the Hamiltonian formulation we find that $(p \cdot \partial)\chi = N \cdot p = p^+$, so that, from (2.38), the Lagrange multiplier λ becomes

$$\lambda = -1/2p^+ , \quad (3.10)$$

implying the equation of motion $\dot{x}^\mu = p^\mu/p^+$. Note that p^+ must be non-zero. The equations of motion are generated by the Hamiltonian

$$H = n \cdot p = p^-/2 = \frac{p_\perp^2 + m^2}{2p^+} , \quad (3.11)$$

by means of the Poisson brackets

$$\dot{x}^- = \{x^-, H\} = \frac{p^-}{p^+} , \quad \dot{x}^i = \{x^i, H\} = \frac{p^i}{p^+} . \quad (3.12)$$

and

$$\dot{p}^+ = \{p^+, H\} = 0 , \quad \dot{p}^i = \{p^i, H\} = 0 . \quad (3.13)$$

The solution is

$$x^-(x^+) = \frac{p^-}{p^+}x^+ + x^-(0) = v^-x^+ + x^-(0) \quad (3.14)$$

$$x^i(x^+) = \frac{p^i}{p^+}x^+ + x^i(0) = v^ix^+ + x^i(0) , \quad (3.15)$$

and again corresponds to linear motion with constant (front-form) velocity. Note that the Hamiltonian (3.11) is *not* the normal projection $N \cdot p$ of the momentum, because $N \cdot p$ lies within Σ and thus corresponds to a kinematical direction. As for the instant form, the light-cone representation of the Poincaré generators can be obtained by solving the constraint $p^2 = m^2$ for p^- , inserting the result into the elementary representation (2.64) of the generators and setting $x^+ = 0$. The kinematical generators are

$$P^i = p^i , \quad P^+ = p^+ , \quad M^{+i} = -x^ip^+ , \quad M^{+-} = -x^-p^+ , \quad M^{12} = x^1p^2 - x^2p^1 . \quad (3.16)$$

The dynamical ones (or Hamiltonians) are given by

$$P^- = \frac{p_\perp^2 + m^2}{p^+} , \quad M^{-i} = x^-p^i - x^ip^- . \quad (3.17)$$

We thus have found *seven* kinematical generators, so that the front form leads to the largest stability group. Furthermore, it is important to note that the Hamiltonian P^- does not contain a square root as already pointed out by Dirac. However, there is a singularity at $p^+ = 0$ which therefore is a somewhat peculiar point. In the gauge theory language, it corresponds to a vanishing Faddeev-Popov ‘matrix’. We will quite frequently be concerned with this issue in what follows.

3.2 Lorentz Boosts

Consider the following boost in z -direction with rapidity ω written in instant-form coordinates,

$$t \rightarrow t \cosh \omega + z \sinh \omega , \quad (3.18)$$

$$z \rightarrow t \sinh \omega + z \cosh \omega . \quad (3.19)$$

As stated before, such a boost mixes space and time coordinates z and t , respectively. However, if we add and subtract these equations, we obtain the action of the boost for the front form,

$$x^+ \rightarrow e^\omega x^+ , \quad (3.20)$$

$$x^- \rightarrow e^{-\omega} x^- . \quad (3.21)$$

We thus find the important result that a boost in z -direction does not mix light-cone space and time but rather rescales the coordinates! Note that x^+ and x^- are rescaled inversely with respect to each other. The scaling factor can be written as

$$e^\omega = \sqrt{\frac{1-v}{1+v}} , \quad (3.22)$$

if the rapidity ω is defined in the usual manner in terms of the boost velocity v , $\tanh \omega = v$. One should note in particular, that one has the fixed point hypersurface $\Sigma : x^+ = 0$ which is mapped onto itself, so that the relevant generator, $M^{+-} = 2M^{30} = -2K^3$, is kinematical. However, we see explicitly that this is not true for $x^+ \neq 0$, where we get a rescaling. Stated differently, the transformation to light-cone coordinates diagonalizes the boosts in z -direction. Therefore, the behavior under such boosts becomes especially simple. A pedagogical discussion of some elementary applications can be found in [40].

In our discussion of the boosts we are still left with the other two kinematical generators, the M^{+i} . These act on the transverse coordinates as

$$\delta x^i = \{x^i, M^{+j}\} \delta \beta^j = x^+ \delta \beta^i = 0 , \quad (3.23)$$

where the last identity holds on Σ . Translated to instant-form coordinates, M^{+i} is a combination of boosts and angular momenta,

$$M^{+i} = K^i + \epsilon^{ij} L^j . \quad (3.24)$$

The finite transformation corresponding to (3.23) is found via exponentiation. In general, if the infinitesimal transformation is written as

$$\delta_B A = \{A, B\} \equiv i \operatorname{ad} B(A) , \quad (3.25)$$

the finite transformation is given by

$$A' = \exp(i \operatorname{ad} B)(A) . \quad (3.26)$$

From (3.23) we thus obtain

$$x^i \rightarrow x^i + x^+ \beta^i, \quad (3.27)$$

which coincides with the infinitesimal transformation because the exponential series terminates after the term linear in the boost parameter β^i . Again, on Σ the x^i remain unchanged.

We are actually more interested in the transformation properties of the momenta, as these, being Poincaré generators, are more fundamental quantities than the coordinates, in particular in the quantum theory [31]. As P^μ transforms as a four-vector we just have to replace x^μ by P^μ in the boost transformations (3.20, 3.21, 3.27) and obtain,

$$P^+ \rightarrow e^\omega P^+, \quad (3.28)$$

$$P^- \rightarrow e^{-\omega} P^-, \quad (3.29)$$

$$P^i \rightarrow P^i + \beta^i P^+. \quad (3.30)$$

We remark that $P^+ = 0$ is a fixed point under longitudinal boosts. In quantum field theory, it corresponds to the vacuum. Note further, that in the third identity describing the action of M^{+i} , longitudinal and transverse momenta (which are both kinematical) get mixed.

We can now ask the question how to boost from (P^+, P^i) to momenta (Q^+, Q^i) . This can be done by fixing the boost parameters ω and β^i as

$$\omega = -\log \frac{Q^+}{P^+}, \quad \beta^i = \frac{Q^i - P^i}{P^+}. \quad (3.31)$$

Obviously, this is only possible for $P^+ \neq 0$. We emphasize that in the construction above there is no dynamics involved. For the quantum theory, this means that we can build states of arbitrary light-cone momenta with very little effort. All we have to do is applying some kinematical boost operators. The simple behavior of light-cone momenta under boosts will be important for the discussion of bound states. The same is true for the subject of the next section.

3.3 Galilei Subgroups and Consequences

Let us analyze the Poincaré algebra in light-cone coordinates in more detail. We are particularly interested in the Poisson bracket relations of the seven generators P^μ, M^{12}, M^{+i} ,

$$\begin{aligned} \{M^{12}, M^{+i}\} &= \epsilon^{ij} M^{+j}, \\ \{M^{12}, P^i\} &= \epsilon^{ij} P^j, \\ \{M^{+i}, P^-\} &= -2P^i, \\ \{M^{+i}, P^j\} &= -\delta^{ij} P^+ \end{aligned} \quad (3.32)$$

All other brackets of the seven generators vanish. Consider now the two-dimensional Galilei group. Its generators (for a free particle of mass m) are: two momenta k^i , one angular momentum $L = \epsilon^{ij}x^ik^j$, two Galilei boosts $G^i = mx^i$, the Hamiltonian $H = k^ik^i/2m$ and the mass m , which is the Casimir generator. Upon using $\{x^i, k^j\} = \delta^{ij}$ and identifying $P^i \leftrightarrow k^i$, $M^{12} \leftrightarrow L$, $M^{+i} \leftrightarrow -2G^i$, $P^+ \leftrightarrow 2m$ and $P^- \leftrightarrow H$, one easily finds that (3.32) forms a subalgebra of the Poincaré algebra which is isomorphic to the Lie algebra of the two-dimensional Galilei group. (A second isomorphic subalgebra is obtained via identifying $M^{-i} \leftrightarrow 2G^i$ and exchanging P^+ with P^- .) The first two identities in (3.32), for instance, state that M^{+i} and P^i transform as ordinary two-dimensional vectors.

One thus expects that light-cone kinematics will partly show a non-relativistic behavior which is associated with the transverse dimensions and governed by the two-dimensional Galilei group.

A very important example where these features are profitably at work is provided by a system of many particles which, for the time being, will be assumed as non-interacting. Let the i^{th} particle have mass m_i and light-cone four momentum

$$p_i = (p_i^+, \mathbf{p}_{\perp i}, p_i^-) . \quad (3.33)$$

As the particles are free, the total four momentum is given by the sum of the individual momenta,

$$P = \sum_i p_i . \quad (3.34)$$

Note that both the individual particles and the total system are on-shell,

$$p_i^- = \frac{p_{\perp i}^2 + m_i^2}{p_i^+} , \quad (3.35)$$

$$P^- = \frac{P_{\perp}^2 + M_0^2}{P^+} , \quad (3.36)$$

where M_0 denotes the total mass. We introduce *relative* momentum coordinates x_i and $\mathbf{k}_{\perp i}$ via

$$p_i^+ \equiv x_i P^+ , \quad (3.37)$$

$$\mathbf{p}_{\perp i} \equiv x_i \mathbf{P}_{\perp} + \mathbf{k}_{\perp i} . \quad (3.38)$$

Thus, x_i and $\mathbf{k}_{\perp i}$ denote the longitudinal momentum fraction and the relative transverse momentum of the i^{th} particle, respectively. Comparing with (3.34) we note that these variables have to obey the constraints

$$\sum_i x_i = 1 , \quad \sum_i \mathbf{k}_{\perp i} = 0 . \quad (3.39)$$

A particularly important property of the relative momenta is their boost invariance. To show this we calculate, using (3.28),

$$x'_i = e^{\omega} p_i^+ / e^{\omega} P^+ = x_i . \quad (3.40)$$

From this and (3.30) we find in addition

$$\mathbf{k}'_{\perp i} = \mathbf{p}'_{\perp i} - x_i \mathbf{P}'_{\perp} = \mathbf{p}_{\perp i} + \beta p_i^+ - x_i(\mathbf{P}_{\perp} + \beta P^+) = \mathbf{k}_{\perp i} , \quad (3.41)$$

which indeed proves the frame independence of x_i and $\mathbf{k}_{\perp i}$.

Let us calculate the total light-cone energy of the system in terms of the relative coordinates. Making use of the constraints (3.39), we obtain

$$\begin{aligned} P^- &= \sum_i p_i^- = \sum_i \frac{p_{\perp i}^2 + m_i^2}{p_i^+} = \sum_i \frac{(x_i \mathbf{P}_{\perp} + \mathbf{k}_{\perp i})^2 + m_i^2}{x_i P^+} \\ &= \frac{1}{P^+} \left(P_{\perp}^2 + \sum_i \frac{k_{\perp i}^2 + m_i^2}{x_i} \right) \equiv P_{\text{CM}}^- + P_r^- . \end{aligned} \quad (3.42)$$

This is another important result: the light-cone Hamiltonian P^- separates into a center-of-mass term,

$$P_{\text{CM}}^- = P_{\perp}^2 / P^+ , \quad (3.43)$$

and a term containing only the relative coordinates,

$$P_r^- = \frac{1}{P^+} \left(\sum_i \frac{k_{\perp i}^2 + m_i^2}{x_i} \right) \equiv \frac{M_0^2}{P^+} . \quad (3.44)$$

The second identity, which states that P_r^- is essentially the free invariant mass squared, follows upon multiplying (3.42) by P^+ ,

$$P^+ P_r^- = P^+ P^- - P_{\perp}^2 = M_0^2 = \sum_i \frac{k_{\perp i}^2 + m_i^2}{x_i} . \quad (3.45)$$

To simplify things even more, one often goes to the ‘transverse rest frame’ where $\mathbf{P}_{\perp}$ and therefore the center-of-mass Hamiltonian P_{CM}^- from (3.43) vanish.

Summarizing we note that the special behavior under boosts together with the transverse Galilei invariance leads to frame independent relative coordinates and a separation of the center-of-mass motion reminiscent of ordinary non-relativistic physics. This is at variance with the instant form, where the appearance of the notorious square root in the energy, $P^0 = (\mathbf{P}^2 + M_0^2)^{1/2}$, prohibits a similar separation of variables.

For a system of interacting particles, we add an interaction W to the free invariant mass squared defined in (3.45),

$$P^2 = M^2 = M_0^2 + W . \quad (3.46)$$

The total light-cone energy P^- is thus no longer given by the sum of the individual energies, but rather by

$$P^- = \frac{P_{\perp}^2 + M^2}{P^+} = \sum_i \frac{p_{\perp i}^2 + m_i^2}{p_i^+} + W/P^+ . \quad (3.47)$$

Though we will not enter an intensive discussion of quantization at this point, we note that, when translated to the quantum theory, (3.46) gives rise to the following light-cone Schrödinger equation, $(\hat{P}^2 - M^2)|\psi\rangle = 0$, or

$$(M^2 - \hat{M}_0^2)|\psi\rangle = \hat{W}|\psi\rangle, \quad (3.48)$$

where M^2 now denotes the mass-squared eigenvalue. In later chapters we will derive and solve such equations within quantum field theory.

Of course, in the interacting case, all dynamical Poincaré generators differ from their free counterpart by some ‘potential’ like W . This has already been pointed out by Dirac [32], who also stated that finding potentials which are consistent with the commutation relations of the Poincaré algebra is the “real difficulty in the construction of a theory of a relativistic dynamical system” with a fixed number of particles.

It has turned out, however, that Poincaré invariance alone is not sufficient to guarantee a reasonable Hamiltonian formulation. There are no-go theorems both for the instant [41] and the front form [42], which state that the inclusion of any potential into the Poincaré generators, even if consistent with the commutation relations, spoils relativistic *covariance*. The latter is a stronger requirement as it enforces particular transformation laws for the particle trajectories (world-lines). Thus, covariance imposes rather severe restrictions on the dynamical system [31].

The physical reason for these problems is that potentials imply an instantaneous interaction-at-a-distance which is in conflict with the existence of a limiting velocity and retardation effects. Relativistic causality is thus violated. This is equivalently obvious from the fact that a *fixed* number of particles is in conflict with the necessity of particle creation and annihilation and the appearance of anti-particles.

Nevertheless, with considerable effort, it is possible to construct dynamical quantum systems with a fixed number of constituents which are consistent with the requirements of Poincaré invariance and relativistic covariance [31, 43, 44].

At this point one might finally ask the question whether the different forms of relativistic dynamics are physically equivalent. From the point of view that different time choices correspond to different gauge fixings it is clear that equivalence must hold. After all, we are just dealing with different coordinate systems. People have tried to make this equivalence more explicit by working with coordinates which smoothly interpolate between the instant and the front form [45, 46, 47, 48]. In the context of relativistic quantum mechanics, it has been shown that the Poincaré generators for different forms are unitarily equivalent [43].

We are, however, more interested in what might be called a ‘top-down approach’. Our aim is to describe few-body systems not within quantum mechanics but quantum field theory to which we now turn.

Chapter 4

Light-Cone Quantization of Fields

4.1 Construction of the Poincaré Generators

We want to derive the representation of the Poincaré generators within field theory and their dependence on the hypersurface Σ chosen to define the time evolution. To this end we follow [33] and describe the hypersurface mathematically through the equation

$$\Sigma : F(x) = \tau . \quad (4.1)$$

The surface element on Σ is implicitly defined via

$$\int_{\Sigma} d\sigma_{\mu} u(x) = \int d^4x N^{\mu} \delta(F(x) - \tau) u(x) , \quad (4.2)$$

where, as before, $N^{\mu} = \partial^{\mu} F(x)$ is the normal on Σ and u some arbitrary function. We will write this expression symbolically as

$$d\sigma_{\mu} = d^4x N^{\mu} \delta(F(x) - \tau) . \quad (4.3)$$

The central object of this section will be the energy-momentum tensor,

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_{\mu}\phi)} \partial^{\nu} \phi - g^{\mu\nu} \mathcal{L} , \quad (4.4)$$

with $\mathcal{L}$ being the Lagrangian which depends on fields that are collectively denoted by ϕ . With the help of the energy-momentum tensor we can define a generator

$$A[f] = \int_{\Sigma} d\sigma_{\mu} f_{\nu}(x) T^{\mu\nu}(x) , \quad (4.5)$$

where A and f can be tensorial quantities carrying dummy indices $\alpha, \beta, \dots$ which we have suppressed. $A[f]$ generates the infinitesimal transformations

$$\delta_f B(x) = f_{\mu}(x) \partial^{\mu} B(x) , \quad (4.6)$$

$$\delta_f x^{\mu} = f^{\mu}(x) , \quad (4.7)$$

where f is now understood as being infinitesimal. In the same way as for a finite number of degrees of freedom, the generator A is called kinematical, if it leaves Σ invariant, that is,

$$\delta_f F = f_\mu \partial^\mu F = 0 . \quad (4.8)$$

Otherwise, A is dynamical. With the energy-momentum tensor $T^{\mu\nu}$ at hand we can easily show that kinematical generators are interaction independent. We decompose $T^{\mu\nu}$,

$$T^{\mu\nu} = T_0^{\mu\nu} - g^{\mu\nu} \mathcal{L}_{\text{int}} , \quad (4.9)$$

into a free part $T_0^{\mu\nu} = T^{\mu\nu}(g = 0)$, g denoting the coupling, and an interacting part (we exclude the case of derivative coupling). If A is kinematical, we have from (4.3, 4.5, 4.8),

$$A_{\text{int}}[f] = - \int d^4x \delta(F - \tau) \mathcal{L}_{\text{int}} f^\mu \partial_\mu F = - \int d^4x \delta(F - \tau) \mathcal{L}_{\text{int}} \delta_f F = 0 , \quad (4.10)$$

which indeed shows that A does not depend on the interaction. Of course, we are particularly interested in the Poincaré generators, P^α and $M^{\alpha\beta}$. The momenta P^α and Lorentz transformations $M^{\mu\nu}$ correspond to the choices $f_\mu^\alpha = g_\mu^\alpha$ and $f_\mu^{\alpha\beta} = x^\alpha g_\mu^\beta - x^\beta g_\mu^\alpha$, respectively so that, from (4.5), they are given in terms of $T^{\mu\nu}$ as

$$P^\alpha = \int_\Sigma d\sigma_\mu T^{\mu\alpha} , \quad (4.11)$$

$$M^{\alpha\beta} = \int_\Sigma d\sigma_\mu (x^\alpha T^{\mu\beta} - x^\beta T^{\mu\alpha}) . \quad (4.12)$$

From (4.8) it is easily seen that the Poincaré generators defined in (4.11, 4.12) act on $F(x) = \tau$ exactly as described in (2.67). The remarks of Chapter 2 on the kinematical or dynamical nature of the generators in the different forms are therefore equally valid in quantum field theory. Let us discuss the three main forms of relativistic dynamics explicitly.

The Instant Form

We recall the hypersurface of equal time,

$$\Sigma : F(x) \equiv N \cdot x \equiv x^0 = \tau , \quad (4.13)$$

which leads to a surface element

$$d\sigma^\mu = d^4x N^\mu \delta(x^0 - \tau) , \quad N^\mu = (1, \mathbf{0}) . \quad (4.14)$$

Using (4.11, 4.12), the Poincaré generators are obtained as

$$P^\mu = \int_\Sigma d^3x T^{0\mu} , \quad (4.15)$$

$$M^{\mu\nu} = \int_\Sigma d^3x (x^\mu T^{0\nu} - x^\nu T^{0\mu}) . \quad (4.16)$$

The Front Form

Hypersurface and surface element are given by

$$\Sigma : F(x) \equiv N \cdot x \equiv x^+ = \tau , \quad d\sigma^\mu = d^4x N^\mu \delta(x^+ - \tau) , \quad (4.17)$$

where N is the light-like four-vector $N = (1, 0, 0, -1)$, $N^2 = 0$ of (3.6). In terms of $T^{\mu\nu}$, the Poincaré generators are

$$P^\mu = \frac{1}{2} \int_\Sigma dx^- d^2x_\perp T^{+\mu} , \quad (4.18)$$

$$M^{\mu\nu} = \frac{1}{2} \int_\Sigma dx^- d^2x_\perp (x^\mu T^{+\nu} - x^\nu T^{+\mu}) . \quad (4.19)$$

The somewhat peculiar factor $1/2$ is the Jacobian which arises upon transforming to light-cone coordinates.

The Point Form

As we have already seen, the point form is slightly more involved. The hypersurface of equal time,

$$\Sigma : F(x) = \sqrt{x^2} = \tau > 0 , \quad x^2 > 0 , \quad (4.20)$$

has the radial vector $N^\mu = x^\mu/\tau$ as its normal. In terms of the hyperbolic coordinates (2.78), the Poincaré generators become

$$P^\mu = \tau^2 \int_\Sigma d\omega d\theta d\phi \sinh^2 \omega \sin \theta x_\alpha T^{\mu\alpha} , \quad (4.21)$$

$$M^{\mu\nu} = \tau^2 \int_\Sigma d\omega d\theta d\phi \sinh^2 \omega \sin \theta x_\alpha (x^\mu T^{\alpha\nu} - x^\nu T^{\alpha\mu}) . \quad (4.22)$$

Note again that the limit $\tau \rightarrow 0$ does not make sense.

4.2 Schwinger's (Quantum) Action Principle

Our next task is to actually quantize the fields on the hypersurfaces $\Sigma : \tau = F(x)$ of equal time τ . There is more than one possibility to do so, and we will explain a few of these. We begin with a method that is essentially due to Schwinger [49, 50, 51]. We define a four-momentum density

$$\Pi^\mu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} , \quad (4.23)$$

so that the energy-momentum tensor $T^{\mu\nu}$ can be written as

$$T^{\mu\nu} = \Pi^\mu \partial^\nu \phi - g^{\mu\nu} \mathcal{L} . \quad (4.24)$$

In some sense, this can be viewed as a covariant generalization of the usual Legendre transformation between Hamiltonian and Lagrangian. Using the normal N^μ of the hypersurface Σ , we define the canonical momentum (density) as the projection of Π^μ ,

$$\pi \equiv N \cdot \Pi . \quad (4.25)$$

Schwinger's action principle states that upon variation the action $S = \int d^d x \mathcal{L}$ changes at most by a surface term which (if Σ is not varied, i.e. $\delta x^\mu = 0$) is given by

$$\delta G(\tau) = \int_{\Sigma} d\sigma_\mu \Pi^\mu \delta\phi = \int d^d x \delta(\tau - F) \pi \delta\phi \quad (4.26)$$

taken between the initial and final hypersurfaces Σ_1 and Σ_2 . The quantity δG is interpreted as the generator of field transformations, so that we have

$$\delta\phi = \{\phi, \delta G\} , \quad (4.27)$$

in case that Σ is entirely space-like (with time-like normal) [49, 50]. We note in passing that the generator δG is a field theoretic generalization of the canonical one-form used in analytical mechanics. In the last expression we have been a bit sloppy with the space-time arguments, so let us improve on this. As in the preceding chapter, we parametrize space-time with foliation coordinates ξ^μ , with $\xi^0 = \tau$ and the ξ^i living on Σ . Then we calculate

$$\begin{aligned} \delta\phi(\xi) &= \{\phi(\xi), \delta G(\tau)\} \\ &= \int d\xi^0 \int d^{d-1}\xi' J(\xi') \delta(\tau - \xi'^0) \{\phi(\xi), \pi(\xi')\} \delta\phi(\xi') \\ &= \int_{\Sigma} d^{d-1}\xi' J(\xi') \{\phi(\xi), \pi(\xi')\} \delta\phi(\xi') \Big|_{\xi^0=\xi'^0=\tau} , \end{aligned} \quad (4.28)$$

where $J(\xi)$ denotes the Jacobian associated with the change of variables $x^\mu \rightarrow \xi^\alpha$. From the last expressions we infer that the canonical equal-time bracket must be

$$\{\phi(\xi), \pi(\xi')\}_{\xi, \xi' \in \Sigma} = \{\phi(\xi), \pi(\xi')\}_{\xi^0=\xi'^0=\tau} = J^{-1}(\xi) \delta^{d-1}(\xi - \xi') . \quad (4.29)$$

As usual, commutators are abstracted from the basic bracket by invoking Dirac's correspondence principle, that is, by replacing the bracket by i times the commutator. For arbitrary classical observables, A, B , this means explicitly,

$$[\hat{A}, \hat{B}] = i \widehat{\{A, B\}} , \quad (4.30)$$

With minor modifications, the approach leading to (4.29) can also be used for the front form, resulting in what is called light-cone or light-front quantization. The canonical light-cone momentum is

$$\pi = N \cdot \Pi = N \cdot \partial\phi = \partial^+ \phi \equiv 2 \frac{\partial}{\partial x^-} \phi , \quad (4.31)$$

which is peculiar to the extent that it does not involve a (light-cone) time derivative. Therefore, π is a dependent quantity which does not provide additional information, being known on Σ when the field is known there, so that π is merely an abbreviation for $\partial^+\phi$. Again, the reason is that the normal N^μ of the null-plane Σ lies within Σ . As a result, ϕ and $\partial^+\phi$ have to be treated on the same footing within Schwinger's approach which leads to an additional factor 1/2 compared to (4.27),

$$\frac{1}{2}\delta\phi = \{\phi, \delta G\} , \quad (4.32)$$

with a front-form generator

$$\delta G(x^+) = \frac{1}{2} \int_{\Sigma} dx^- d^2x_{\perp} \partial^+\phi \delta\phi . \quad (4.33)$$

The appearance of the peculiar factor 1/2 in (4.32) has been discussed at length by Schwinger in [51] (see also [52]). This factor cancels the light-cone Jacobian $J = 1/2$ in (4.33), so that we are left with the equal-time commutator (for $d = 4$),

$$[\phi(x), \pi(y)]_{x^+=y^+=\tau} = i\delta(x^- - y^-)\delta^2(\mathbf{x}_{\perp} - \mathbf{y}_{\perp}) \quad (4.34)$$

As the independent quantities are the fields themselves, we invert the derivative ∂^+ and obtain the more fundamental commutator

$$[\phi(x), \phi(y)]_{x^+=y^+=\tau} = -\frac{i}{4}\text{sgn}(x^- - y^-)\delta^2(\mathbf{x}_{\perp} - \mathbf{y}_{\perp}) . \quad (4.35)$$

In deriving (4.35) we have chosen the anti-symmetric Green function $\text{sgn}(x^-)$ satisfying

$$\frac{\partial}{\partial x^-}\text{sgn}(x^-) = 2\delta(x^-) , \quad (4.36)$$

so that (4.34) is re-obtained upon differentiating (4.35) with respect to y^- . We will see later that the *field* commutator (4.35) can be directly obtained from Schwinger's method. Before that, however, we will study the relation between the choice of initializing hypersurfaces, the problem of field quantization and the solutions of the dynamical equations.

4.3 Quantization as an Initial/Boundary-Value Problem

As a prototype field theory we consider a massive scalar field in 1+1 dimensions. Its dynamics is encoded in the action

$$S[\phi] = \int d^2x \mathcal{L} = \int d^2x \left(\frac{1}{2}\partial_{\mu}\phi\partial^{\mu}\phi - \frac{1}{2}m^2\phi^2 - \mathcal{V}[\phi] \right) , \quad (4.37)$$

where $\mathcal{V}$ is some interaction term like e.g. $\lambda\phi^4$ and $\mathcal{L} = \mathcal{L}_0 + \mathcal{V}$. By varying the free action in the standard way we obtain

$$\delta S = \int_{\partial M} d\sigma_\mu \Pi^\mu \delta\phi + \int_M \left[\frac{\partial \mathcal{L}_0}{\partial \phi} - \partial_\mu \frac{\partial \mathcal{L}_0}{\partial (\partial_\mu \phi)} \right] \delta\phi. \quad (4.38)$$

If we do not vary on the boundary of our integration region M , $\delta\phi|_{\partial M} = 0$, the surface term in δS vanishes and we end up with the (massive) Klein-Gordon equation in 1+1 dimensions,

$$(\square + m^2)\phi = 0. \quad (4.39)$$

We are going to solve this equation by specifying initial and/or boundary conditions for the scalar field ϕ on different hypersurfaces Σ . In particular we will clarify the relation between the associated initial value problems and the determination of ‘equal-time’ commutators.

It may look rather trivial to consider just the free theory, but this is not really true. Let us analyze what quantization of a field theory means in the light of the different forms of relativistic dynamics. One specifies canonical commutators like $[\phi(x), \phi(0)]_{x \in \Sigma}$, where the equation for the hypersurface Σ : $\tau = \text{const}$ defines an evolution parameter τ and the notion of ‘equal time’ as already pointed out several times. Σ should be chosen such that the interaction does not change the commutator on it, i.e. the latter is a kinematical quantity.

Now, if ϕ is a *free* field, the commutator,

$$[\phi(x), \phi(0)] = i\Delta(x), \quad (4.40)$$

is exactly known: it is the Pauli-Jordan or Schwinger function Δ which is a special solution of the Klein-Gordon equation (4.39). It can be obtained directly from the action in a covariant manner as a Peierls bracket [53, 54]. Alternatively, one can find it from its Fourier representation which yields [55]

$$\begin{aligned} \Delta(x) &= -\frac{i}{2\pi} \int d^2p \delta(p^2 - m^2) \text{sgn}(p^0) e^{-ip \cdot x} \\ &= -\frac{1}{2} \text{sgn}(x^0) \theta(x^2) J_0(m\sqrt{x^2}) \\ &= -\frac{1}{4} [\text{sgn}(x^+) + \text{sgn}(x^-)] J_0(m\sqrt{x^+ x^-}), \end{aligned} \quad (4.41)$$

We note that it is antisymmetric, $\Delta(x) = -\Delta(-x)$, and vanishes outside the light-cone, i.e. for $x^2 < 0$.

If ϕ is an *interacting* field one still has some information from causality. For space-like hypersurfaces (instant and point form) the equal-time commutator vanishes like in the free field case,

$$[\phi(x), \phi(0)]_{x^2 < 0} = 0. \quad (4.42)$$

This expresses the fact that fields which are separated by a space-like distance cannot communicate with each other. For the front form with hypersurface $\Sigma : x^+ = 0$, the situation is slightly different. In 1+1 dimensions, Σ is the light-cone and therefore entirely light-like. In higher dimensions, Σ still contains light-like directions namely where $x^- = \mathbf{x}_\perp = 0$. This can also be seen from the (free) commutator (4.35) which does not vanish at these points corresponding to a light-like separation. It may therefore well be that the interaction changes the light-cone behavior of the commutator. This is actually seen in deep-inelastic scattering, where the assumption of free-field commutation relations leads to the parton model and scaling [56], which, however, is violated in the interacting theory, i.e. QCD (though only weakly).

Concluding, we can say that, if the fields are separated by a space-like distance, the free field ‘equal-time’ commutators are boundary conditions also for the interacting theory and thus of sizable interest.

4.3.1 The Cauchy Problem

Before discussing the case of the front form, we would like to recall that the conventional quantization on a space-like surface (based on the instant form) corresponds to a Cauchy problem: if one specifies the field ϕ and its time derivative $\dot{\phi}$ on $\Sigma : x^0 = 0$, the solution of the Klein-Gordon equation is uniquely determined as [57, 37]

$$\phi(x) = \int_{y^0=0} dy^1 \left[\phi(y) \frac{\overleftrightarrow{\partial}}{\partial y^0} \Delta(x-y) \right]. \quad (4.43)$$

This solution expresses the well-known fact that the creation and annihilation operators in the Fock expansion of the quantum field are given in terms of the field *and* the velocity on Σ ,

$$a(p^1) = \int dx^1 e^{-ip^1 x^1} \left[\omega_p \phi(x^0 = 0, x^1) + i\dot{\phi}(x^0 = 0, x^1) \right], \quad (4.44)$$

with $\omega_p = (p_1^2 + m^2)^{1/2}$. If we compare (4.43) with the two independent canonical commutators

$$\begin{aligned} [\phi(x), \phi(0)]_{x^0=0} &= i\Delta(x)|_{x^0=0} = 0, \\ [\dot{\phi}(x), \phi(0)]_{x^0=0} &= i\dot{\Delta}(x)|_{x^0=0} = -i\delta(x^1), \end{aligned} \quad (4.45)$$

we see that they can be viewed as Cauchy data for the Pauli-Jordan function Δ . Using the data (4.45) in formula (4.43) we obtain a consistency condition,

$$[\phi(x), \phi(0)] = \int_{y^0=0} dy^1 [\phi(y), \phi(0)] \frac{\overleftrightarrow{\partial}}{\partial y^0} \Delta(x-y), \quad (4.46)$$

which expresses the commutator for arbitrary time differences x^0 in terms of its initial values given by (4.45).

4.3.2 The Characteristic Initial-Value Problem

In the following we will perform an analogous discussion for the hypersurfaces $\Sigma : x^\pm = \text{const}$, which, in $d+1+1$, constitute the light-cone. In Dirac's classification, the light-cone corresponds to a degenerate point form with parameter $a = 0$. One thus does not have transitivity as points on different 'legs' of the cone are not related by a kinematical operation. The Hamiltonians are P^+ and P^- , so that a Hamiltonian formulation appears a little bit strange and will involve two time parameters. We will not pursue this issue here. A Lagrangian formulation, however, is no problem; it just leads to the Klein-Gordon equation. Let us discuss how to solve it using initial data on Σ .

The light-fronts $x^\pm = 0$ are characteristics of the Klein-Gordon equation [58]. Therefore, one is dealing with a characteristic initial-value problem [59, 60]: if one specifies the field ϕ on $x^+ = x_0^+$ and $x^- = x_0^-$, say $\phi(x^+, x_0^-) = f(x^+)$ and $\phi(x_0^+, x^-) = g(x^-)$, the solution of the Klein-Gordon equation is [37, 38]

$$\phi(x) = \int_{x_0^+}^{\infty} dy^+ f(y^+) \frac{\overleftrightarrow{\partial}}{\partial y^+} \Delta(x-y) \Big|_{y^- = x_0^-} + \int_{x_0^-}^{\infty} dy^- g(y^-) \frac{\overleftrightarrow{\partial}}{\partial y^-} \Delta(x-y) \Big|_{y^+ = x_0^+}. \quad (4.47)$$

This amounts to quantization on *two* characteristics, $x^\pm = 0$, i.e., in $d = 1 + 1$, really *on* the light cone, $x^2 = 0$. The following two independent commutators,

$$[\phi(x), \phi(0)]_{x^\pm=0} = i\Delta(x)|_{x^\pm=0} = -\frac{i}{4}\text{sgn}(x^\mp), \quad (4.48)$$

are then characteristic data for the Pauli-Jordan function. Again this can be cast in the form of a consistency condition,

$$\begin{aligned} [\phi(x), \phi(0)] &= \int_0^\infty dy^+ [\phi(y), \phi(0)] \frac{\overleftrightarrow{\partial}}{\partial y^+} \Delta(x-y) \Big|_{y^- = 0} + \\ &+ \int_0^\infty dy^- [\phi(y), \phi(0)] \frac{\overleftrightarrow{\partial}}{\partial y^-} \Delta(x-y) \Big|_{y^+ = 0}, \end{aligned} \quad (4.49)$$

which expresses the commutator in terms of its characteristic initial values.

It turns out that, in case the field ϕ is massless, the above quantization procedure is the only consistent one (in $d=1+1$), if one wants to use light-like hypersurfaces [61].

4.3.3 The Hamiltonian Formulation

In this subsection we want to analyze whether a (Hamiltonian) quantization on a *single* light-front is possible in the massive case. Note that in the instant form

with its corresponding Cauchy problem everything is quite clear. The data for $\dot{\phi}$ and $\dot{\phi}$ translate into data for ϕ and the momentum π , which is independent of ϕ . We already know that within the front form there is no independent momentum as the normal of a null-plane lies within the null-plane. If one does not want to use a second light-front in order to avoid an awkward Hamiltonian formulation with two times, it seems that one only needs half of the data as compared to the Cauchy or the characteristic initial-value problem [52]. This seems strange and, indeed, is not the case as we will see.

Let us write the Klein-Gordon equation (4.39) in terms of light-cone coordinates

$$(\partial^+ \partial^- + m^2)\phi = 0 , \quad (4.50)$$

and solve for $\dot{\phi} \equiv \partial\phi/\partial x^+$ (which amounts to inverting $\partial^+ = 2\partial/\partial x^-$). This gives

$$\dot{\phi}(x^+, x^-) = -\frac{m^2}{4} \int dy^- G(x^-, y^-) \phi(x^+, y^-) , \quad (4.51)$$

where the Green function G is an inverse of the derivative $\partial/\partial x^-$,

$$\frac{\partial}{\partial x^-} G(x^-, y^-) = \delta(x^- - y^-) . \quad (4.52)$$

The Hamiltonian equation of motion is given by the Poisson bracket with $H = \frac{1}{2}m^2 \int dx^- \phi^2$,

$$\dot{\phi}(x^+, x^-) = \frac{m^2}{2} \int dy^- \{ \phi(x^+, x^-) , \phi(x^+, y^-) \} \phi(x^+, y^-) \quad (4.53)$$

with a presently unknown bracket of the fields ϕ . Clearly, Euler-Lagrange and Hamiltonian equation of motion, (4.51) and (4.53) have to be equivalent. This is realized if one identifies

$$\{ \phi(x^+, x^-) , \phi(x^+, y^-) \} \equiv -\frac{1}{2}G(x^-, y^-) , \quad (4.54)$$

Upon quantization, this should, of course, coincide with (4.35) specialized to $d = 1 + 1$. To make the whole procedure of solving the equations of motion unambiguous, we need an initial condition for ϕ on Σ and a *further* condition which makes the inversion of ∂^+ unique. Thus, as announced, we need again *two* conditions.

After quantization the basic bracket (4.54) becomes the free field commutator according to the correspondence principle. The free field commutator on the light front $x^+ = 0$ thus essentially is the Green function G . In momentum space, G is given by (a regularization of) the expression $1/p^+$. In this sense, quantization on $x^+ = 0$ amounts to inversion of the light-cone momentum p^+ (viewed as a distribution). It is well known that there is no unique inverse (or regularization) of $1/p^+$ [62]. The possible regularizations differ by terms proportional to $\delta(p^+)$

which correspond to homogeneous solutions of (4.52). To solve this problem, one has to specify additional conditions with respect to the variable x^- which is conjugate to p^+ .

It turns out that data on a second characteristic, say $\phi(x^+, x^- = x_0^-) = f(x^+)$, do not work: they do not lead to an antisymmetric Green function in (4.51) which is needed for the commutator. Therefore one does not obtain equivalence between Euler-Lagrange and Hamiltonian equations of motion, which is a severe inconsistency. As a result, there is no Hamiltonian formulation with data on *two* characteristics and a *single* light-cone Hamiltonian.

There exists, however, an alternative. We follow the philosophy of an approach called ‘discretized light-cone quantization’ [63, 64, 65] and enclose the system in a finite volume, $-L \leq x^- \leq L$. Upon demanding periodic boundary conditions for the field,

$$\phi(x^+, x^- = -L) = \phi(x^+, x^- = L) , \quad (4.55)$$

which are to hold for all light-cone times x^+ , the longitudinal momenta become discrete,

$$k_n^+ = 2\pi n/L , \quad n \in \mathbb{Z} . \quad (4.56)$$

As the spatial boundaries are not separated by a space-like but a light-like distance, one has to check whether the boundary conditions (4.55) are consistent with the equation of motion and thus with relativistic causality. This will be done in the next subsection. Assuming this consistency, the Euler-Lagrange equation (4.51) becomes

$$\dot{\phi}(x^+, x^-) = -\frac{m^2}{4} \int_{-L}^L dy^- G_p(x^-, y^-) \phi(x^+, y^-) , \quad (4.57)$$

where G_p is uniquely determined as the periodic sign function,

$$G_p(x^-, y^-) = \frac{1}{2} \text{sgn}(x^- - y^-) - \frac{x^- - y^-}{2L} , \quad (4.58)$$

so that there is no arbitrariness left. From (4.54) we obtain a unique and consistent commutator,

$$[\phi(x), \phi(0)]_{x^+=0} = -\frac{i}{2} G_p(x^-) = -\frac{i}{2} \left[\frac{1}{2} \text{sgn}(x^- - y^-) - \frac{x^- - y^-}{2L} \right] . \quad (4.59)$$

Apart from the second term, which is a finite volume correction, this coincides with (4.35). We are left with checking the consistency of the procedure above.

4.3.4 The Initial-Boundary-Value Problem

The attempt to solve the Klein-Gordon equation with periodic boundary conditions in x^- poses an initial-boundary-value problem which is explicitly defined

through the conditions

$$\phi(x^+ = 0, x^-) = g(x^-), \quad (4.60)$$

$$\phi(x^+, x^- = L) = \phi(x^+, x^- = -L), \quad (4.61)$$

where (4.60) represents the initial data and (4.61) the boundary conditions. The unique solution of the Klein-Gordon equation obeying these conditions is obtained via Fourier transformation,

$$\phi(x^+, x^-) = \sum_{n \neq 0} g_n e^{-ik_n^+ x^- / 2 - i\hat{k}_n^- x^+ / 2} \equiv \sum_{n \neq 0} g_n e^{-i\hat{k}_n \cdot x}. \quad (4.62)$$

The on-shell energy is $k_n^- = m^2/k_n^+ = m^2 L / 2\pi n$, and the g_n are the Fourier modes of the function $g(x^-)$ specified in (4.60), i.e.

$$g_n = \frac{1}{2L} \int_{-L}^L dx^- g(x^-) e^{in\pi x^- / L} \quad (4.63)$$

Investigating the Klein-Gordon equation in momentum space one finds that the zero mode g_0 is vanishing (see also Section 4.4). It is therefore already omitted in (4.62). From this solution it is obvious that data on a second characteristic are not necessary. In the quantum theory, (4.63) corresponds to the peculiar fact that the light-cone Fock operators are expressible in terms of the field alone; the velocity is not needed [66, 52].

Another consistency check is to demand periodic boundary conditions for the solution (4.47) of the characteristic initial-value problem. Together with the data $f(x^+)$ this would over-determine the problem, which can only be avoided, if f is expressible in terms of g . This is indeed the case,

$$f(x^+) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dp e^{px^+} \frac{H_L[g, p]}{1 - \exp(-m^2 L / 2p)}, \quad (4.64)$$

where $H_L[g, p]$ is a complicated but known functional of the data g [55].

In addition, let us check the consistency of the solution (4.53) for an important example, the periodic Pauli-Jordan function,

$$\Delta_L(x^+, x^-) \equiv - \sum_{n \neq 0} \frac{i}{4\pi n} e^{-i\hat{k}_n^- x^+ / 2 - in\pi x^- / L}. \quad (4.65)$$

In analogy with (4.49) this leads to the light-cone commutators

$$\left[\phi(x), \phi(0) \right]_{x^+=0} = -\frac{i}{2} \left[\frac{1}{2} \text{sgn}(x^-) - \frac{x^-}{2L} \right] \equiv g(x^-), \quad (4.66)$$

$$\left[\phi(x), \phi(0) \right]_{x^-=-L} = \frac{1}{4\pi} \sum_{n \neq 0} \frac{(-1)^n}{n} \exp\left(-i \frac{m^2 L}{4\pi n} x^+\right) \equiv f(x^+), \quad (4.67)$$

where (4.66) coincides with (4.59), as expected. Due to the periodic boundary conditions the second commutator given by $f(x^+)$ is not independent but satisfies exactly (4.64) with g given by (4.66). As in the case of the Cauchy problem and the characteristic initial-value problem one can write down a consistency condition,

$$[\phi(x), \phi(0)] = \int_{-L}^L dy^- [\phi(y), \phi(0)] \frac{\partial}{\partial y^-} \Delta_L(x - y) \Big|_{y^+=0}, \quad (4.68)$$

which expresses the commutator through its value on *one* characteristic ($y^+ = 0$) and the *periodic* Pauli-Jordan function.

4.3.5 Fermions

For fermions, light-cone quantization is slightly more involved because not all of the fermion field components are independent. This is easily seen by considering the free massive Dirac equation (again, for simplicity, in 1+1 dimensions),

$$(i\partial - m)\psi = 0. \quad (4.69)$$

If we write the two-spinor ψ in terms of a lower and upper component as

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad (4.70)$$

the Dirac equation (4.69) splits up into

$$\partial^- \psi_1 = -im\psi_2, \quad (4.71)$$

$$\partial^+ \psi_2 = -im\psi_1. \quad (4.72)$$

The important point to note is that (4.72) represents a constraint as ∂^+ is a spatial rather than a temporal derivative. It should be solved for the dependent lower component ψ_2 in terms of the independent upper component ψ_1 ,

$$\psi_2 = -im(\partial^+)^{-1}\psi_1. \quad (4.73)$$

Obviously, at this point, the ubiquitous inverse of ∂^+ arises again, and we have to make sure that it becomes well defined. Like for the scalar field, the mentioned ambiguity affects the quantization procedure. The covariant anti-commutator of two fermion fields with arbitrary space-time separation x is given in terms of an invariant function S ,

$$\{\psi(x), \bar{\psi}(0)\} = iS(x), \quad (4.74)$$

where S can be inferred from the Pauli-Jordan function as

$$S(x) = (i\partial + m)\Delta(x). \quad (4.75)$$

From this representation it is obvious that S obeys the Dirac equation (as it must), as Δ is a solution of the Klein-Gordon equation. We thus expect that S will also contain dependent degrees of freedom. To make these explicit, we introduce a two-by-two matrix notation and write (4.74) in the form

$$\left\{ \psi_i(x), \psi_j^\dagger(0) \right\} = iS_{ij}(x), \quad i, j = 1, 2. \quad (4.76)$$

According to (4.75), the components iS_{ij} are

$$iS_{11}(x) = -\partial^+ \Delta(x), \quad (4.77)$$

$$iS_{12}(x) = iS_{21}(x) = im\Delta(x), \quad (4.78)$$

$$iS_{22}(x) = -\partial^- \Delta(x). \quad (4.79)$$

Comparing the second and the third identity we learn that S_{22} is a dependent quantity satisfying the constraint (4.72),

$$\partial^+ S_{22} = -imS_{12}. \quad (4.80)$$

Of course, taking into account (4.76), this is consistent with the fact that ψ_2 is the dependent field component. There is, however, one subtlety that deserves a separate analysis. If one performs the derivatives in (4.75) and writes the Pauli-Jordan function (4.41) as

$$\Delta(x) \equiv D(x)J_0(m\sqrt{x^2}), \quad (4.81)$$

the iS_{ij} are explicitly found to be

$$iS_{11}(x) = \delta(x^-) + mD(x)\sqrt{x^+/x^-}J_1(m\sqrt{x^2}), \quad (4.82)$$

$$iS_{12}(x) = imD(x)J_0(m\sqrt{x^2}), \quad (4.83)$$

$$iS_{22}(x) = \delta(x^+) + mD(x)\sqrt{x^-/x^+}J_1(m\sqrt{x^2}). \quad (4.84)$$

There are several remarks in order. First of all, the first identity provides the canonical anti-commutator of the independent field components,

$$\left\{ \psi_1(x), \psi_1^\dagger(0) \right\}_{x^+=0} = iS_{11}(x^+ = 0, x^-) = \delta(x^-). \quad (4.85)$$

In addition, we note that S_{11} and S_{22} are related by an exchange of x^+ and x^- . Going back to instant-form coordinates, this amounts to the spatial inversion $x^1 \rightarrow -x^1$, that is, a *parity* transformation. Obviously, as parity exchanges light-cone space and time, it is a dynamical and therefore complicated transformation. Furthermore, the leading light-cone singularities of the S_{ii} are mass independent, as is well known from, say, the space-time analysis of deep-inelastic scattering [67]. In particular, we have

$$\lim_{m \rightarrow 0} iS_{22}(x) \equiv iS_{22}^0 = iS_{22}(x^- = 0) = \delta(x^+) \quad (4.86)$$

On the other hand, if we formally solve (4.80) for S_{22} , we obtain

$$S_{22}(x) = -im(\partial^+)^{-1}S_{12} , \quad (4.87)$$

which, at first glance, seems to be in conflict with (4.86) upon performing the limit $m \rightarrow 0$. One might be tempted to add the homogeneous solution S_{22}^0 to the right-hand-side of (4.87). This would imply that S_{22} is not entirely determined in terms of S_{12} , and one would have to provide $S_{22}(x^+, x^- = 0)$ as a boundary condition for *all light-cone times* x^+ . Analogous arguments apply to the relation between ψ_2 and ψ_1 . In the language of the preceding subsections this would correspond to a characteristic initial-value problem. Fortunately, these difficulties can be circumvented, if one makes use of parity, as we are now going to show.

We assume knowledge of the field ψ_1 at light-cone time $x^+ = 0$. For arbitrary times x^+ it is then given by

$$\psi_1(x) = \frac{1}{\sqrt{2\pi}} \int dk^+ \left[b(k^+)e^{-i\hat{k}\cdot x} + d^\dagger(k^+)e^{i\hat{k}\cdot x} \right] , \quad (4.88)$$

where

$$\hat{k} \cdot x = \frac{1}{2}k^+x^- + \frac{1}{2}\hat{k}^-x^+ , \quad \hat{k}^- = m^2/k^+ , \quad (4.89)$$

and the Fock operators obey the anti-commutation relations

$$\{b(k^+), b^\dagger(q^+)\} = \{d(k^+), d^\dagger(q^+)\} = \delta(k^+ - q^+) . \quad (4.90)$$

Plugging (4.88) into (4.73), the dependent field becomes

$$\psi_2(x) = \frac{m}{\sqrt{2\pi}} \int \frac{dk^+}{k^+} \left[b(k^+)e^{-i\hat{k}\cdot x} + d^\dagger(k^+)e^{i\hat{k}\cdot x} \right] , \quad (4.91)$$

from which one infers the anti-commutator

$$iS_{22}(x) = \frac{m^2}{2\pi} \int \frac{dk^+}{(k^+)^2} [e^{-i\hat{k}\cdot x} + e^{i\hat{k}\cdot x}] . \quad (4.92)$$

Note the m^2 in front of the integral which is consistent with (4.87) as S_{12} is of order m . The trick is now to invoke parity by substituting $k^+ = m^2/k^-$ with k^- as the new integration variable. This absorbs the m^2 in front of the integral and results in

$$iS_{22}(x) = \int \frac{dk^-}{2\pi} \exp(-im^2x^-/k^- - ik^-x^+) . \quad (4.93)$$

Evaluating this for $x^- = 0$, we finally obtain

$$iS_{22}(x^+, x^- = 0) = \{\psi_2(x), \psi_2(0)\}_{x^-=0} = \delta(x^+) . \quad (4.94)$$

We reemphasize that this result was obtained by only assuming initial data (in terms of the usual Fock expansion) for the independent field component ψ_1 , and parity. Data on the second characteristic are once more unnecessary.

Altogether, we have thus shown that, for massive bosons and fermions in space-time dimension $d = 1+1$, quantization on a *single* light front $x^+ = 0$ is possible. However, there are special requirements to be met. In the bosonic case, periodic boundary conditions in the spatial direction lead to a consistent and simple light-cone Hamiltonian formulation. Data on a second characteristic are not necessary. The result justifies the approach of discretized light-cone quantization [63, 64, 65]. For fermions, we have used parity as an additional input.

For $d = 3+1$ analogous arguments apply because the solution of the Klein-Gordon equation is still of the form (4.47) with just additional integrations over the transverse dimensions [37, 38].

4.4 First-Order Systems

In this section we will encounter yet another method of Hamiltonian reduction, i.e. of identifying and quantizing the physical degrees of freedom. It will serve us to construct the light-cone Fock space. We will restrict our attention to the bosonic case. According to (4.37), the Lagrangian of a scalar field in 1+1 dimensions, when written in terms of light-cone coordinates, takes the following form,

$$L[\phi] = \frac{1}{2} \int dx^- \left(\frac{1}{2} \partial^+ \phi \partial^- \phi - \frac{1}{2} m^2 \phi^2 - \mathcal{V}[\phi] \right) . \quad (4.95)$$

The important thing to note is the fact that the Lagrangian is *linear* in the light-cone velocity $\partial^- \phi = 2\partial\phi/\partial x^+$. This means that the appropriate method for quantization is the one of Faddeev and Jackiw for first order systems [68] called “(Constrained) Quantization Without Tears” by Jackiw [69]. It avoids many of the technicalities of the Dirac-Bergmann formalism and is in general more economic. It reduces phase-space right from the beginning as there are no ‘primary constraints’ introduced. The method is essentially equivalent to Schwinger’s action principle, especially in the form presented by Schwinger in [51].

The approach becomes particularly transparent for the case at hand if we again discretize the system in momentum space [63, 64, 65] by enclosing it in a finite spatial volume, $-L \leq x^- \leq L$. We proceed by Fourier expanding the classical free field as in (4.62),

$$\phi(x^+, x^-) = a_0 + \sum_{n \neq 0} g_n e^{-i(k_n^+ x^- / 2 + \hat{k}_n^- x^+ / 2)} . \quad (4.96)$$

Note that we have allowed for a zero-momentum mode a_0 in the expansion (4.96). In what follows we will show that it actually vanishes. If we define $a_n \equiv (4\pi|n|)^{-1/2}$, separate positive and negative modes in (4.96), and use the fact that $a_{-n} = a_n^*$ (as ϕ is real), the summation can be restricted to run over

only positive modes. At light-cone time $x^+ = 0$ the field ϕ thus becomes

$$\phi(x^+ = 0, x^-) = a_0 + \sum_{n>0} \frac{1}{\sqrt{4\pi n}} \left(a_n e^{-in\pi x^-/L} + a_n^* e^{in\pi x^-/L} \right), \quad (4.97)$$

Plugging this into the free part of the Lagrangian we get (discarding a total time derivative)

$$L_0[a_n, a_0] = -i \sum_{n>0} a_n \dot{a}_n^* - m^2 L a_0^2 - \sum_{n>0} \frac{m^2 L}{4\pi n} a_n^* a_n \equiv -i \sum_{n>0} a_n \dot{a}_n^* - H, \quad (4.98)$$

with H denoting the Hamiltonian. The Euler-Lagrange equations are

$$-i\dot{a}_n + \frac{m^2 L}{4\pi n} a_n = 0, \quad (4.99)$$

$$2m^2 L a_0 = 0, \quad (4.100)$$

where we have defined $\dot{a}_n = \partial a_n / \partial x^+$. The first equation, (4.99), is just the free Klein-Gordon equation which can be seen upon multiplying by k_n^+ . The second identity, (4.100), is non-dynamical and thus a *constraint* which states that a free field does not have a zero mode, $a_0 = 0$.

The method of Faddeev and Jackiw is based on equivalence of the Euler-Lagrange and Hamiltonian equations of motion. The latter are

$$\dot{a}_n = \{a_n, H\} = \sum_{k>0} \frac{m^2 L}{4\pi k} \{a_n, a_k^*\} a_k, \quad (4.101)$$

Obviously, this coincides with (4.99), if the canonical bracket is

$$\{a_k, a_n^*\} = -i\delta_{kn}. \quad (4.102)$$

The constraint (4.100) is obtained by differentiating the Hamiltonian,

$$\frac{\partial H}{\partial a_0} = 2m^2 L a_0 = 0. \quad (4.103)$$

Let us briefly show that the approach presented above is equivalent to Schwinger's [51]. From (4.98) we read off a generator

$$\delta G = -i \sum_{n>0} a_n \delta a_n^* \quad (4.104)$$

effecting the transformation

$$\delta a_n^* = \{a_n^*, \delta G\} = -i \sum_{k>0} \{a_n^*, a_k\} \delta a_k^*, \quad (4.105)$$

which in turn implies the canonical bracket (4.102).

Quantization is performed as usual by employing the correspondence principle, so that, from (4.102), the elementary commutator is given by

$$[a_m, a_n^\dagger] = \delta_{mn} . \quad (4.106)$$

The Fock space expansion for the (free) scalar field ϕ thus becomes

$$\phi(x^+ = 0, x^-) = \sum_{n>0} \frac{1}{\sqrt{4\pi n}} \left(a_n e^{-in\pi x^-/L} + a_n^\dagger e^{in\pi x^-/L} \right) , \quad (4.107)$$

Note that the Fock ‘measure’ $1/\sqrt{4\pi n}$ does not involve any scale like the mass m or the volume L . This is at variance with the analogous expansion in the instant form which reads

$$\phi(x, t = 0) = \frac{1}{\sqrt{2L}} \sum_n \frac{1}{\sqrt{2(k_n^2 + m^2)}} \left(a_n e^{ik_n x} + a_n^\dagger e^{-ik_n x} \right) , \quad (4.108)$$

where $-L \leq x \leq L$, $k_n = \pi n/L$, and $[a_n, a_m^\dagger] = \delta_{mn}$. Obviously, ϕ *does* depend on m and L . We will discuss some consequences of this difference in the next chapter.

We can use the result (4.107) to calculate the free field commutator at equal light-cone time x^+ ,

$$[\phi(x), \phi(0)]_{x^+=0} = \sum_{n \neq 0} \frac{1}{4\pi n} e^{-in\pi x^-/L} = -\frac{i}{2} \left[\frac{1}{2} \text{sgn}(x^-) - \frac{x^-}{2L} \right] . \quad (4.109)$$

which coincides with (4.35), (4.59) and (4.66). The commutator (4.109) has originally been obtained in [63] using the Dirac-Bergmann algorithm for constrained systems. The Faddeev-Jackiw method, however, is much more economic and transparent. In particular, it makes clear that the basic canonical variables of a light-cone field theory are the Fock operators or their classical counterparts. The a_n with, say, $-N \leq n \leq N$ in (4.96) can be viewed as defining a $(2N+1)$ -dimensional phase space. A phase space, however, should have even dimension. This is accomplished by choosing a polarization in terms of positions and momenta, here a_n and $a_n^\dagger$, with $n > 0$, and by the vanishing of the zero mode, $a_0 = 0$. We will later see that this vanishing is a peculiarity of the free theory.

Chapter 5

The Light-Cone Vacuum

5.1 Basics

One of the basic axioms of quantum field theory states that the spectrum of the four-momentum operator is contained within the closure of the forward light-cone (in momentum space) [70, 71]. The four-momentum P^μ of any physical, that is, observable particle thus obeys

$$P^2 \geq 0, \quad P^0 \geq 0, \quad (5.1)$$

which is, of course, consistent with the mass-shell constraint, $p^2 = m^2$. In other words, there are no tachyons having imaginary mass ($m^2 < 0$). The tip of the cone, the point $P^2 = P^0 = 0$, corresponds to the vacuum. From the spectrum condition (5.1) we infer that

$$P_0^2 - P_3^2 \geq P_\perp^2 \geq 0 \quad \text{or} \quad P^0 \geq |P^3|. \quad (5.2)$$

This implies for the longitudinal light-cone momentum,

$$P^+ = P^0 + P^3 \geq |P^3| + P^3 \geq 0. \quad (5.3)$$

We thus have the important kinematical constraint that physical states must have non-negative longitudinal momentum,

$$\langle \text{phys} | P^+ | \text{phys} \rangle \geq 0. \quad (5.4)$$

The spectrum of P^+ is thus bounded from below. Due to Lorentz invariance, the vacuum $|0\rangle$ must have vanishing four-momentum, and in particular

$$P^+ |0\rangle = 0. \quad (5.5)$$

Therefore, the vacuum is an eigenstate of P^+ with the lowest possible eigenvalue, namely zero. We will be interested in the phenomenon of spontaneous symmetry

breaking, i.e. in the question whether—roughly speaking—the vacuum is degenerate. We are thus going to analyze whether there is another state, $|p^+ = 0\rangle$, having the same eigenvalue, $p^+ = 0$, as the vacuum. If so, it must be possible to create this state from the vacuum with some operator U ,

$$|p^+ = 0\rangle = U|0\rangle, \quad (5.6)$$

where U must not produce any longitudinal momentum. Note that within ordinary quantization such a construction is straightforward and quite common, for example, in BCS theory. A state with vanishing three-momentum can be obtained via

$$|\mathbf{p} = 0\rangle = \int d^3k f(\mathbf{k}) a^\dagger(\mathbf{k}) a^\dagger(-\mathbf{k}) |0\rangle, \quad (5.7)$$

where f is an arbitrary wave function. Evidently, the contributions from modes with positive and negative momenta cancel each other. It is obvious as well, that within light-cone quantization things must be different as there cannot be an analogous cancelation for the longitudinal momenta which are always non-negative. Instead, one could imagine something like

$$|p^+ = \mathbf{p}_\perp = 0\rangle = \int_0^\infty dk^+ \int d^2k_\perp f(\mathbf{k}_\perp) \delta(k^+) a^\dagger(k^+, \mathbf{k}_\perp) a^\dagger(k^+, -\mathbf{k}_\perp) |0\rangle. \quad (5.8)$$

The problem thus boils down to the question whether there are Fock operators carrying light-cone momentum $k^+ = 0$. As we have seen in the preceding section there are no such operators, and a construction like (5.8) is impossible.

The only remaining possibility is that, if U contains a creation operator $a^\dagger(k^+ > 0)$ carrying longitudinal momentum $k^+ \neq 0$, there must be annihilators that annihilate exactly the same amount k^+ of momentum. Thus, after Wick ordering, U must have the general form

$$\begin{aligned} U = \langle 0|U|0\rangle &+ \int_{k^+ > 0} dk^+ f_2(k^+) a^\dagger(k^+) a(k^+) \\ &+ \int_{p^+ > 0} dp^+ \int_{k^+ > 0} dk^+ f_3(k^+, p^+) a^\dagger(p^+ + k^+) a(p^+) a(k^+) \\ &+ \int_{p^+ > 0} dp^+ \int_{k^+ > 0} dk^+ \tilde{f}_3(k^+, p^+) a^\dagger(p^+) a^\dagger(k^+) a(k^+ + p^+) \\ &+ \dots \end{aligned} \quad (5.9)$$

It follows that the light-cone vacuum $|0\rangle$ is an eigenstate of U ,

$$U|0\rangle = \langle 0|U|0\rangle |0\rangle. \quad (5.10)$$

As we only deal with rays in Hilbert space, the action of U on the vacuum does not create a state distinct from the vacuum. This result is actually rather general.

Any quantity that is obtained by integrating some functional of the fields over longitudinal space, i.e.,

$$F[\phi] = \int dx^- \mathcal{F}[\phi] , \quad (5.11)$$

is of the form (5.9), because the integration can be viewed as a projection onto the longitudinal momentum sector $k^+ = 0$. The most important examples for such quantities are the Poincaré generators, as is obvious from the representations (4.18, 4.19). This implies in particular that the trivial light-cone vacuum is an eigenstate of the fully interacting light-cone Hamiltonian P^- ,

$$P^-|0\rangle \equiv 2H|0\rangle = \langle 0|P^-|0\rangle|0\rangle . \quad (5.12)$$

This can be seen alternatively by considering

$$P^+P^-|0\rangle = P^-P^+|0\rangle = 0 , \quad (5.13)$$

which says that $P^-|0\rangle$ is a state with $k^+ = 0$, so that P^- must have a Fock representation like U in (5.9).

The actual value of $\langle 0|P^-|0\rangle$ is not important at this point as it only defines the zero of light-cone energy. Note that within the instant form the Fock or trivial vacuum is *not* an eigenstate of the full Hamiltonian as the latter usually contains terms with only creation operators where positive and negative three-momenta compensate to zero as in (5.7). The instant-form vacuum thus is unstable under time evolution. Such a vacuum, a typical example of which is provided by (5.7), is called ‘nontrivial’.

In the next section we perform an explicit comparison of nontrivial instant-form vacua and their trivial light-cone counterparts.

5.2 Trivial versus Nontrivial Vacua

Let us discuss some simple field theoretic models which are exactly solvable so that one can explicitly determine the ground state and the particle spectrum built on top of it.

5.2.1 Constant Source

Consider first a massive scalar field ϕ in $d = 1 + 1$ coupled to a constant external source ρ . We solve this model field theory both in the instant and front form. Using coordinates interpolating between the two forms the same model has been studied by Hornbostel [48]. We work in a finite spatial volume to circumvent any possible infrared problems. The Lagrangian density is

$$\mathcal{L} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}m^2\phi^2 + \rho\phi = \mathcal{L}_0 + \rho\phi . \quad (5.14)$$

Note that the external source explicitly breaks the reflection symmetry $\phi \rightarrow -\phi$ of the free Lagrangian $\mathcal{L}$. Upon completing the square, the Lagrangian (5.14) becomes

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 (\phi - \rho/m^2)^2 + \rho^2/2m^2 . \quad (5.15)$$

The minimum of the (classical) potential is thus located at $\phi = \rho/m^2 \equiv \phi_0$, which, in the quantum theory, corresponds to a non-vanishing vacuum expectation value,

$$\langle \Omega | \phi | \Omega \rangle = \phi_0 . \quad (5.16)$$

The state $|\Omega\rangle$ denotes a ‘non-trivial’ vacuum in the following sense. If we expand the field ϕ in Fock space like in (4.108),

$$\phi(x, t=0) = \frac{1}{\sqrt{2L}} \sum_n \frac{1}{\sqrt{2\omega_n}} (A_n e^{ik_n x} + A_n^\dagger e^{-ik_n x}) , \quad (5.17)$$

where $-L \leq x \leq L$, $k_n = \pi n/L$, $\omega_n^2 = k_n^2 + m^2$ and

$$[A_n, A_m^\dagger] = \delta_{mn} , \quad (5.18)$$

we must have

$$A_n |\Omega\rangle \neq 0 \quad (5.19)$$

for at least one n . The vacuum state $|\Omega\rangle$ must therefore contain particles corresponding to the A -modes of ϕ . If we denote the ‘trivial’ vacuum, i.e. the one annihilated by the A_n , as usual by $|0\rangle$, the two vacua must be related as in (5.7) via

$$|\Omega\rangle = U[A_n, A_n^\dagger] |0\rangle . \quad (5.20)$$

Let us determine the explicit form of U . To this end we note that the system under consideration is a field theoretic version of the shifted harmonic oscillator. If we define a translated field

$$\varphi \equiv \phi - \phi_0 , \quad (5.21)$$

this does not have an expectation value in $|\Omega\rangle$, as

$$\langle \Omega | \varphi | \Omega \rangle = \langle \Omega | \phi | \Omega \rangle - \phi_0 = 0 . \quad (5.22)$$

Thus, from the Fock expansion of φ

$$\varphi(x, t=0) = \frac{1}{\sqrt{2L}} \sum_n \frac{1}{\sqrt{2\omega_n}} (a_n e^{ik_n x} + a_n^\dagger e^{-ik_n x}) , \quad (5.23)$$

we infer that the a_n annihilate the non-trivial vacuum $|\Omega\rangle$ and thus correspond to quasi-particle operators built on the ground state $|\Omega\rangle$. Put differently, φ can be viewed as a free field with vacuum $|\Omega\rangle$, as is also obvious from the Klein-Gordon equation,

$$(\square + m^2)\phi = (\square + m^2)\varphi + \rho = \rho , \quad (5.24)$$

which implies that φ obeys a *free* equation of motion. The relation between the A - and a -modes is easily found. As ϕ and φ are related by a translation, we must have

$$\phi = U^\dagger \varphi U , \quad (5.25)$$

where U is the translation operator

$$U = \exp(-i\rho\Pi/m^2) = \exp(-i\phi_0\Pi) . \quad (5.26)$$

Accordingly, the operator U from (5.20) is determined in terms of the momentum Π conjugate to ϕ , which is the integral over the momentum density

$$\pi(x, t=0) = \frac{-i}{\sqrt{2L}} \sum_n \sqrt{\omega_n/2} (A_n e^{ik_n x} - A_n^\dagger e^{-ik_n x}) . \quad (5.27)$$

The momentum Π is thus essentially the zero mode of the density π ,

$$\Pi(t=0) = \int_{-L}^L dx \pi(x, t=0) = i\sqrt{mL} (A_0^\dagger - A_0) . \quad (5.28)$$

Putting everything together we obtain the explicit representation of U in terms of Fock operators,

$$U[A_n, A_n^\dagger] = \exp[\sqrt{mL} \phi_0 (A_0^\dagger - A_0)] . \quad (5.29)$$

Note that this quantity is entirely determined by the zero momentum modes $A_0^\dagger$ and A_0 . Plugging (5.29) into (5.20), one finds that $|\Omega\rangle$ is a coherent state of zero momentum modes,

$$|\Omega\rangle = \exp(-mL\phi_0^2/2) \exp(\sqrt{mL}\phi_0 A_0^\dagger) |0\rangle , \quad (5.30)$$

where we have used the Baker-Campbell-Hausdorff formula to factorize U . Being a coherent state, $|\Omega\rangle$ must be an eigenstate of the annihilators A_n . This is confirmed by a simple calculation yielding

$$A_{n \neq 0} |\Omega\rangle = 0 , \quad A_0 |\Omega\rangle = \sqrt{mL} \phi_0 |\Omega\rangle . \quad (5.31)$$

The non-vanishing vacuum expectation value thus arises from the action of the zero mode operators,

$$\langle \Omega | \phi | \Omega \rangle = \frac{1}{2L} \int_{-L}^L dx \langle \Omega | \phi | \Omega \rangle = \frac{1}{2\sqrt{mL}} \langle \Omega | A_0 + A_0^\dagger | \Omega \rangle = \phi_0 . \quad (5.32)$$

The quasi-particle operators are obtained by the unitary transformation (5.25),

$$\begin{aligned} a_n &= U A_n U^\dagger = A_n , & (n \neq 0) \\ a_0 &= U A_0 U^\dagger = A_0 - \sqrt{mL} \phi_0 , \end{aligned} \quad (5.33)$$

As might have been expected, only the zero modes get shifted through the action of U , namely by the c -number $-\sqrt{mL}\phi_0$. This transformation is a particular example of a canonical (or Bogolubov) transformation. It conserves the commutation relations (5.18), i.e.

$$[a_n, a_m^\dagger] = \delta_{mn} , \quad (5.34)$$

and diagonalizes the Hamiltonian, as is shown by the following calculation,

$$\begin{aligned} H &= \frac{1}{2} \int_{-L}^L dx \left[\pi^2 + \phi \left(-\frac{\partial^2}{\partial x^2} + m^2 \right) \phi \right] - \int_{-L}^L dx \rho \phi \\ &= \sum_n \omega_n A_n^\dagger A_n - \frac{\rho}{m} \sqrt{mL} (A_0 + A_0^\dagger) \\ &= \sum_n \omega_n a_n^\dagger a_n - (2L) \frac{\rho^2}{2m^2} . \end{aligned} \quad (5.35)$$

Obviously, the Hamiltonian becomes diagonal in the a -modes. In addition, we find a vacuum energy

$$E_{\text{vac}} \equiv \langle \Omega | H | \Omega \rangle = -(2L) \frac{\rho^2}{2m^2} , \quad (5.36)$$

which is consistent with the shift in energy density in (5.15). We thus have solved the problem entirely: the Hamiltonian is diagonal and its ground state as well as the excited states are determined. The entire effect of adding a constant source is a rearrangement of the vacuum state accompanied by a lowering of the ground state energy. The spectrum of excited states which is determined by the mass m (i.e. the curvature of the oscillator potential) remains unchanged.

Let us see now whether we can do the same within light-cone quantization. Again, we enclose the system in a finite spatial box, $-L \leq x^- \leq L$ and demand periodic boundary conditions in x^- . We perform the same decomposition of the fields,

$$\phi = \varphi + \phi_0 , \quad (5.37)$$

where ϕ_0 represents the vacuum expectation value, and look for the translation operator U relating ϕ and φ . The momentum (density) conjugate to ϕ is

$$\pi \equiv \partial^+ \phi . \quad (5.38)$$

We know from the discussion of the initial-value problem that π does not provide any new information: knowledge of ϕ implies knowledge of π as ∂^+ is a spatial derivative. Its zero mode is therefore determined through the spatial boundary conditions according to

$$\Pi = \int_{-L}^L dx^- \pi(x^-) = 2 \int_{-L}^L dx^- \frac{\partial}{\partial x^-} \phi(x^-) = 2[\phi(x^- = L) - \phi(x^- = -L)] = 0 , \quad (5.39)$$

and we note that it is vanishing. This has far-reaching consequences. First of all we note that U is the identity,

$$U = \exp(-i\phi_0\Pi) = 1 , \quad (5.40)$$

so that the vacuum does not change under action of U ,

$$|\Omega\rangle = |0\rangle , \quad (5.41)$$

nor do the Fock operators. How can this observation be reconciled with the decomposition (5.37), i.e. the existence of a non-vanishing vacuum expectation value ϕ_0 ? To this end we consider the Klein-Gordon equation for ϕ ,

$$(\square + m^2)\phi = (\partial^+ \partial^- + m^2)\phi = \rho , \quad (5.42)$$

where the external source ρ provides an inhomogeneous term. Upon integrating this over x^- the derivative term vanishes as in (5.39), and we are left with

$$\frac{1}{2L} \int_{-L}^L dx^- \phi(x^-) = \rho/m^2 \equiv \phi_0 . \quad (5.43)$$

which is nothing but a straightforward generalization of (4.100). The zero mode of ϕ thus coincides with the vacuum expectation value ϕ_0 . It is a c -number, entirely determined by the source (or ‘interaction’) ρ . This is at variance with ordinary quantization where we have

$$\frac{1}{2L} \int_{-L}^L dx \phi(x) = \frac{1}{2\sqrt{mL}}(A_0 + A_0^\dagger) \neq \phi_0 . \quad (5.44)$$

Here, the zero mode is *operator valued* and yields the vacuum expectation value only after acting on the non-trivial vacuum $|\Omega\rangle$ as is evident from (5.32).

The decomposition (5.37), when written in Fock space, becomes

$$\phi(x^-) = \sum_{n>0} \frac{1}{4\pi n} \left(a_n e^{ik_n^+ x^-/2} + a_n^\dagger e^{-ik_n^+ x^-/2} \right) + \phi_0 . \quad (5.45)$$

Obviously, the Fock operators in ϕ and φ are the same, and the zero mode of φ , which is a free field, vanishes in accordance with (4.100). As a result we conclude that, in order to determine the vacuum expectation value of ϕ , we have to calculate its zero mode which is constrained by the Euler-Lagrange equation of motion. This will be discussed exhaustively in Chapter 7.

5.2.2 Mass Shift: Bosons

Let us consider two free scalar fields, ϕ_0 and ϕ_1 with different masses, m_0 and m_1 [72]. We stay in $d = 1 + 1$ but go over to a continuum formulation. The Heisenberg fields obey the Klein-Gordon equations

$$(\square + m_i^2) \phi_i = 0, \quad i = 0, 1, \quad (5.46)$$

and have the instant-form Fock expansions

$$\phi(x, t) = \int \frac{dk}{2\pi} \frac{1}{\sqrt{2\omega_i(k)}} \left[a_i(k) e^{-ik_i \cdot x} + a_i^\dagger(k) e^{ik_i \cdot x} \right], \quad (5.47)$$

with $\omega_i(k) \equiv (k^2 + m_i^2)^{1/2}$, $k_i \cdot x \equiv \omega_i(k)t - kx$, and the commutation relations

$$[a_i(k), a_i^\dagger(p)] = 2\pi \delta(k - p). \quad (5.48)$$

It is important to note that the instant-form fields even at $t = 0$ (i.e. the Schrödinger fields) depend on the masses m_i both explicitly via ω_i and implicitly via the Fock operators a_i and $a_i^\dagger$. Accordingly, also the (Fock) vacua defined through

$$a_i(k)|\Omega_i\rangle = 0, \quad (5.49)$$

are mass dependent. This can be seen explicitly in the functional Schrödinger picture [73] in which the states are functionals $\Psi[\varphi] \equiv \langle \varphi | \Psi \rangle$ of the classical field φ , and the momentum operator $\pi(x)$ acts as the functional derivative $-i\delta/\delta\varphi(x)$. In this very intuitive formulation of quantum field theory, the free field vacua (5.49) are represented by Gaussian functionals [73],

$$\begin{aligned} \langle \varphi | \Omega_i \rangle &= \det^{1/4}(\omega_i/\pi) \exp \left[-\frac{1}{2} \int dx dy \varphi(x) \omega_i(x - y) \varphi(y) \right] \\ &= \det^{1/4}(\omega_i/\pi) \exp \left[-\frac{1}{2} \int \frac{dk}{2\pi} \varphi(-k) \omega_i(k) \varphi(k) \right], \end{aligned} \quad (5.50)$$

where the mass dependence resides in the ‘covariances’ ω_i . If we rewrite the Klein-Gordon equation for ϕ_1 as

$$(\square + m_0^2)\phi_1 = -(m_1^2 - m_0^2)\phi_1 \equiv \delta m^2 \phi_1, \quad (5.51)$$

we can equivalently view ϕ_1 as an interacting field with the interaction given by the Hamiltonian

$$H_{\text{int}} \equiv \frac{1}{2} \int dx \delta m^2 \phi^2. \quad (5.52)$$

The field ϕ_1 in the interaction picture, and the Heisenberg field ϕ_0 thus satisfy the same free field equation of motion. This implies that the associated fields in

the Schrödinger picture are the same. If we choose $t = 0$ as the time where the different dynamical pictures coincide, we have the common initial conditions,

$$\phi_0(x, t = 0) = \phi_1(x, t = 0) , \quad (5.53)$$

and analogously for the momenta π_i . This relates the Fock operators according to

$$a_1(k) = \alpha_k a_0(k) + \beta_k a_0^\dagger(-k) , \quad (5.54)$$

with the coefficients α_k and β_k given by

$$\alpha_k = \frac{1}{2} \left(\sqrt{\omega_1(k)/\omega_0(k)} + \sqrt{\omega_0(k)/\omega_1(k)} \right) , \quad (5.55)$$

$$\beta_k = \frac{1}{2} \left(\sqrt{\omega_1(k)/\omega_0(k)} - \sqrt{\omega_0(k)/\omega_1(k)} \right) . \quad (5.56)$$

As in the preceding subsection, we can interpret the $a_1(k)$ as the annihilators of quasi- or dressed particles which by some (unspecified) mechanism acquire an effective mass $m_1 \neq m_0$. Using the fact that $\alpha_k^2 - \beta_k^2 = 1$, one can define an angle θ_k such that

$$\cosh \theta_k \equiv \alpha_k , \quad \sinh \theta_k = \beta_k . \quad (5.57)$$

Therefore, the transformation (5.54) can be rewritten as

$$a_1(k) = a_0(k) \cosh \theta_k + a_0^\dagger(-k) \sinh \theta_k , \quad (5.58)$$

which is the standard Bogolubov transformation for bosons. As this transformation is unitary, it can equivalently be written with the help of a unitary operator U ,

$$a_1(k) = U a_0(k) U^\dagger , \quad (5.59)$$

given by

$$U \equiv U(\theta_k) = \exp \int \frac{dk}{4\pi} \theta_k \left[a_0(k) a_0(-k) - a_0^\dagger(k) a_0^\dagger(-k) \right] . \quad (5.60)$$

The vacuum $|\Omega_1\rangle$ is thus found to be

$$|\Omega_1\rangle = U |\Omega_0\rangle = \exp \int \frac{dk}{4\pi} \theta_k \left[a_0(k) a_0(-k) - a_0^\dagger(k) a_0^\dagger(-k) \right] |\Omega_0\rangle . \quad (5.61)$$

In quantum optics, such a state is called a ‘squeezed’ state [74]. It corresponds to a harmonic oscillator where the parabola representing the quasi-particle potential is ‘squeezed’ compared to the parabola of the particles (a_0 -modes). It is exactly this ‘squeezing’ which leads to a different curvature at the minimum thus implying a shift in the mass, $m_0 \rightarrow m$. From the point of view of the particles, the vacuum $|\Omega_1\rangle$ is non-trivial, as $a_0|\Omega_1\rangle \neq 0$.

Again, within light-cone quantization, the situation is drastically different. We have seen that the initial field at $x^+ = 0$ does not involve any scale, in particular no mass scale. As a consequence, both the light-cone Fock operators and the light-cone vacuum are mass independent [66, 75, 76]. Heuristically, one can argue that the front form is equivalent to a formulation in the ‘infinite-momentum frame’ [77] which is obtained from the instant form through the limit of a Lorentz boost with velocity $v \rightarrow c$ [78]. In this frame, all masses can be neglected compared to the longitudinal momentum, and thus should only play a minor role [76].

Technically, for the case at hand, the initial condition analogous to (5.53),

$$\phi_1(x^-, x^+ = 0) = \phi_0(x^-, x^+ = 0) , \quad (5.62)$$

implies that the a_0 - and a_1 -modes coincide so that U becomes the identity. Once again we thus find that the light-cone vacuum is not changed by the interaction, but remains trivial.

5.2.3 Mass Shift: Fermions

Consider finally the same situation for fermions, however, now in $d = 3 + 1$ [79]. Our starting point are the two Dirac equations

$$(i\not{\partial} - m_i) \psi_i = 0 , \quad i = 0, 1 . \quad (5.63)$$

We expand the Heisenberg fields as

$$\psi_i(x) = \sum_{\lambda} \int \frac{d^3k}{(2\pi)^{3/2}} \frac{1}{\sqrt{2\omega_i}} \left[b_i(\mathbf{k}, \lambda) u_i(\mathbf{k}, \lambda) e^{-ik_i \cdot x} + d_i^\dagger(\mathbf{k}, \lambda) v_i(\mathbf{k}, \lambda) e^{ik_i \cdot x} \right] , \quad (5.64)$$

with ω_i and k_i as in (5.47). The elementary spinors u and v are those of Itzykson and Zuber [80], multiplied by a factor $(2m_i)^{1/2}$ to allow for a straightforward massless limit. The Fock operators obey the canonical anti-commutation relations,

$$\left\{ b_i(\mathbf{k}, \lambda), b_i^\dagger(\mathbf{p}, \lambda) \right\} = \left\{ d_i(\mathbf{k}, \lambda), d_i^\dagger(\mathbf{p}, \lambda) \right\} = \delta_{\lambda\lambda'} \delta^3(\mathbf{k} - \mathbf{p}) . \quad (5.65)$$

As before, the Fock operators with subscripts $i = 0$ and $i = 1$ (corresponding to bare and dressed particles, respectively) are connected by a Bogolubov transformation,

$$b_1(\mathbf{k}, \lambda) = U b_0(\mathbf{k}, \lambda) U^\dagger , \quad d_1(\mathbf{k}, \lambda) = U d_0(\mathbf{k}, \lambda) U^\dagger , \quad (5.66)$$

which conserves the canonical anti-commutation relations. Like in the bosonic case, U is found from the common initial condition,

$$\psi_0(x, t = 0) = \psi_1(x, t = 0) . \quad (5.67)$$

A somewhat lengthy calculation involving a number of spinor matrix elements finally yields [79]

$$b_1(\mathbf{k}, \lambda) = \alpha_k b_0(\mathbf{k}, \lambda) + \beta_k d_0^\dagger(-\mathbf{k}, \lambda) , \quad (5.68)$$

$$d_1(\mathbf{k}, \lambda) = \alpha_k d_0(\mathbf{k}, \lambda) - \beta_k b_0^\dagger(-\mathbf{k}, \lambda) , \quad (5.69)$$

with coefficients

$$\alpha_k = \sqrt{\frac{1}{2}(1 + \gamma_k)} , \quad (5.70)$$

$$\beta_k = \sqrt{\frac{1}{2}(1 - \gamma_k)} , \quad (5.71)$$

$$\gamma_k = \frac{m_0 m_1 + k^2}{\omega_0 \omega_1} . \quad (5.72)$$

Obviously, we have $\alpha_k^2 + \beta_k^2 = 1$, so that this time there is an angle θ_k allowing to write

$$b_1(\mathbf{k}, \lambda) = b_0(\mathbf{k}, \lambda) \cos \theta_k + d_0^\dagger(-\mathbf{k}, \lambda) \sin \theta_k , \quad (5.73)$$

$$d_1(\mathbf{k}, \lambda) = d_0(\mathbf{k}, \lambda) \cos \theta_k - b_0^\dagger(-\mathbf{k}, \lambda) \sin \theta_k . \quad (5.74)$$

The unitary operator U is found to be

$$U = \exp \left\{ - \sum_{\lambda} \int d^3 k \theta_k \left[b_0(-\mathbf{k}, \lambda) d_0(\mathbf{k}, \lambda) + b_0^\dagger(\mathbf{k}, \lambda) d_0^\dagger(-\mathbf{k}, \lambda) \right] \right\} , \quad (5.75)$$

defining the nontrivial vacuum $|\Omega_1\rangle = U|\Omega_0\rangle$, with $b_0|\Omega_1\rangle, d_0|\Omega_1\rangle \neq 0$.

Upon initializing the fields on the null-plane $x^+ = 0$, however, things again become very different. The fermion fields ψ_i are decomposed into $\psi_i = \psi_{i,+} + \psi_{i,-}$, where only the $\psi_{i,+}$ are independent fields (see Section 4.3.5 and App. A). These obey a Klein-Gordon equation,

$$i\partial^- \psi_{i,+} = (i\partial^+)^{-1} (-\partial_\perp^2 + m_i^2) \psi_{i,+} , \quad i = 0, 1 , \quad (5.76)$$

which is obtained after eliminating the $\psi_{i,-}$. The (independent) Heisenberg fields have the Fock expansion

$$\psi_{i,+}(x) = \sum_{\lambda} \int_0^\infty \frac{dk^+}{k^+} \int \frac{d^2 k_\perp}{16\pi^3} \left[b_i(\mathbf{k}, \lambda) u_{i,+}(\mathbf{k}, \lambda) e^{-ik_i \cdot x} + d_i^\dagger(\mathbf{k}, \lambda) v_{i,+}(\mathbf{k}, \lambda) e^{ik_i \cdot x} \right] \quad (5.77)$$

where $\mathbf{k} = (k^+, \mathbf{k}_\perp)$ and

$$k_i \cdot x = \frac{1}{2} \hat{k}_i^- x^+ + \frac{1}{2} k^+ x^- - \mathbf{k}_\perp \cdot \mathbf{x}_\perp \equiv \frac{1}{2} \hat{k}_i^- x^+ + \mathbf{k} \cdot \mathbf{x} \quad (5.78)$$

with the $\hat{k}_i^-$ being on shell, $\hat{k}_i^- = (k_\perp^2 + m_i^2)/k^+$. Again, it is crucial to note that both the Fock operators and the spinors $u_{i,+}$ and $v_{i,+}$ are mass independent (see App. A). As a result, the common initial condition,

$$\psi_{0,+}(\mathbf{x}, x^+ = 0) = \psi_{1,+}(\mathbf{x}, x^+ = 0) , \quad (5.79)$$

leads to a trivial Bogolubov transformation with $U = \mathbb{1}$ [79, 81], like in the bosonic case. The light-cone vacuum, therefore, is stable under mass generation and the same for both fields $\psi_{i,+}$, namely the trivial Fock vacuum. This is corroborated by taking the infinite-momentum limit of the coefficient γ_k ,

$$\lim_{|\mathbf{k}| \rightarrow \infty} \gamma_k = 1, \quad (5.80)$$

which also makes the Bogolubov transformation (5.68, 5.69) trivial. In Chapters 8 and 9 we will discuss how one can obtain a fermion condensate in the trivial light-cone vacuum (see also [81]).

Chapter 6

Light-Cone Wave Functions

In this chapter we collect some basic facts about the eigenvalue problem of the light-cone Hamiltonian, or, in other words, about the light-cone Schrödinger equation and its solutions, the light-cone wave functions. We follow the discussion of Brodsky and Lepage [82] and also use their conventions. Accordingly, this chapter contains only few original results, but we think it worthwhile to include to make our presentation more self-contained.

6.1 Definition

Let us first stick to a discrete notation and, for the time being, stay in 1+1 dimensions. We thus have a Fock basis of states

$$\begin{aligned} |n\rangle &= a_n^\dagger |0\rangle, \\ |m, n\rangle &= a_m^\dagger a_n^\dagger |0\rangle, \\ &\vdots \\ |n_1, \dots, n_N\rangle &= a_{n_1}^\dagger \dots a_{n_N}^\dagger |0\rangle. \end{aligned} \tag{6.1}$$

This leads to a completeness relation defining the unit operator in Fock space,

$$\begin{aligned} \mathbb{1} &= |0\rangle\langle 0| + \sum_{n>0} |n\rangle\langle n| + \frac{1}{2} \sum_{m, n>0} |m, n\rangle\langle m, n| + \dots \\ &= |0\rangle\langle 0| + \sum_{N=1}^{\infty} \frac{1}{N!} \sum_{n_1, \dots, n_N>0} |n_1, \dots, n_N\rangle\langle n_1, \dots, n_N|. \end{aligned} \tag{6.2}$$

An arbitrary state $|\psi\rangle$ can thus be expanded as

$$|\psi\rangle = \sum_{n>0} \langle n|\psi\rangle |n\rangle + \frac{1}{2} \sum_{m, n>0} \langle m, n|\psi\rangle |m, n\rangle + \dots \tag{6.3}$$

The sums are such that the longitudinal momenta in each Fock sector add up to the total longitudinal momentum of $|\psi\rangle$. Note that the vacuum state does not contribute as it is orthogonal to any particle state, $\langle 0|\psi\rangle = 0$. The normalization of this state is obtained as

$$\begin{aligned}\langle\psi|\psi\rangle &= \sum_{n>0} |\langle n|\psi\rangle|^2 + \frac{1}{2} \sum_{m,n>0} |\langle m,n|\psi\rangle|^2 + \dots \\ &= \sum_{N=1}^{\infty} \frac{1}{N!} \sum_{n_1,\dots,n_N>0} |\langle n_1,\dots,n_N|\psi\rangle|^2.\end{aligned}\quad (6.4)$$

Let us assume that the state $|\psi\rangle$ corresponds to a bound-state obeying the light-cone Schrödinger equation (3.48),

$$(M^2 - \hat{M}_0^2)|\psi\rangle = \hat{W}|\psi\rangle. \quad (6.5)$$

We want to project this equation onto the different Fock sectors. For this we need the eigenvalues of the free invariant mass squared when applied to an N -particle state

$$|N\rangle \equiv |n_1, \dots, n_N\rangle. \quad (6.6)$$

We find that $\hat{M}_0^2$ is diagonal as expected,

$$\langle N|\hat{M}_0^2|N'\rangle = \langle N|P^+ \sum_{i=1}^N \frac{m^2}{k_{n_i}^+} |N'\rangle \equiv \sum_{i=1}^N \frac{m^2}{x_i} \langle N|N'\rangle \equiv M_N^2 \langle N|N'\rangle. \quad (6.7)$$

The light-cone Schrödinger equation thus becomes a system of coupled eigenvalue equations,

$$\begin{bmatrix} (M^2 - M_1^2) & \langle l|\psi\rangle \\ (M^2 - M_2^2) & \langle kl|\psi\rangle \\ \vdots & \end{bmatrix} = \begin{bmatrix} \langle l|W|m\rangle & \langle l|W|mn\rangle & \dots \\ \langle kl|W|m\rangle & \langle kl|W|mn\rangle & \dots \\ \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \langle l|\psi\rangle \\ \langle kl|\psi\rangle \\ \vdots \end{bmatrix}. \quad (6.8)$$

Clearly, this represents an infinite number of equations which in general will prove impossible to solve unless the matrix is very sparse and/or the matrix elements are small. The former condition is usually fulfilled as the interaction W in the light-cone Hamiltonian typically changes particle number at most by two¹. Assuming the matrix elements to be small amounts to dealing with a perturbative situation. This will be true for non-relativistic bound states of heavy constituents, but not for light hadrons which we are mainly interested in. We shall, however, encounter situations where the magnitude of the amplitudes decreases enormously with the

¹Note that terms with only creation operators are forbidden by k^+ -conservation. Still, in 1+1 dimensions, things can become messy as interactions with polynomials of arbitrary powers in ϕ are allowed [83]

particle number N , so that it is a good approximation to restrict to the lowest Fock sectors. In instant form field theory this has long been known as the Tamm-Dancoff method [84, 85].

Let us turn to the more realistic case of 3+1 dimensions in a continuum formulation. In the discussion of the Klein-Gordon equation we have seen that the solutions are of the form

$$\phi(x) = \int \frac{d^4 p}{2\pi^4} 2\pi \delta(p^2 - m^2) \theta(p^+) e^{-ip \cdot x} \chi(p) = \int_0^\infty \frac{dp^+}{16\pi^3 p^+} \int d^2 p_\perp e^{-i\hat{p} \cdot x} \chi(\hat{p}) , \quad (6.9)$$

where $\hat{p}$ denotes the on-shell four-momentum with $\hat{p}^2 = m^2$. From this expression we read off that the invariant normalization of a momentum eigenstate $|P^+, \mathbf{P}_\perp\rangle \equiv |\mathbf{P}\rangle$ is given by

$$\langle \mathbf{P} | \mathbf{K} \rangle = 16\pi^3 P^+ \delta^3(\mathbf{P} - \mathbf{K}) . \quad (6.10)$$

We already know that the bare Fock vacuum is an eigenstate of the interacting Hamiltonian. It thus serves as an appropriate ground state on top of which we can build a reasonable Fock expansion. If we specialize immediately to the case of QCD, we are left with the Fock basis states

$$\begin{aligned} |0\rangle , \\ |q\bar{q} : \mathbf{k}_i, \alpha_i\rangle &= b^\dagger(\mathbf{k}_1, \alpha_1) d^\dagger(\mathbf{k}_2, \alpha_2) |0\rangle , \end{aligned} \quad (6.11)$$

$$\begin{aligned} |q\bar{q}g : \mathbf{k}_i, \alpha_i\rangle &= b^\dagger(\mathbf{k}_1, \alpha_1) d^\dagger(\mathbf{k}_2, \alpha_2) a^\dagger(\mathbf{k}_3, \alpha_3) |0\rangle , \\ &\vdots \end{aligned} \quad (6.12)$$

In these expressions, $b^\dagger$, $d^\dagger$ and $a^\dagger$ create quarks q , antiquarks $\bar{q}$ and gluons g with momenta $\mathbf{k}_i$ from the trivial vacuum $|0\rangle$. The α_i denote all other relevant quantum numbers, like helicity, flavor and color.

In a more condensed notation we can thus describe, say, a pion with momentum $\mathbf{P} = (P^+, \mathbf{P}_\perp)$, as

$$|\pi(\mathbf{P})\rangle = \sum_{n, \lambda_i} \int \prod_i \frac{dx_i}{\sqrt{x_i}} \frac{d^2 k_{\perp i}}{16\pi^3} \psi_{n/\pi}(x_i, \mathbf{k}_{\perp i}, \lambda_i) \left| n : x_i P^+, x_i \mathbf{P}_\perp + \mathbf{k}_{\perp i}, \lambda_i \right\rangle , \quad (6.13)$$

where we have suppressed all discrete quantum numbers apart from the helicities λ_i . The integration measure takes care of the constraints (3.39) which the relative momenta in each Fock state (labeled by n) have to obey,

$$\prod_i dx_i \equiv \prod_i dx_i \delta\left(1 - \sum_j x_j\right) , \quad (6.14)$$

$$\prod_i d^2 k_{\perp i} \equiv 16\pi^3 \prod_i d^2 k_{\perp i} \delta^2\left(\sum_j \mathbf{k}_{\perp j}\right) . \quad (6.15)$$

As a mnemonic rule, we note that any measure factor $d^2k_{\perp i}$ is always accompanied by $1/16\pi^3$.

The most important quantities in (6.13) are the *light-cone wave functions*

$$\psi_{n/\pi}(x_i, \mathbf{k}_{\perp i}) \equiv \langle n : x_i P^+, x_i \mathbf{P}_{\perp} + \mathbf{k}_{\perp i}, \lambda_i | \pi(\mathbf{P}) \rangle, \quad (6.16)$$

which are the amplitudes to find n constituents with relative momenta $p_i^+ = x_i P^+$, $\mathbf{p}_{\perp i} = x_i \mathbf{P}_{\perp} + \mathbf{k}_{\perp i}$ and helicities λ_i in the pion. Due to the separation properties of the light-cone Hamiltonian the wave functions do not depend on the total momentum $\mathbf{P}$ of the pion. Applying (6.10) to the pion state (6.13), we obtain the normalization condition

$$\sum_{n, \lambda_i} \int \prod_i dx_i \frac{d^2k_{\perp i}}{16\pi^3} |\psi_{n/\pi}(x_i, \mathbf{k}_{\perp i}, \lambda_i)|^2 = 1. \quad (6.17)$$

The light-cone bound-state equation for the pion is a straightforward generalization of (6.8),

$$\begin{bmatrix} (M^2 - M_{q\bar{q}}^2) & \langle q\bar{q} | \pi \rangle \\ (M^2 - M_{q\bar{q}g}^2) & \langle q\bar{q}g | \pi \rangle \\ \vdots & \end{bmatrix} = \begin{bmatrix} \langle q\bar{q} | W | q\bar{q} \rangle & \langle q\bar{q} | W | q\bar{q}g \rangle & \dots \\ \langle q\bar{q}g | W | q\bar{q} \rangle & \langle q\bar{q}g | W | q\bar{q}g \rangle & \dots \\ \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \langle q\bar{q} | \pi \rangle \\ \langle q\bar{q}g | \pi \rangle \\ \vdots \end{bmatrix}. \quad (6.18)$$

If a constituent picture for the pion were true, its wave function would be entirely given by the projection $\langle q\bar{q} | \pi \rangle$ onto the valence state. All the higher Fock contributions would vanish and the unitarity sum (6.17) would simply reduce to

$$\sum_{\lambda\lambda'} \int_0^1 dx \int \frac{d^2k_{\perp}}{16\pi^3} |\psi_{q\bar{q}/\pi}(x, \mathbf{k}_{\perp}, \lambda, \lambda')|^2 = 1. \quad (6.19)$$

We will later discuss a model where this is indeed a good approximation to reality.

6.2 Properties

Let us rewrite the light-cone bound-state equation (6.8) by collecting all light-cone wave functions $\psi_n = \langle n | \psi \rangle$ into a vector Ψ ,

$$\Psi = \frac{W\Psi}{M^2 - M_0^2}. \quad (6.20)$$

From this expression it is obvious that all light-cone wave functions tend to vanish whenever the denominator

$$\epsilon \equiv M^2 - M_0^2 = M^2 - \left(\sum_i p_i \right)^2 = M^2 - \sum_i \frac{k_{\perp i}^2 + m_i^2}{x_i} \quad (6.21)$$

becomes very large. We know from Section 3.3 that this quantity measures how far off energy shell the total system, i.e. the bound-state is,

$$P^- - \sum_i p_i^- = \epsilon / P^+ . \quad (6.22)$$

For this reason, ϵ is sometimes called the ‘off-shellness’ [86, 87]. We thus learn from (6.20) that there is only a small overlap of the bound-state with Fock states that are far off shell. This implies the limiting behavior

$$\psi(x_i, \mathbf{k}_{\perp i}, \lambda_i) \rightarrow 0 \quad \text{for } x_i \rightarrow 0, k_{\perp i}^2 \rightarrow \infty . \quad (6.23)$$

These boundary conditions are related to the self-adjointness of the light-cone Hamiltonian and to the finiteness of its matrix elements. Analogous criteria have been used recently to relate wave functions for different Fock states n (via “ladder relations”, [88]) and to analyze the divergence structure of light-cone perturbation theory [89].

Suppressing spin, flavor and color degrees of freedom, a light-cone wave function will be a scalar function $\phi(x_i, \mathbf{k}_{\perp i})$ of the parameter ϵ . This is used for building models, the most common one being to assume a Gaussian behavior [90, 91, 92, 93],

$$\phi(x_i, \mathbf{k}_{\perp i}) = N \exp(-|\epsilon|/\beta^2) , \quad (6.24)$$

where β measures the size of the wave function in momentum space. Note, however, that a Gaussian ansatz is in conflict with perturbation theory which is the appropriate tool to study the high- $\mathbf{k}_{\perp}$ behavior and indicates a power decay of the *renormalized* wave function (up to possible logarithms, [82]). For the unrenormalized wave functions the boundary conditions (6.23) are violated unless one uses a cutoff as a regulator.

As the off-shellness ϵ is the most important quantity characterizing a light-cone wave function let us have a closer look by specializing to the simplest possible system, namely two bound particles of equal mass. One can think of this, for instance, as the valence wave function of the pion. The off-shellness becomes

$$\epsilon = M^2 - \frac{k_{\perp}^2 + m^2}{x(1-x)} = -\frac{1}{x(1-x)} \left[M^2(x - \frac{1}{2})^2 + \underbrace{\frac{4m^2 - M^2}{4}}_{\geq 0} + k_{\perp}^2 \right] . \quad (6.25)$$

The second term on the right-hand-side is positive because, for a bound state, the binding energy,

$$E = M - 2m , \quad (6.26)$$

is negative so that $2m > M$. As a result, the off-shellness is always negative. Only for free particles it is zero, because *all* momentum components (including the energy) sum up to the total momentum. In this case, each individual term in (6.25) vanishes,

$$M = 2m , \quad x = \frac{1}{2} , \quad \mathbf{k}_{\perp} = 0 . \quad (6.27)$$

It follows that the light-cone wave function of a free two-particle system (with equal-mass constituents) is of the form

$$\phi(x, \mathbf{k}_\perp) \sim \delta(x - 1/2) \delta^2(\mathbf{k}_\perp) . \quad (6.28)$$

One expects that for weak binding, in particular in the non-relativistic case, the wave functions will not look too differently, and thus will be highly peaked around $x = 1/2$ (in the equal mass case) and $\mathbf{k}_\perp = 0$. Let us check this explicitly. We go to the particle rest frame with $\mathbf{P} = 0$ or $P^+ = P^- = M$ and $\mathbf{P}_\perp = 0$, implying $\mathbf{p}_{\perp i} = \mathbf{k}_{\perp i}$. In this frame, the non-relativistic limit is defined by the following inequalities for the constituent masses and momenta (in ordinary instant-form coordinates),

$$p_i^0 - m_i \simeq \frac{\mathbf{p}_i^2}{2m_i} \ll |\mathbf{p}_i| \ll m_i . \quad (6.29)$$

The prototype systems in this class are of hydrogen type where we have for binding energy and r.m.s. momentum,

$$|E| = \langle \frac{\mathbf{p}^2}{2m} \rangle = \frac{m\alpha^2}{2} , \quad (6.30)$$

$$\langle p \rangle = m\alpha , \quad (6.31)$$

with $\alpha = 1/137$ the fine structure constant and m the reduced mass. In this case, the hierarchy (6.29) becomes

$$\frac{\alpha^2}{2} \ll \alpha \ll 1 , \quad (6.32)$$

which is fulfilled to a very good extent in view of the smallness of α .

Consider now the longitudinal momentum of the i^{th} constituent,

$$p_i^+ = p_i^0 + p_i^3 \simeq m_i + \frac{\mathbf{p}_i^2}{2m_i} + p_i^3 = x_i P^+ = x_i M . \quad (6.33)$$

We thus find that we should replace p_i^3 in instant-form non-relativistic wave functions by

$$p_i^3 = x_i M - m_i , \quad (6.34)$$

where we neglect terms of order $\mathbf{p}_i^2/m_i$. Let us analyze the consequences for the off-shellness. The latter is in ordinary coordinates

$$\epsilon = M^2 - M_0^2 = (M + \sum_i p_i^0)(M - \sum_i p_i^0) . \quad (6.35)$$

We thus need

$$\sum_i p_i^0 \simeq \sum_i m_i + \sum_i \frac{\mathbf{p}_i^2}{2m_i} = (M - E) + \sum_i \frac{\mathbf{p}_i^2}{2m_i} , \quad (6.36)$$

with $E = M - \sum m_i$, so that the off-shellness (6.35) becomes

$$\epsilon \simeq 2M \left(E - \sum_i \mathbf{p}_i^2 / 2m_i \right) \simeq 2M \left[E - \sum_i \frac{k_{\perp i}^2 + (Mx_i - m_i)^2}{2m_i} \right], \quad (6.37)$$

where we have performed the replacement (6.34) in the second identity. The light-cone wave functions will be peaked where the off-shellness is small, that is, for

$$x_i = m_i/M, \quad \text{and} \quad \mathbf{k}_{\perp i} = 0, \quad (6.38)$$

as expected from the non-interacting case.

As an explicit example we consider the ground state wave function of positronium, which in ordinary momentum space is given by the Fourier transform of $\psi(r) = N \exp(-m\alpha r)$. Performing (6.34) once more and normalizing to one, we obtain

$$\psi(x, \mathbf{k}_{\perp}) = \sqrt{\gamma^3/\pi} \frac{8\pi\gamma}{\left[k_{\perp}^2 + (xM - m_e)^2 + \gamma^2 \right]^2}, \quad (6.39)$$

where m_e is the electron mass and $\gamma = m\alpha$ the r.m.s. momentum. The result (6.39) is valid for small momenta, i.e. when $k_{\perp}^2, (xM - m_e)^2 \ll m_e^2$. It is obvious from (6.39) that the positronium wave function is sharply peaked around $x = m_e/M \simeq 1/2$ and $k_{\perp}^2 = 0$.

Part II

Applications

Chapter 7

Zero Modes and the Light-Cone Vacuum

Our first application will not (yet) deal with light-cone wave functions but rather with the issue of Chapter 5, the triviality of the light-cone vacuum. We will, however, be concerned with a non-trivial example of an interacting field theory, namely ϕ^4 -theory in 1+1 dimensions. As before we will address the question how a non-vanishing vacuum expectation value of the field ϕ can arise in the light-cone vacuum. As a further motivation we note that the dynamical system under consideration may be viewed as a low-dimensional caricature of the Higgs sector of the standard model which is also given by a ϕ^4 -type scalar field theory.

Historically, it was Weinberg in 1966, who realized that within “old-fashioned” Hamiltonian perturbation theory the infinite-momentum limit of many diagrams, in particular vacuum diagrams, is vanishing [94]. This limit was later on shown to be (at least perturbatively) equivalent to light-cone quantization [78, 77]. Within the context of current algebra [95, 96, 97] the success of this limit was traced to the fact that light-like charges always annihilate the vacuum, irrespective of whether they are conserved or not [66]. Thus, Coleman’s theorem [98], “the symmetry of the vacuum is the symmetry of the world”, does not apply [99]. As we have seen in Chapter 5, the Fock vacuum, i.e. the ground state of the free Hamiltonian, is stable under interaction. In the same manner as the light-like charges, the fully interacting Hamiltonian annihilates the vacuum, the formal reason being that both quantities conserve longitudinal momentum and thus are of the form (5.9).

With the advent of QCD as the theory of strong interactions, however, people began to feel somewhat uneasy. There was (and is) growing evidence, that many of the phenomenological aspects of hadron physics, like confinement and chiral symmetry breaking are related to the non-trivial features of the QCD vacuum (within standard instant-form quantization). Let us only mention features like quark- and gluon condensates, instantons, monopole condensation etc., which all indicate that the vacuum is densely populated by non-trivial quantum fluctua-

tions, which furthermore are not accessible to perturbation theory.

The concern that arose at these points can be put into the question: can the existence of these large vacuum fluctuations be reconciled with the triviality of the light-front vacuum? For QCD, the answer to this question is not (yet) known. For model field theories, some answers have already been found. Not surprisingly, it is the zero modes that are to some extent the carriers of non-trivial vacuum properties. One generally distinguishes between constrained and unconstrained zero modes. The latter typically arise in (low-dimensional) gauge theories [100, 101, 102, 103, 104, 105, 106]. The former we have already met in the discussion of the constant source (Subsection 5.2.1). For ϕ^4 -theory in 1+1 dimensions, it has been shown that the zero modes are responsible for spontaneous symmetry breaking of the reflection symmetry $\phi \rightarrow -\phi$ [107, 108, 109, 110, 111]. Let us review the basic features of what is going on [112].

7.1 The Constrained Zero Mode

We recall that the Lagrangian of a scalar field theory in 1+1 dimensions, written in terms of light-cone coordinates, cf. (4.95),

$$L[\phi, \partial^- \phi] = \frac{1}{2} \int dx^- \frac{1}{2} \partial^+ \phi \partial^- \phi - U[\phi] , \quad (7.1)$$

is linear in the velocity $\partial^- \phi$. According to the discussion of Chapter 4, the elementary bracket between the fields is essentially given by the inverse of the spatial derivative, $\partial^+ = 2\partial/\partial x^-$. In order to uniquely define this inverse, one has to specify the domain of ∂^- and boundary conditions. As before, we enclose our spatial variable x^- in a box, $-L \leq x^- \leq L$, and impose periodic boundary conditions on our fields. We know that in this case the operator ∂^+ has zero modes, namely all spatially constant functions. To deal with this aspect, we split our field ϕ into a zero mode ω and its complement φ ,

$$\phi(x^+, x^-) = \omega(x^+) + \varphi(x^+, x^-) , \quad (7.2)$$

$$\omega(x^+) \equiv \frac{1}{2L} \int_{-L}^L dx^- \phi(x^+, x^-) , \quad (7.3)$$

such that φ does *not* have a zero mode,

$$\int_{-L}^L dx^- \varphi(x^+, x^-) = 0 . \quad (7.4)$$

The Lagrangian (7.1) can then be rewritten as

$$L[\varphi, \omega] = \int_{-L}^L dx^- \frac{1}{2} \varphi (-2\partial_-) \dot{\varphi} - H[\varphi, \omega] . \quad (7.5)$$

For brevity, we use the derivative $\partial_- \equiv \partial/\partial x^-$ and $\dot{\varphi} \equiv \partial\varphi/\partial x^+$. Note that ω does not appear in the kinetic term. The Hamiltonian $H \equiv P^-$ coincides with the potential U from (7.1) after the replacement (7.2) has been performed. We thus find that the zero mode ω is constrained by its ‘equation of motion’,

$$\theta \equiv \frac{\delta H}{\delta \omega} = 0. \quad (7.6)$$

The basic bracket can be read off from the kinetic term in the Lagrangian (à la Schwinger or Faddeev/Jackiw, cf. Chapter 4),

$$\{\varphi(x^+, x^-), \varphi(x^+, y^-)\} \equiv -\frac{1}{2}\langle x^- | \partial_-^{-1} | y^- \rangle = -\frac{1}{2} \left[\frac{1}{2} \text{sgn}(x^- - y^-) - \frac{x^- - y^-}{2L} \right], \quad (7.7)$$

which, of course, coincides with (4.59) upon quantization via the correspondence principle.

As is well known, not all classical observables can be quantized unambiguously due to possible operator ordering problems [113, 114, 115, 116]. Such problems do not arise for the field φ and the bracket (7.7), where the field-independent right-hand-side leads to a c -number commutator. The constraint (7.6), however, implies a functional dependence of the zero mode ω on φ and thus a non-vanishing commutator of ω with φ . This can be explicitly verified by calculating the associated Dirac bracket within the Dirac-Bergmann algorithm [117, 118, 119]. For the quantum theory, it results in an ordering ambiguity with respect to ω and φ [107, 110]. Therefore, a definite ordering has to be prescribed. We chose Weyl (or symmetric) ordering [113, 120] which is explicitly hermitian. Using this prescription, one finds the quantum Hamiltonian for light-cone ϕ_{1+1}^4 -theory,

$$\begin{aligned} H = & \int_{-L}^L dx^- \left(\frac{1}{2} m^2 \varphi^2 + \frac{\lambda}{4!} \varphi^4 \right) + \\ & + \int_{-L}^L dx^- \left[\frac{1}{2} m^2 \omega^2 + \frac{\lambda}{4!} \left(\omega^4 + \omega \varphi^3 + \varphi \omega \varphi^2 + \varphi^2 \omega \varphi + \varphi^3 \omega + \right. \right. \\ & \left. \left. + \omega^2 \varphi^2 + \varphi^2 \omega^2 + \omega \varphi \omega \varphi + \varphi \omega \varphi \omega + \omega \varphi^2 \omega + \varphi \omega^2 \varphi \right) \right]. \quad (7.8) \end{aligned}$$

Note that we have chosen the sign of the mass term(s) in such a way that there is no spontaneous symmetry breaking at tree level as the classical potential does not have any turning points. With this Hamiltonian, equation (7.6) for the constrained zero mode ω reads explicitly

$$\theta = \frac{\delta H}{\delta \omega} = m^2 \omega + \frac{\lambda}{3!} \omega^3 + \frac{\lambda}{3!} \frac{1}{2L} \int_{-L}^L dx^- [\varphi^3 + \varphi^2 \omega + \varphi \omega \varphi + \omega \varphi^2] = 0. \quad (7.9)$$

This is nothing but the zero mode of the Euler-Lagrange equation of motion for the total field ϕ decomposed into ω and φ [63, 121, 110]. The remainder of this chapter is concerned with different approaches to solve this equation for ω .

7.2 Perturbative Solution

To obtain a perturbative solution for ω we expand it in a power series in λ ,

$$\omega \equiv \sum_{n=0}^{\infty} \lambda^n \omega_n \quad (7.10)$$

Inserting this into (7.9) determines the coefficients ω_n recursively. For the first three we find

$$\omega_0 = 0, \quad (7.11)$$

$$\omega_1 = -\frac{1}{6m^2} \frac{1}{2L} \int_{-L}^L dx^- \varphi^3(x), \quad (7.12)$$

$$\omega_2 = \frac{1}{36m^4} \frac{1}{(2L)^2} \int_{-L}^L dx^- dy^- \left[\varphi^2(x) \varphi^3(y) + \varphi(x) \varphi^3(y) \varphi(x) + \varphi^3(y) \varphi^2(x) \right] \quad (7.13)$$

All higher orders may be obtained similarly. Unfortunately, however, we have not been able to find a closed formula for ω_n which would allow for summing up the whole series (7.10).

If we expand the quantum field φ in terms of Fock operators, cf. (4.107),

$$\varphi(x) = \sum_{n=1}^{\infty} \frac{1}{\sqrt{4\pi n}} \left[a_n e^{-ik_n^+ x^-/2} + a_n^\dagger e^{ik_n^+ x^-/2} \right], \quad (7.14)$$

with the discretized longitudinal momentum $k_n^+ = 2\pi n/L$, one notes that the vacuum expectation value of ω is zero to all orders in λ , since ω_n contains an *odd* number of Fock operators, i.e. $\langle 0 | \omega_n | 0 \rangle = 0$. Thus

$$\langle 0 | \omega | 0 \rangle = \sum_{n=0}^{\infty} \lambda^n \langle 0 | \omega_n | 0 \rangle = 0. \quad (7.15)$$

This seems to imply that a non-vanishing vacuum expectation value for ω can only arise non-perturbatively and must be non-analytic in the coupling λ . We will discuss this issue in the next sections and continue for the time being within the framework of perturbation theory. In particular, we will study the effect

of the zero mode ω on the mass renormalization. To this end, we split up the Hamiltonian H from (7.8) into two pieces,

$$H = H_0 + H_\omega , \quad (7.16)$$

where H_0 is independent of ω ,

$$H_0 = \int_{-L}^L dx^- \left(\frac{1}{2} m^2 \varphi^2 + \frac{\lambda}{4!} \varphi^4 \right) , \quad (7.17)$$

and H_ω is the ω -dependent interaction given by the last two lines in (7.8). H_ω can be simplified using the constraint equation (7.9),

$$\begin{aligned} H_\omega &= H_\omega - \frac{L}{2} (\omega \theta + \theta \omega) \\ &= \frac{\lambda}{4!} \int_{-L}^L dx^- (-\omega^4 + \varphi \omega \varphi^2 + \varphi^2 \omega \varphi + \varphi \omega^2 \varphi - \omega \varphi^2 \omega) . \end{aligned} \quad (7.18)$$

Because ω is of order λ , the zero mode dependent part H_ω is of order λ^2 . Explicitly, we find, neglecting terms of order λ^3 ,

$$\begin{aligned} H_\omega &= \frac{\lambda}{4!} \int_{-L}^L dx^- (\varphi \omega \varphi^2 + \varphi^2 \omega \varphi) \\ &= -\frac{\lambda^2}{144 m^2 2L} \int_{-L}^L dx^- dy^- [\varphi(x) \varphi^3(y) \varphi^2(x) + \varphi^2(x) \varphi^3(y) \varphi(x)] , \end{aligned} \quad (7.19)$$

where we have used the first order term (7.12). This induces a mass shift of order λ^2 which is given by [119] (see also [108])

$$\delta m_n^2 \equiv \frac{2\pi n}{L} \left(\langle n | H_\omega | n \rangle - \langle 0 | H_\omega | 0 \rangle \right) . \quad (7.20)$$

Here, $|n\rangle \equiv a_n^\dagger |0\rangle$ denotes a one-particle state of longitudinal momentum $k_n^+ = 2\pi n/L$. We have subtracted the constant vacuum energy, $\langle 0 | H_\omega | 0 \rangle$, which diverges linearly with the volume. This is sufficient to render the mass shift finite. After inserting the Fock-expansion (7.14) into (7.19) one obtains for (7.20)

$$\begin{aligned} \delta m_n^2 &= -\frac{\lambda^2}{6m^2} \frac{L}{(4\pi)^3} \frac{1}{n} \left[\sum_{m=1}^{n-1} \frac{1}{(n-m)m} + 4 \sum_{m=1}^{\infty} \frac{1}{(n+m)m} \right] \frac{2\pi n}{L} \\ &= -\frac{\lambda^2}{6m^2} \frac{1}{16\pi^2 n} \left[3\gamma + \Psi(n) + 2\Psi(1+n) \right] < 0 , \end{aligned} \quad (7.21)$$

where Ψ denotes the Digamma-function [122], and $\gamma = -\Psi(1) = 0.577216\dots$ is Euler's constant. The question now is, whether the result (7.21) is a finite size effect which vanishes in the infinite-volume limit. The latter is obtained by replacing

$$\sum_{n=1}^N f(n) \rightarrow \lim_{L \rightarrow \infty} \frac{L}{2\pi} \int_{2\pi/L}^{2\pi N/L} dk^+ f(k^+ L/2\pi) \quad (7.22)$$

in such a way that k_n^+ approaches a finite limit k^+ , i.e. $n, L \rightarrow \infty$ with n/L finite. For the case at hand, this amounts to replacing the Digamma-function by its asymptotics [122]

$$\Psi(z) \simeq \log(z) - \frac{1}{2z} + O(1/z^2), \quad (z \rightarrow \infty \text{ in } |\arg(z)| < \pi). \quad (7.23)$$

Denoting the infinite-volume limit of the mass shift (7.20) by δm^2 , we finally obtain

$$\delta m^2 = -\frac{\lambda^2}{m^2} \frac{1}{16\pi k^+} \lim_{L \rightarrow \infty} \frac{1}{L} \left[\log(k^+ L/2\pi) + \gamma + O(1/k^+ L) \right] = 0. \quad (7.24)$$

The vanishing of this expression implies that the zero mode induced second order mass shift δm_n^2 is indeed a finite size effect. We do not have a general proof that this is also true for higher orders in λ . However, it seems to be plausible that a *single* mode like ω constitutes a “set of measure zero” within the infinite number of modes as long as one applies perturbation theory. From the theory of condensation, however, it is well known that single modes, especially those with vanishing momentum, can significantly alter the perturbative results. To analyze this possibility one clearly has to use non-perturbative methods. This will be done in the next sections.

7.3 Mean-Field Ansatz

As ϕ_{1+1}^4 -theory is super-renormalizable, the number of divergent diagrams is finite, namely one: the tadpole resulting from a self-contraction of two fields at the same space-time point. Therefore, renormalization can, at least within perturbation theory, be done via normal-ordering, which is nothing but making use of Wick's theorem: expanding the appearing powers of ϕ in a sum of normal-ordered terms with more and more self-contractions, one separates the finite term (with no contraction) from the divergent ones (with at least one contraction). The latter terms then are just the negative of the required counter-terms. For conventional ϕ^4 -theory one finds

$$\begin{aligned} \frac{1}{2} m^2 \phi^2 + \frac{\lambda}{4!} \phi^4 &= \frac{1}{2} m^2 \left(:\phi^2: + T \right) + \frac{\lambda}{4!} \left(:\phi^4: + 6 T :\phi^2: + 3 T^2 \right) \\ &= \frac{1}{2} \left(m^2 + \frac{\lambda}{2} T \right) :\phi^2: + \frac{\lambda}{4!} :\phi^4: + \frac{m^2}{2} T + \frac{\lambda}{8} T^2, \end{aligned} \quad (7.25)$$

where

$$T \equiv \langle \phi^2 \rangle = \int \frac{dk}{4\pi} \frac{1}{\sqrt{k^2 + m^2}} \quad (7.26)$$

formally denotes the logarithmically divergent tadpole contribution (in $d = 1+1$) which coincides with the vacuum expectation value of ϕ^2 or a self-contraction of the field.

It is obvious that only mass and vacuum energy get renormalized. The renormalized Hamiltonian is obtained by adding the counter-terms

$$\delta\mathcal{H} \equiv -\frac{\lambda}{4}T : \phi^2 : -\frac{1}{2}m^2T - \frac{\lambda}{8}T^2. \quad (7.27)$$

If the vacuum expectation value of ϕ^2 is taken in a Fock vacuum corresponding to the bare mass m , the latter coincides with the renormalized mass to order λ .

This rather trivial renormalization procedure cannot be straightforwardly extended to light-cone field theory, simply because we do not know the Hamiltonian! The zero mode ω is a complicated functional of the Fock operators $a_n, a_n^\dagger$, which has to be found from the constraint (7.9) before we can normal-order. There is, however, a way around this obstacle, such that an *exact* knowledge of ω is not needed for renormalization.

To this end we use the following general ansatz for ω ,

$$\begin{aligned} \omega &= \omega_0 + \sum_{n>0} \omega_n a_n^\dagger a_n + \sum_{m,n>0} \omega_{mn} a_m^\dagger a_n^\dagger a_{m+n} + \sum_{m,n>0} \omega_{mn}^* a_{m+n}^\dagger a_m a_n + \\ &+ \sum_{l,m,n>0} \omega_{lmn} a_{l+m+n}^\dagger a_l a_m a_n + \sum_{l,m,n>0} \omega_{lmn}^* a_l^\dagger a_m^\dagger a_n^\dagger a_{l+m+n} + \\ &+ \sum_{k,l,m,n>0} \delta_{k+l,m+n} \omega_{klmn} a_k^\dagger a_l^\dagger a_m a_n + \dots, \end{aligned} \quad (7.28)$$

where $\omega_0, \omega_n, \dots$ are c -numbers to be determined. Obviously, ω_0 is the vacuum expectation value of the operator ω ,

$$\omega_0 = \langle 0 | \omega | 0 \rangle, \quad (7.29)$$

whereas the coefficients ω_n etc. contribute to matrix elements of ω in higher Fock sectors. The ansatz (7.28) is hermitian and takes care of the fact that ω cannot transfer any momentum. Thus, it is a discrete analogue of (5.9) and can be understood as a Wick expansion written in the opposite of the usual order: the first term ω_0 is the sum of all contractions, the second the sum of all contractions but one and so on. Accordingly, each individual term in the expansion is a normal-ordered operator. Plugging the ansatz (7.28) into H_ω one notes that the latter may be viewed as an effective Hamiltonian where additional operators φ^n , $n \geq 2$ have been induced by the zero mode. This is reminiscent of a renormalization group analysis in which the same operators, being all marginal, are generated

by the renormalization group flow [123, 83]. In what follows, we will, however, truncate the ansatz (7.28) after the second term ($\sim a_n^\dagger a_n$) which only introduces additional φ^2 contributions and thus corresponds to a mean-field treatment.

Inserting the ansatz (7.28) into (7.8) and (7.9) we obtain for the constraint and the Hamiltonian

$$\theta = \theta_0 + \sum_{n>0} \theta_n a_n^\dagger a_n + \dots, \quad (7.30)$$

$$H = H_0 + \sum_{n>0} H_n a_n^\dagger a_n + \dots, \quad (7.31)$$

where, in accordance with the truncation of our ansatz (7.28), we have omitted terms containing more than two Fock operators. The coefficients H_0 , H_n and θ_0 , θ_n are functions of ω_0 and ω_n . Thus, (7.31) is an effective one-body or mean-field Hamiltonian describing the influence of the zero mode ω . Explicitly, one finds for the coefficients of the constraint,

$$\theta_0 = \left(m^2 + \frac{\lambda}{2}T\right) \omega_0 + \frac{\lambda}{3!} \left(\omega_0^3 + \sum_{n>0} \frac{\omega_n}{4\pi n}\right), \quad (7.32)$$

$$\theta_n = \left(m^2 + \frac{\lambda}{2}T\right) \omega_n + \frac{\lambda}{3!} \left(\omega_n^3 + 3\omega_0\omega_n^2 + 3\omega_0^2\omega_n + \frac{6\omega_0}{4\pi n} + \frac{6\omega_n}{4\pi n}\right), \quad (7.33)$$

and of the Hamiltonian (scaled by $2L$)

$$\begin{aligned} \frac{H_0}{2L} &= \frac{1}{2} \left(m^2 + \frac{\lambda}{2}T\right) \omega_0^2 + \frac{\lambda}{4!} \left(\omega_0^4 + 4 \sum_{n>0} \frac{\omega_0\omega_n}{4\pi n} + \sum_{n>0} \frac{\omega_n^2}{4\pi n}\right) \\ &+ \frac{m^2}{2}T + \frac{\lambda}{8}T^2, \end{aligned} \quad (7.34)$$

$$\begin{aligned} \frac{H_n}{2L} &= \frac{1}{2} \left(m^2 + \frac{\lambda}{2}T\right) \left(\omega_n^2 + 2\omega_0\omega_n + \frac{1}{2\pi n}\right) + \\ &+ \frac{\lambda}{4!} \left((\omega_0 + \omega_n)^4 - \omega_0^4 + \frac{3}{\pi n}(\omega_0 + \omega_n)^2 + \frac{\omega_n^2}{2\pi n}\omega_n \sum_{m>0} \frac{\omega_m}{\pi m}\right). \end{aligned} \quad (7.35)$$

T now denotes the light-cone tadpole (in discretized form)

$$T = \langle \varphi^2 \rangle = \sum_{n>0} \frac{1}{4\pi n}, \quad (7.36)$$

which is mass independent in contrast to (7.26) (unless one regularizes with a mass dependent cutoff, cf. Chapter 9). Note that the one-particle matrix elements of θ and H are given as the *sum* of two coefficients,

$$\langle n|\theta|n\rangle = \theta_0 + \theta_n = \left(m^2 + \frac{\lambda}{2}T\right) (\omega_0 + \omega_n)$$

$$+ \frac{\lambda}{3!} \left[(\omega_0 + \omega_n)^3 + 6 \frac{\omega_0 + \omega_n}{4\pi n} + \sum_{k>0} \frac{\omega_k}{4\pi k} \right], \quad (7.37)$$

and

$$\begin{aligned} \langle n|H/2L|n \rangle &= (H_0 + H_n)/2L = \\ &= \frac{1}{2} \left(m^2 + \frac{\lambda}{2} T \right) (\omega_0 + \omega_n)^2 + \frac{1}{2} \left(m^2 + \frac{\lambda}{2} T \right) \frac{1}{2\pi n} + \\ &+ \frac{\lambda}{4!} \left[(\omega_0 + \omega_n)^4 + (12\omega_0^2 + 24\omega_0\omega_n + 14\omega_n^2) \frac{1}{4\pi n} + \right. \\ &\quad \left. + 4(\omega_0 + \omega_n) \sum_{k>0} \frac{\omega_k}{4\pi k} + \sum_{k>0} \frac{\omega_k^2}{4\pi k} \right] + \\ &+ \frac{m^2}{2} T + \frac{\lambda}{8} T^2. \end{aligned} \quad (7.38)$$

From the divergence structure above it is clear that all coefficients can be made finite by adding the counter-term

$$\delta H/2L = -\frac{\lambda}{4} T \sum_{n=1}^{\infty} \frac{1}{2\pi n} a_n^\dagger a_n - \frac{\lambda}{4} T \omega^2 - \frac{1}{2} m^2 T - \frac{\lambda}{8} T^2, \quad (7.39)$$

which can be obtained from (7.27) by integrating over x^- and decomposing the field. The renormalization is thus standard, i.e. performed by normal-ordering and formally achieved by setting $T = 0$ in the expressions above.

It is convenient to rescale the coefficients

$$\omega_0 \rightarrow \frac{\omega_0}{\sqrt{4\pi}}, \quad (7.40)$$

$$\omega_n \rightarrow \frac{\omega_n}{\sqrt{4\pi}}, \quad (7.41)$$

and define a dimensionless coupling g as

$$g \equiv \frac{\lambda}{24\pi m^2} > 0, \quad (7.42)$$

such that (7.32 - 7.35) become

$$H_0/2L = \frac{m^2}{4\pi} \left[\frac{1}{2} \omega_0^2 + \frac{g}{4} \left(\omega_0^4 + 4\omega_0 \sum_{n=1}^{\infty} \frac{\omega_n}{n} + \sum_{n=1}^{\infty} \frac{\omega_n^2}{n} \right) \right], \quad (7.43)$$

$$\begin{aligned} H_n/2L &= \frac{m^2}{4\pi} \left[\frac{1}{2} \left(\omega_n^2 + 2\omega_0\omega_n + \frac{2}{n} \right) + \right. \\ &+ \left. \frac{g}{4} \left[(\omega_0 + \omega_n)^4 - \omega_0^4 + \frac{12}{n} (\omega_0 + \omega_n)^2 + 4\omega_n \sum_{k=1}^{\infty} \frac{\omega_k}{k} + 2\frac{\omega_n^2}{n} \right] \right] \end{aligned} \quad (7.44)$$

and

$$\omega_0 + g \left(\omega_0^3 + \sum_{n=1}^{\infty} \frac{\omega_n}{n} \right) = 0 , \quad (7.45)$$

$$\omega_n + g \left(\omega_n^3 + 3\omega_0\omega_n^2 + 3\omega_0^2\omega_n + \frac{6}{n}(\omega_0 + \omega_n) \right) = 0 . \quad (7.46)$$

This system of equations has a trivial solution

$$\omega_0 = \omega_n = 0 , \quad (7.47)$$

corresponding to the symmetric phase with vanishing vacuum expectation value of the field. Note further, that $\omega_n = 0$ implies $\omega_0 = 0$, which means that it is the *operator* part of ω that drives a possible phase transition. In the latter case, where $\omega_n \neq 0$, there are non-trivial solutions of (7.45, 7.46) which, however, cannot be obtained exactly. If we assume that there is a critical coupling where the field starts to develop a vacuum expectation value, then, very close to this coupling, ω_0 should be small so that we can expand ω_n as a power series in ω_0 [107, 112],

$$\omega_n = \alpha_n(g) \omega_0 + \beta_n(g) \omega_0^3 + O(\omega_0^5) . \quad (7.48)$$

Plugging this into (7.46), the expansion coefficients are determined recursively as,

$$\alpha_n = -\frac{6g}{n+6g} , \quad (7.49)$$

$$\begin{aligned} \beta_n &= -\frac{ng}{n+6g} \alpha_n (3 + 3\alpha_n + \alpha_n^2) \\ &= \frac{n}{6} \left(\frac{6g}{n+6g} \right)^2 \left[3 - \frac{18g}{n+6g} + \left(\frac{6g}{n+6g} \right)^2 \right] . \end{aligned} \quad (7.50)$$

Inserting ω_n into (7.45), one obtains ω_0 as a function of the coupling g via

$$a_0(g) + a_2(g) \omega_0^2 \equiv 1 + g \sum_{n>0} \frac{\alpha_n}{n} + g \left(1 + \sum_{n>0} \frac{\beta_n}{n} \right) \omega_0^2 = 0 . \quad (7.51)$$

The coefficients a_0 and a_2 can be evaluated analytically with the result

$$a_0(g) = 1 - g[\Psi(1+6g) + \gamma] , \quad (7.52)$$

$$a_2(g) = g \left[1 + 18g^2 [\Psi^{(1)}(1+6g) + 3g\Psi^{(2)}(1+6g) + 2g^2\Psi^{(3)}(1+6g)] \right] \quad (7.53)$$

where the $\Psi^{(n)}$ are Polygamma functions [122]. As β_n and therefore a_2 are positive, equation (7.51) develops two real solutions for ω_0 (with opposite sign) if

$$a_0(g) \leq 0 . \quad (7.54)$$

The critical coupling g_c is determined if equality holds. Using the explicit form of a_0 from (7.52), one finds

$$a_0(g_c) = 1 - g_c[\Psi(1 + 6g_c) + \gamma] = 0, \quad (7.55)$$

with a numerical value for the critical coupling of

$$g_c = 0.53070059 \dots, \quad (7.56)$$

which in terms of the original parameters is (cf. [124])

$$\lambda_c = 24\pi g_c m^2 \simeq 40.0 m^2. \quad (7.57)$$

Above this coupling, the vacuum expectation value ω_0 is non-vanishing and acquires one of two possible signs so that the reflection symmetry is spontaneously broken.

It is very illuminating to visualize the discussion above in terms of an effective potential V_{eff} [107, 125]. To this end we interpret (7.51), which determines the vacuum expectation value ω_0 , as the equation localizing the minima of V_{eff} . The latter is thus obtained via integration,

$$V_{\text{eff}}[\Omega] = \frac{1}{2}a_0(g)\Omega^2 + \frac{1}{4}a_2(g)\Omega^4. \quad (7.58)$$

The integration constant has been chosen such that $V_{\text{eff}}[\Omega = 0] = 0$. At the critical coupling, g_c , the coefficient a_0 of Ω^2 changes its sign from positive to negative. Therefore, below g_c , the effective potential has no turning points and its minimum is located at $\omega_0 = 0$ (symmetric phase). On the other hand, above g_c , V_{eff} develops two non-trivial minima at $\pm\omega_0$ (broken phase). As the order parameter ω_0 changes continuously, the phase transition is second order. Obviously, V_{eff} has precisely the form of a Landau-Ginzburg free energy which corroborates our mean-field interpretation. The final confirmation is provided by the calculation of the critical exponents. In a thermodynamic/magnetic analogy we view g as a temperature, Ω as a magnetization, and $\partial V_{\text{eff}}/\partial\Omega$ as an external magnetic field. The coefficient a_0 is then an inverse susceptibility which, near the critical coupling, should behave like $a_0 \sim (g - g_c)^\gamma$. (Evidently, the exponent γ has nothing to do with Euler's constant.) From (7.52) and (7.55) we obtain via Taylor expansion around g_c ,


$$a_0(g) \simeq (g - g_c) a'_0(g_c), \quad (7.59)$$

yielding the expected mean field result, $\gamma = 1$. The order parameter goes like $\omega \sim (g - g_c)^\beta$. Using (7.51) and (7.59), we find

$$\omega_0 = \pm [-a_0(g)/a_2(g)]^{1/2} \simeq \pm (g - g_c)^{1/2} [-a'_0(g_c)/a_2(g_c)]^{1/2}, \quad (7.60)$$

i.e. a mean field exponent $\beta = 1/2$. The dependence of the rescaled vacuum expectation value ω_0 on g is plotted in Fig. 7.1. From both (7.60) and the figure it is evident that ω_0 is non-analytic in the coupling g .

Figure 7.1: Behavior of the rescaled VEV ω_0 close to the critical coupling g_c . Graphs 1,2 and 3 refer to the expansion (7.48) of ω_n in ω_0 up to third, fifth and seventh order, respectively.



The last exponent, δ , is defined through the behavior of the magnetic field at g_c , $\partial V_{\text{eff}}/\partial\Omega \sim \Omega^\delta$. We calculate

$$\partial V_{\text{eff}}/\partial\Omega|_{g=g_c} = a_2(g_c) \Omega^3, \quad (7.61)$$

so that $\delta = 3$. As a cross-check, we note that our critical exponents obey the correct scaling law [123],

$$\delta = 1 + \frac{\gamma}{\beta}. \quad (7.62)$$

At this point it is instructive to make contact with perturbation theory. For this purpose, we expand $a_0(g)$ as given in (7.52) in powers of g using the series representation [122]

$$\Psi(1+z) = -\gamma + \sum_{n=2}^{\infty} (-1)^n \zeta(n) z^{n-1}, \quad (7.63)$$

from which we read off that $\Psi(1+z)$ is the generating function of the sequence $\zeta(n)$, ζ being the Riemann zeta function. The convergence region of the right-hand-side of (7.63) is $|z| < 1$. For $a_0(g)$ this yields,

$$a_0(g) = 1 - 6\zeta(2)g^2 + 36\zeta(3)g^3 + O(g^4), \quad (7.64)$$

which converges for $g < 1/6$. Keeping just the first two terms, one can find a critical coupling perturbatively [107, 125, 126] via

$$a_0(g_{cp}) = 1 - 6 \zeta(2) g_{cp}^2 = 0, \quad (7.65)$$

leading to

$$g_{cp} = 1/\pi = 0.3183098... > 1/6 \quad (7.66)$$

which is beyond the radius of convergence of (7.64). If we include the third order in g , the equation $a_0(g) = 0$ does no longer have a positive root like g_{cp} . We conclude that in the present framework, perturbation theory is not applicable for studying the critical behavior, and the value (7.66) for g_{cp} has to be abandoned. In other words, upon using perturbation theory one stays in the symmetric phase with ω_0 being zero, in agreement with the discussion of Section 7.2.

Finally, from the one-particle energies (7.44), we calculate the mass-shift induced by the non-perturbative zero mode ω as given by the ansatz (7.28),

$$\begin{aligned} \delta m_n^2 &= P_n^+ P_n^- - m^2 = \frac{2\pi n}{L} H_n - m^2 = m^2 \left[\frac{n}{2} (\omega_n^2 + 2\omega_0 \omega_n) + \right. \\ &\quad \left. + \frac{ng}{4} \left((\omega_0 + \omega_n)^4 - \omega_0^4 + \frac{12}{n} (\omega_0 + \omega_n)^2 + 4\omega_n \sum_{k=1}^{\infty} \frac{\omega_k}{k} + 2\frac{\omega_n^2}{n} \right) \right] \end{aligned} \quad (7.67)$$

Using the expression (7.48) for ω_n this becomes (near the critical coupling)

$$\begin{aligned} \delta m_n^2 &= 3g m^2 \omega_0^2 \left\{ 1 - 2 \left[1 - \frac{g}{1 + 6g/n} \left(\Psi(1 + 6g) + \gamma \right) \right] + \right. \\ &\quad \left. + 6g \frac{n + 7g}{(n + 6g)^2} \right\}. \end{aligned} \quad (7.68)$$

In contrast to the perturbative result (7.21), this expression is *not* vanishing in the continuum limit,

$$\delta m^2 = 3g m^2 \omega_0^2 \left[1 - 2a_0(g) \right]. \quad (7.69)$$

Due to (7.54) and (7.55), $a_0(g)$ is a small (negative) quantity. The mass shift induced by ω is thus positive. We want to emphasize that the zero mode ω has a non-trivial influence on the spectrum.

7.4 The Tamm-Dancoff Approximation

7.4.1 General Remarks

The Tamm-Dancoff approximation was originally designed within the conventional formulation of quantum field theory [84]. The principle of the method was to enormously reduce the particle number to a finite (and small) one. Due to

the complicated many-body structure of the equal-time vacuum, however, the approximation failed to lead to quantitative results and was abandoned thereafter. Although it has been noted rather early that light-cone field theory might be better suited for a Tamm-Dancoff approximation [127], it was not until recently that people began to start systematic investigations [128]. The hope was (and is), of course, that many of the problems of the original Tamm-Dancoff approximation can be avoided due to the simplicity of the light-cone vacuum. Specifically, people first tried to calculate bound states for 1+1 dimensional field theories where the method turned out to work extremely well (see Chapter 8 and [127, 129, 130, 131, 132]). Considerably more effort is needed in 3+1 dimensions [133, 134], the essential problem to solve being renormalization within a Hamiltonian formulation [135, 136, 137, 138, 139], as the structure of counter-terms as well as their required number is unclear. Furthermore, dynamical symmetries like rotational invariance or parity are violated in general [140]. Nevertheless, the first phenomenological applications have appeared recently [141, 142, 143].

In the following we will use the model at hand to shed some light on the issue of renormalization, in particular on the construction of the counter-terms needed. These could, in principle, be different for different particle number sectors ("sector-dependent renormalization" [128]). For a super-renormalizable theory like ϕ_{1+1}^4 , which conventionally is renormalizable by normal-ordering, this would be a rather undesired feature since it would complicate the renormalization procedure enormously. Therefore, in the following, we will try to keep the renormalization as simple as possible. To this end, we attempt to incorporate the normal-ordering prescription into the Tamm-Dancoff approximation. First we need some definitions. The projection operators on the lowest particle number states are

$$P_0 = |0\rangle\langle 0| \quad (7.70)$$

$$P_1 = \sum_{n>0} |n\rangle\langle n| \quad (7.71)$$

$$P_2 = \frac{1}{2} \sum_{n,m>0} |n, m\rangle\langle m, n| \quad (7.72)$$

$$\vdots$$

$$P_N = \frac{1}{N!} \sum_{n_1, \dots, n_N > 0} |n_1, \dots, n_N\rangle\langle n_1, \dots, n_N|, \quad (7.73)$$

with the Fock basis states defined in (6.1). Most important to us will be the Tamm-Dancoff projector

$$\mathbb{P}_N \equiv \sum_{\alpha=1}^N P_\alpha, \quad (7.74)$$

which projects onto the direct sum of all sectors with particle number less than or

equal to N . Before we perform any detailed calculation, let us make a few remarks about the relativistic invariance of the Tamm-Dancoff approximation [144].

Let $\mathcal{P}$ denote the Poincaré group. In d space-time dimensions the number of Poincaré generators is

$$\dim \mathcal{P} = \frac{d(d+1)}{2} \equiv D . \quad (7.75)$$

If the Poincaré generators are $G_0, G_1, \dots, G_{D-1}$, the Tamm-Dancoff approximation is relativistically invariant if

$$[\mathbb{P}_N G_i \mathbb{P}_N, \mathbb{P}_N G_k \mathbb{P}_N] = \mathbb{P}_N [G_i, G_k] \mathbb{P}_N . \quad (7.76)$$

It is easy to see, that this expression can only hold if at most one of the Poincaré generators fails to commute with $\mathbb{P}_N$. So we must have, e.g.

$$[G_0, \mathbb{P}_N] \neq 0 , \quad (7.77)$$

$$[G_k, \mathbb{P}_N] = 0 , \quad k = 1, \dots, D-1 . \quad (7.78)$$

The last identity implies that the G_k conserves particle number and therefore must be kinematical, i.e. interaction independent. Thus, G_0 must be the only dynamical generator. Clearly, this can only happen if the dimension of the stability group (of kinematical generators) is $D-1$. There is only one single case where this requirement is met, namely the front form in $d = 1+1$, which is the basis of the present discussion. Here, according to (7.75), there are only three generators; the dynamical generator is $G_0 \equiv P^-$, the light-cone energy; the kinematical generators are the momentum P^+ and the (longitudinal) boost M^{+-} . We think that this unique feature is one of the reasons, why the Tamm-Dancoff approximation (or more generally: Fock space truncation methods) work so well for light-cone field theories in $d = 1+1$.

Encouraged by this observations we continue and analyze the impact of the Tamm-Dancoff approximation on the quantum nature of the scalar field. This will be important for the issue of normal-ordering. We imagine that any operator $\mathcal{O}$ can be built from the elementary Fock operators $a_n, a_n^\dagger$, no matter how complicated its form may be. Thus, we *define* the N -particle Tamm-Dancoff approximation by the replacement

$$a_n \rightarrow \mathbb{P}_N a_n \mathbb{P}_N , \quad (7.79)$$

$$a_n^\dagger \rightarrow \mathbb{P}_N a_n^\dagger \mathbb{P}_N . \quad (7.80)$$

In this way, however, one is mutilating the quantum structure of the theory. This can be seen by calculating the N -particle Tamm-Dancoff approximation of the elementary commutator,

$$[a_m, a_n^\dagger]_N \equiv [\mathbb{P}_N a_m \mathbb{P}_N, \mathbb{P}_N a_n^\dagger \mathbb{P}_N] . \quad (7.81)$$

A somewhat tedious calculation leads to the final result [112]

$$[a_m, a_n^\dagger]_{\mathbf{N}} = \delta_{mn} \mathbb{P}_{\mathbf{N}-1} - P_N a_n^\dagger a_m P_N . \quad (7.82)$$

Taking matrix elements of this expression one readily sees that the correct result (δ_{mn}) for the commutator within N -particle Tamm-Dancoff approximation is reproduced only up to the $(N-1)$ -particle sector. The additional term on the right-hand-side of (7.82) is acting in the N -particle subspace only. In other words, matrix elements of expressions involving the elementary commutator, which are calculated within N -particle Tamm-Dancoff approximation, should not be trusted beyond the $(N-1)$ -particle sector. This will be relevant for the problem of normal-ordering to be discussed shortly.

To be a little bit more explicit, we list the lowest order expressions for (7.82)

$$[a_m, a_n^\dagger]_1 = \delta_{mn} |0\rangle \langle 0| - |n\rangle \langle m| , \quad (7.83)$$

$$[a_m, a_n^\dagger]_2 = \delta_{mn} \left[|0\rangle \langle 0| + \sum_{l>0} |l\rangle \langle l| \right] - \sum_{l>0} |l, n\rangle \langle m, l| . \quad (7.84)$$

What are now the implications of all that for the renormalization, in particular mass renormalization? As can be seen from (7.25) and (7.27), the latter is encoded in the normal-ordering prescription of the expression

$$\begin{aligned} \frac{1}{2L} \int_{-L}^L dx^- \varphi^2(x) &= \frac{1}{2L} \int_{-L}^L dx^- : \varphi(x)^2 : + T \\ &= \sum_{n>0} \frac{1}{2\pi n} a_n^\dagger a_n + \sum_{n>0} \frac{1}{4\pi n} [a_n, a_n^\dagger] . \end{aligned} \quad (7.85)$$

Normal-ordering thus amounts to splitting off the divergent tadpole contribution T , which, in the Fock space language, is given by an elementary commutator (or contraction). Thus the remarks above, leading to (7.82), apply. Let us calculate the N -particle Tamm-Dancoff approximation of (7.85),

$$\begin{aligned} \frac{1}{2L} \int_{-L}^L dx^- \varphi^2(x) \stackrel{N\text{PTDA}}{\simeq} &= \sum_{n>0} \frac{1}{2\pi n} \mathbb{P}_{\mathbf{N}-1} a_n^\dagger \mathbb{P}_{\mathbf{N}-1} a_n \mathbb{P}_{\mathbf{N}-1} + \mathbb{P}_{\mathbf{N}-1} T - \\ &- \sum_{n>0} \frac{1}{4\pi n} P_N a_n^\dagger a_n P_N . \end{aligned} \quad (7.86)$$

The same remarks as for (7.82) are in order. Matrix elements of the last expression should not be expected to be consistent beyond the $(N-1)$ -particle sector. Furthermore, one should note that, as φ^2 is a one-body operator, there is only *one* commutator (or contraction) involved in the above normal-ordering. For a k -body operator we therefore conjecture that its renormalization will be correct only up to the $(N-k)$ -particle sector. For example, in order to get the renormalization of the two-body operator φ^4 correct up to the one-particle sector, a three-particle Tamm-Dancoff approximation will be needed.

7.4.2 One-Particle Tamm-Dancoff Approximation

The one-particle light-cone Tamm-Dancoff approximation is defined by the replacements

$$a_n \simeq \mathbb{P}_1 a_n \mathbb{P}_1 = |0\rangle\langle n| \quad (7.87)$$

$$a_n^\dagger \simeq \mathbb{P}_1 a_n^\dagger \mathbb{P}_1 = |n\rangle\langle 0| . \quad (7.88)$$

From our general results we expect that within one-particle light-front Tamm-Dancoff approximation we will get a consistent renormalization of the one-body operator φ^2 in the vacuum (zero-particle) sector only.

To solve for the zero mode ω we make the following Tamm-Dancoff ansatz

$$\omega_{TD} = c_0 |0\rangle\langle 0| + \sum_{n>0} c_n |n\rangle\langle n| . \quad (7.89)$$

We similarly expand the constraint θ and the Hamiltonian H

$$\theta \simeq \bar{\theta}_0 |0\rangle\langle 0| + \sum_{n>0} \bar{\theta}_n |n\rangle\langle n| , \quad (7.90)$$

$$H \simeq \bar{H}_0 |0\rangle\langle 0| + \sum_{n>0} \bar{H}_n |n\rangle\langle n| , \quad (7.91)$$

where the bars simply indicate the distinction of the coefficients above from those of the mean field ansatz (7.30) and (7.31). The coefficients can be found as functions of c_0, c_n upon inserting the ansatz (7.89) into the constraint (7.9) and the Hamiltonian (7.8) yielding

$$\bar{\theta}_0 = \left(m^2 + \frac{\lambda}{3} T \right) c_0 + \frac{\lambda}{3!} \left(c_0^3 + \sum_{n>0} \frac{c_n}{4\pi n} \right) , \quad (7.92)$$

$$\bar{\theta}_n = \left(m^2 + \frac{\lambda}{12\pi n} \right) c_n + \frac{\lambda}{3!} \left(c_n^3 + \frac{c_0}{4\pi n} \right) , \quad (7.93)$$

and

$$\begin{aligned} \frac{\bar{H}_0}{2L} &= \frac{1}{2} \left(m^2 + \frac{\lambda}{4} T \right) c_0^2 + \frac{\lambda}{4!} \left(c_0^4 + 2 \sum_{n>0} \frac{c_0 c_n}{4\pi n} + \sum_{n>0} \frac{c_n^2}{4\pi n} \right) \\ &+ \frac{m^2}{2} T + \frac{\lambda}{4!} T^2, \end{aligned} \quad (7.94)$$

$$\begin{aligned} \frac{\bar{H}_n}{2L} &= \frac{1}{2} \left(m^2 + \frac{\lambda}{16\pi n} \right) c_n^2 + \frac{1}{2} \left(m^2 + \frac{\lambda}{12} T \right) \frac{1}{4\pi n} \\ &+ \frac{\lambda}{4!} \left(c_n^4 + \frac{c_0 c_n}{2\pi n} + \frac{c_0^2}{4\pi n} \right). \end{aligned} \quad (7.95)$$

At a first look, these expressions for the coefficients appear to be a disaster: the infinities in form of the tadpole T do not appear systematically, the mass gets renormalized in the vacuum sector only, but differently for the constraint θ and the Hamiltonian H . Both expressions differ from the standard expression $m^2 + \lambda T/2$. There is also a divergent contribution from the φ^4 term to $\bar{H}_0$ which differs from the usual $\lambda T^2/8$ of (7.25). As stated above, we do not believe the coefficients $\bar{\theta}_n$ and $\bar{H}_n$ to be correct within one-particle light-cone Tamm-Dancoff approximation. They will be discussed in the next subsection, when we go to higher order.

The way to remedy the situation (for the coefficients $\bar{\theta}_0$ and $\bar{H}_0$) is the following. We insist, firstly, on the standard mass renormalization, $m^2 + \lambda T/2$, however, according to our general discussion, in the vacuum sector only. Secondly, we do not assume that the coefficients c_0 and c_n are independent, and use this freedom to *redefine* them in the following way

$$c_0 \equiv \omega_0 , \quad (7.96)$$

$$c_n \equiv \omega_0 + \omega_n . \quad (7.97)$$

Inserting this into (7.92) and (7.94) one finds

$$\bar{\theta}_0 = \left(m^2 + \frac{\lambda}{2} T \right) \omega_0 + \frac{\lambda}{3!} \left(\omega_0^3 + \sum_{n>0} \frac{\omega_n}{4\pi n} \right) , \quad (7.98)$$

$$\begin{aligned} \bar{H}_0/2L &= \frac{1}{2} \left(m^2 + \frac{\lambda}{2} T \right) \omega_0^2 + \frac{\lambda}{4!} \left(\omega_0^4 + 4 \sum_{n>0} \frac{\omega_0 \omega_n}{4\pi n} + \sum_{n>0} \frac{\omega_n^2}{4\pi n} \right) + \\ &+ \frac{m^2}{2} T + \frac{\lambda}{4!} T^2 . \end{aligned} \quad (7.99)$$

Remarkably, the simple redefinition (7.97) has led to the desired results. The mass renormalization is standard and the same for θ_0 and H_0 . The divergences thus can be made to vanish by adding the counter-term (7.39). Both equations (7.98) and (7.99) coincide with the lowest order results from the mean-field ansatz (7.32) and (7.34) (up to the constant φ^4 -contribution to H_0 given by $\lambda T^2/4!$). Note that there are no two-body (T^2) contributions to the constraint. This is obviously true to all orders, so the renormalization of θ is slightly simpler than that of H , namely just mass renormalization.

The coincidence with the mean field results is not accidental. If one calculates the lowest order matrix elements of the mean field ansatz (7.28), one finds

$$\omega_0 = \langle 0 | \omega_{MF} | 0 \rangle = \langle 0 | \omega_{TD} | 0 \rangle = c_0 , \quad (7.100)$$

$$\omega_0 + \omega_n = \langle n | \omega_{MF} | n \rangle = \langle n | \omega_{TD} | n \rangle = c_n . \quad (7.101)$$

Analogous relations hold for the matrix elements of the constraint and the Hamiltonian,

$$\langle 0 | \theta | 0 \rangle = \theta_0 = \bar{\theta}_0 , \quad (7.102)$$

$$\langle 0|H|0\rangle = H_0 = \bar{H}_0 , \quad (7.103)$$

$$\langle n|\theta|n\rangle = \theta_0 + \theta_n = \bar{\theta}_n , \quad (7.104)$$

$$\langle n|H|n\rangle = H_0 + H_n = \bar{H}_n . \quad (7.105)$$

Thus, after the redefinition (7.97), the zero- and one-particle matrix elements of ω calculated within the mean field ansatz and Tamm-Dancoff approximation coincide. As the renormalization within mean field ansatz was conventional and straightforward, it is not too surprising that the behavior of the redefined Tamm-Dancoff approximation under renormalization gets improved. This will be another guideline in the following.

7.4.3 Two-Particle Tamm-Dancoff Approximation

If we now go one step further and include also two-particle states via

$$a_n \simeq \mathbb{P}_2 a_n \mathbb{P}_2 = |0\rangle\langle n| + \sum_{m>0} |m\rangle\langle m, n| \quad (7.106)$$

$$a_n^\dagger \simeq \mathbb{P}_2 a_n^\dagger \mathbb{P}_2 = |n\rangle\langle 0| + \sum_{m>0} |n, m\rangle\langle m| , \quad (7.107)$$

we should further improve our renormalization program. We expect a consistent renormalization of φ^2 -contributions in the vacuum- and one-particle sector, and of φ^4 -contributions in the vacuum sector. The extended ansatz for ω becomes

$$\begin{aligned} \omega_{TD} = & c_0 |0\rangle\langle 0| + \sum_{n>0} c_n |n\rangle\langle n| + \\ & + \frac{1}{2} \sum_{m,n>0} c_{mn} |m, n\rangle\langle m+n| + \frac{1}{2} \sum_{m,n>0} c_{mn}^* |m+n\rangle\langle m, n| + \\ & + \frac{1}{4} \sum_{k,l,m,n>0} \delta_{k+l,m+n} c_{klmn} |k, l\rangle\langle m, n| . \end{aligned} \quad (7.108)$$

It is now very plausible (though we cannot prove it a priori) that a consistent renormalization requires a redefinition also of the coefficients c_{mn} , c_{mn}^* , and c_{klmn} . To proceed as before, we would need the matrix elements $\bar{\theta}_0$, $\bar{\theta}_n$, $\bar{H}_0$, and $\bar{H}_n$ with all two-particle contributions in order to just get a consistent renormalization up to the one-particle sector. This is very tedious and inefficient, as we are *only* interested in the *divergent* contributions from higher order terms to lower order matrix elements. Fortunately, there is an alternative: we simply demand that the matrix elements of ω_{TD} and ω_{MF} coincide also in the two-particle sector. This gives us the desired redefinitions, namely

$$c_{mn} = 2\omega_{mn} , \quad (7.109)$$

$$c_{mn}^* = 2\omega_{mn}^* , \quad (7.110)$$

$$c_{klmn} = (\omega_0 + \omega_n + \omega_n)(\delta_{km}\delta_{ln} + \delta_{kn}\delta_{lm}) + 4\delta_{k+l,m+n} \omega_{klmn} . \quad (7.111)$$

Thus, essentially, only the two-particle matrix elements c_{klmn} get redefined. Expression (7.108) becomes

$$\begin{aligned}\omega_{TD} = & \omega_0|0\rangle\langle 0| + \sum_{n>0}(\omega_0 + \omega_n)|n\rangle\langle n| + \\ & + \sum_{m,n>0} \omega_{mn}|m,n\rangle\langle m+n| + \sum_{m,n>0} \omega_{mn}^*|m+n\rangle\langle m,n| + \\ & + \frac{1}{2} \sum_{m,n>0} (\omega_0 + \omega_m + \omega_n)|m,n\rangle\langle m,n| + \sum_{k,l,m,n>0} \delta_{k+l,m+n} \omega_{klmn} |k,l\rangle\langle m,n| .\end{aligned}$$

The diagonal two-particle term in the last line above contributes to the equations determining the coefficients ω_0 and ω_n and crucially alters the renormalization behavior. In [110], it was noted that our mean field ansatz amounts to including two-particle matrix elements, and it is just these terms that we have now explicitly displayed. Omitting them, as was done in [110], leads to an uncanceled tadpole contribution and thus to a spurious logarithmic divergence.

If we neglect all terms containing ω_{mn} , ω_{mn}^* , and ω_{klmn} , which are not of interest within two-particle Tamm-Dancoff approximation, we find for the constraint

$$\bar{\theta}_0 = \left(m^2 + \frac{\lambda}{2}T\right)\omega_0 + \frac{\lambda}{3!} \left(\omega_0^3 + \sum_{n>0} \frac{\omega_n}{4\pi n}\right), \quad (7.112)$$

$$\bar{\theta}_n = \left(m^2 + \frac{\lambda}{2}T\right)(\omega_0 + \omega_n) + \frac{\lambda}{3!} \left[(\omega_0 + \omega_n)^3 + 6\frac{\omega_0 + \omega_n}{4\pi n} + \sum_{k>0} \frac{\omega_k}{4\pi k}\right] \quad (7.113)$$

and for the Hamiltonian

$$\begin{aligned}\bar{H}_0/2L = & \frac{1}{2} \left(m^2 + \frac{\lambda}{2}T\right) \omega_0^2 + \frac{\lambda}{4!} \left[\omega_0^4 + 4 \sum_{n>0} \frac{\omega_0 \omega_n}{4\pi n} + \sum_{n>0} \frac{\omega_n^2}{4\pi n}\right] + \\ & + \frac{m^2}{2}T + \frac{\lambda}{8}T^2, \quad (7.114)\end{aligned}$$

$$\begin{aligned}\bar{H}_n/2L = & \frac{1}{2} \left(m^2 + \frac{\lambda}{2}T\right) (\omega_0 + \omega_n)^2 + \frac{1}{2} \left(m^2 + \frac{\lambda}{3}T\right) \frac{1}{2\pi n} + \\ & + \frac{\lambda}{4!} \left[(\omega_0 + \omega_n)^4 + \left[12(\omega_0 + \omega_n)^2 + 2\omega_n^2\right] \frac{1}{4\pi n} + \right. \\ & \quad \left. + 4(\omega_0 + \omega_n) \sum_{k>0} \frac{\omega_k}{4\pi k} + \sum_{k>0} \frac{\omega_k^2}{4\pi k}\right] + \\ & + \frac{m^2}{2}T + \frac{\lambda}{4!}T^2. \quad (7.115)\end{aligned}$$

Several remarks are in order. As θ does not contain two-body components, the renormalization is correct up to the one-particle sector. (7.112) and (7.113) thus

coincide with (7.32) and (7.37). The coefficient θ_0 is not even changed by including the two-particle contributions as can be seen by comparing with (7.98). In the vacuum coefficient $\bar{H}_0$, the mass renormalization (due to the φ^2 contributions) and the vacuum energy $m^2T/2 + \lambda T^2/8$ (with the T^2 contribution stemming from the φ^4 -term) are correct, as expected. So $\bar{H}_0$ is consistently renormalized. In the one-particle coefficient $\bar{H}_n$, which should be compared with (7.38), only the mass renormalization in the ω -sector is correct, as this is due to one-body contributions like $\omega^2\varphi^2$. As anticipated, mass renormalization and vacuum energy stemming from the φ^4 -term differ from the correct values by numerical factors. To get these correctly, one would have to perform a three-particle Tamm-Dancoff approximation. Presumably, this would only change the coefficients of the divergent terms, whereas the coefficients of the finite terms would remain the same. This would then be analogous to the change in $\bar{H}_0$ by going from one-particle Tamm-Dancoff approximation (7.99) to two-particle Tamm-Dancoff approximation (7.114).

Summarizing, we can say that, in order to obtain a consistent renormalization within a N -particle light-cone Tamm-Dancoff approximation, one has to include contributions from $(N + 1)$ -particle matrix elements by appropriately redefining the coefficients in the Tamm-Dancoff ansatz. In this way, the Fock ansatz method (7.28) and the Tamm-Dancoff approximation become completely equivalent.

7.5 Discussion

In this chapter we have analyzed the vacuum structure of light-cone ϕ_{1+1}^4 -theory by comparing different methods of solving for the constrained zero mode of the field operator. Within perturbation theory, the zero mode induces a second order mass correction which is vanishing in the infinite volume limit. We believe that to all orders in perturbation theory the zero mode only induces finite size effects, although we do not have a general proof.

We have presented two non-perturbative methods to obtain a solution for the zero mode. An ansatz in terms of an increasing number of Fock operators, which we have truncated after the one-body term, seems to be the most economic procedure. With considerably more efforts, exactly the same results can be obtained within a light-cone Tamm-Dancoff approximation, if the renormalization procedure is properly designed.

With either method we find a non-vanishing vacuum expectation value ϕ_c of the field if the coupling λ exceeds a critical value of $\lambda_c \simeq 40 m^2$, implying spontaneous breakdown of the reflection symmetry $\phi \rightarrow -\phi$. As the vacuum expectation value changes continuously, the associated phase transition is of second order, which has been rigorously established for the model at hand [145]. The order parameter ϕ_c , for instance, shows a square-root behavior as a function of the coupling, so that the associated critical exponent is $\beta = 1/2$. The critical behavior is thus of mean field type, which is only qualitatively correct,

as the ϕ_{1+1}^4 model is in the universality class of the two-dimensional Ising model; thus, β should be $1/8$. Because the calculations tend to become very messy, it is difficult to say, whether this shortcoming can be removed if one extends our approximations to higher orders. It is, however, plausible that the critical behavior will change if more terms in the ansatz (7.28) are retained as these correspond to marginal operators φ^n with $n > 2$. This issue, of course, deserves further investigations.

Another problem we have to face is the absence of any volume dependence of the phase transition. We have been working in a finite spatial volume of length $2L$, the length scale L , however, which does not appear in the Fock measure, drops out of the equation (7.55) determining the critical coupling. On the other hand, there cannot be a phase transition in a finite volume due to topological fluctuations (kinks and anti-kinks) which have non-vanishing statistical weight for $L < \infty$. We have not incorporated these fluctuations by our choice of (periodic) boundary conditions, so it is perhaps not too astonishing that we do not obtain a volume dependence. It should, in addition, be pointed out that the light-cone spatial volume, $-L \leq x^- \leq L$, is of a very peculiar nature. It corresponds to a light-like Minkowski region, the *invariant* length of which vanishes. This may give rise to features that are not present if the volume is space-like [47].

A better understanding of this problem is certainly desirable. As a first step towards doing better, a pure continuum formulation of the problem has been developed recently that leads to a slightly different critical coupling [146], namely

$$g_c = 0.69812669 \dots \quad (7.116)$$

Note that in [146] a different coupling normalization, $\tilde{g} \equiv 6g$, is used so that $\tilde{g}_c = 4.18876012 \dots$. The value (7.116) can be straightforwardly obtained from our equation (7.55) if the continuum limit is performed as in Section 7.2, i.e. via replacing the Digamma function by its asymptotics (7.23). Instead of (7.55), we find,

$$1 - g_c \log(6g_c) = 0, \quad (7.117)$$

which is solved by (7.116). This agreement of two independent calculations provides additional confidence in our method. We also point out that the critical coupling (7.116) is not too different from the one obtained within sophisticated renormalization group treatments [83], which, using our normalization, is given by $g_{c,RG} = 0.84 \pm 0.11$ [146].

Chapter 8

't Hooft and Schwinger Model

In the last chapter we have seen that zero modes carry at least some of the vacuum structure for a *scalar* field theory. Unfortunately, the methods used there cannot be straightforwardly generalized to the case of fermionic fields. There, a non-trivial vacuum shows up through the appearance of a fermion condensate, which in QCD, for example, signals spontaneous chiral symmetry breaking, as we have discussed in the introduction,. The point now is, that a fermion condensate is bilinear in the fields, and we do not expect fermionic zero modes to play any particular role in the formation of the condensate. This is also supported by the fact that a single fermionic zero mode cannot condense as it does not have vacuum quantum numbers. We will therefore follow a different route in this chapter and try to reconstruct vacuum properties from the particle spectrum. This is where light-cone wave functions enter the stage.

8.1 Prelude: Fermion Condensates

Assume that we have a symmetry which is explicitly broken so that the associated current is not conserved,

$$\partial_\mu j^\mu(x) = A(x) . \quad (8.1)$$

With the help of the corresponding Ward identity [147] one can derive the following formula for an arbitrary operator B ,

$$\langle 0 | \delta B / \delta \alpha | 0 \rangle = -i \int d^4x \langle 0 | T A(x) B(0) | 0 \rangle = - \sum_n \frac{\langle 0 | A(0) | n \rangle \langle n | B(0) | 0 \rangle}{m_n^2} . \quad (8.2)$$

Here, $\delta B / \delta \alpha$ denotes the change of B under the symmetry transformation, and in the last step a complete set of states $|n\rangle$, each of mass m_n , has been inserted. For the case of chiral symmetry, choosing

$$A = B = 2m\bar{\psi}i\gamma_5\psi , \quad (8.3)$$

one finds for a single quark flavor of mass m

$$\langle 0 | \bar{\psi} \psi | 0 \rangle = -m \sum_n \frac{|\langle 0 | \bar{\psi} i \gamma_5 \psi | n \rangle|^2}{m_n^2} . \quad (8.4)$$

In [148, 149], this expression was used as a definition for the quark condensate in the 't Hooft model (see the next sections). From the above derivation, however, it is clear that (8.4) holds quite generally. Using, for instance, the PCAC relation for the axial vector current, j_5^μ ,

$$\partial_\mu j_5^\mu(x) = f_\pi m_\pi^2 \pi(x) , \quad (8.5)$$

where $\pi(x)$ is an interpolating pion field and f_π the pion decay constant, (8.2) involves the pion two-point function and is easily evaluated with the result

$$f_\pi^2 m_\pi^2 = -4m \langle 0 | \bar{\psi} \psi | 0 \rangle . \quad (8.6)$$

This is the famous Gell-Mann-Oakes-Renner relation [150] (to lowest order in the quark mass m [151]), which relates the QCD parameters, m , the current quark mass, and the fermion condensate to the observables f_π and m_π . We have written everything in terms of bare quantities since the right-hand side does not change under renormalization [152]. Thus, m_π in (8.6) is the physical pion mass.

The Gell-Mann-Oakes-Renner relation can of course also be derived from (8.4) if one replaces

$$\bar{\psi}(x) i \gamma_5 \psi(x) = \frac{1}{2m} f_\pi m_\pi^2 \pi(x) , \quad (8.7)$$

and assumes that, for small m_π , the sum is saturated by the pion.

In any case, we want to stress the fact that in (8.4) and (8.6) above a vacuum quantity, the condensate, is expressed in terms of the particle spectrum. So, once the spectrum is known, after, say, diagonalizing the light-cone Hamiltonian by one of the various methods on the market, we can translate back properties of the spectrum into properties of the vacuum. In view of that, we suggest to deemphasize the role of the vacuum, which is natural to the extent that most of its properties are not directly observable. This is particularly true within the light-cone framework, where the vacuum state seems to decouple completely from the particle states. Similar ideas have been put forward long ago, in the context of chiral symmetry in the (light-cone) parton model, by Susskind et al. [153]: “In this framework the spontaneous symmetry breakdown must be attributed to the properties of the hadron’s wave function and not to the vacuum” [154]. A related point of view has also been taken more recently in [47].

Before we pursue the program just outlined we would like to remark that the ‘master equation’ (8.2) cannot be derived by strictly sticking to the light-cone framework. To this end note that the first term in (8.2) can be written with the help of the charge

$$Q(x^0) = \int d^3x j^0(x^0, \mathbf{x}) , \quad (8.8)$$

the generator of the symmetry, as

$$\langle 0 | \delta B / \delta \alpha | 0 \rangle = -i \langle 0 | [B, Q] | 0 \rangle, \quad (8.9)$$

where the commutator is evaluated at equal time $x^0 = 0$. This expression cannot be directly translated into the light-cone language by replacing the ordinary charge $Q(x^0)$ by the light-cone or “light-like” charge,

$$Q(x^+) = \int dx^- d^2 x_\perp j^+(x^+, x^-, \mathbf{x}_\perp), \quad (8.10)$$

and evaluating the commutator at equal *light-cone* time. The reason for this is a peculiar property of light-like charges which was briefly mentioned in the introduction to Chapter 7: they annihilate the vacuum, irrespective of whether they generate a symmetry or not [66, 155], which is in accordance with the triviality of the light-cone vacuum. Thus, the right-hand side of (8.9), evaluated on a null-plane, is always zero. Hence, that part of the operator $\delta B / \delta \alpha$ having a non-vanishing vacuum expectation value cannot be obtained by an infinitesimal transformation generated by the light-like charge Q . For example, in the light-cone sigma model, the relation

$$[\pi, Q_5] = -i\sigma \quad (8.11)$$

does only hold for those modes of the field operators, π and σ , having non-vanishing light-cone three-momenta, $(p^+, \mathbf{p}_\perp) \neq 0$. These non-zero modes do not have a vacuum expectation value. Thus, the vacuum expectation value of (8.11) consistently vanishes on both sides, as it should.

The moral is that we have to *assume* the validity of the identity (8.2) for any possible choice of quantization hypersurface, in particular for null planes. Accordingly, we can use a complete set of eigenstates of the *light-cone* Hamiltonian in the last term of (8.2), as was done in [148, 149]. This amounts to solving the light-cone Schrödinger equation for the corresponding light-cone wave functions.

In the following, we will test the ideas outlined above for two particular models due to 't Hooft and Schwinger, respectively. Our presentation is a slightly modified version of [156].

8.2 State of the Art

The number of papers on the Schwinger [157] and 't Hooft [127, 158] models is legion – for a twofold reason. On the one hand, the two models are particularly simple and become, for special choices of parameters, even exactly solvable. On the other hand, despite their simplicity, both models contain interesting physics analogous or similar to properties of gauge theories in higher dimensions. The

models therefore serve as interesting theoretical laboratories for studying phenomena like confinement and chiral symmetry breaking, to mention only the most prominent ones. For recent reviews on the two models we refer the reader to [159] for the Schwinger model and to [160] for the 't Hooft model and generalizations thereof.

While confinement (or charge screening) is realized in the same way in both models, via a linearly rising Coulomb potential, the second feature, chiral symmetry breaking, is not. In the 't Hooft model, 1+1 dimensional QCD in the limit of large N_C , chiral symmetry is 'almost' spontaneously broken [161, 162] and a *massless* bound state arises in the chiral limit of vanishing quark mass [127, 158]. In the massive Schwinger model, QED in $d = 1+1$ [163, 164], chiral symmetry is anomalously broken, and the contribution of the anomaly to the mass gap survives the chiral limit yielding the free, *massive* boson of the massless model. This boson becomes interacting in the massive model and again can be viewed as a bound state of fermionic constituents. For the sake of brevity we will call the lowest bound state of both models the 'pion' (although, in the real world of $d = 3+1$, the Schwinger model boson rather corresponds to the η' -meson). In this chapter we will try to determine its mass and (light-cone) wave function as accurately as possible. We mention in passing that the Schwinger model chiral anomaly is closely related to the appearance of a vacuum θ -angle parameter [165, 166] which also affects the particle spectrum [164]. Throughout this chapter, however, we implicitly assume that θ is set to zero.

The calculation of bound state masses and wave functions for the two models at hand has a long history, beginning with the exact solution of the massless Schwinger model [157]. For non-vanishing 'electron' mass, m , the model is no longer exactly solvable and one has to resort to approximations. One of them is to assume the mass m being small and expand around the massless solution [163, 164]. For the 'pion' mass squared one expects an expansion of the form

$$M^2(m) = 1 + M_1 m + M_2 m^2 + M_3 m^3 + O(m^4) , \quad (8.12)$$

where all masses are measured in units of the basic scale $\mu_0 = e/\sqrt{\pi}$, the mass of the boson in the massless Schwinger model, which is thus represented by the '1' in (8.12). The first order coefficient M_1 was obtained analytically in [167] via bosonization,

$$M_1 = 2e^\gamma = 3.56215 . \quad (8.13)$$

with $\gamma = 0.577216$ being Euler's constant. Shortly afterwards Bergknoff, using light-cone Hamiltonian techniques (see below), found a value [129]

$$M_1 = 2\pi/\sqrt{3} = 3.62760 , \quad (8.14)$$

which differs from (8.13) by 1.8 %. One topic of this chapter will be the analysis of this discrepancy and an attempt to do better by refining 't Hooft's and Bergknoff's

methods. This is particularly important as the coefficient M_1 is directly related to the vacuum structure of the Schwinger model by [168]

$$M_1 = -4\pi \langle 0 | \bar{\psi} \psi | 0 \rangle \cos \theta , \quad (8.15)$$

where $\langle 0 | \bar{\psi} \psi | 0 \rangle$ is the condensate of the massless Schwinger model [169, 170] (in units of μ_0),

$$\langle 0 | \bar{\psi} \psi | 0 \rangle = -\frac{1}{2\pi} e^\gamma = -0.28347 , \quad (8.16)$$

and θ the vacuum θ -angle. The result (8.13) thus corresponds to $\theta = 0$. In [99], the relation (8.15) (for $\theta = 0$), which is a 1+1 dimensional analogue of the Gell-Mann-Oakes-Renner formula (8.6), has been used to determine the condensate from the ‘pion’ mass squared via

$$\langle 0 | \bar{\psi} \psi | 0 \rangle = -\frac{1}{4\pi} \frac{\partial}{\partial m} M^2(m) \Big|_{m=0} = -\frac{1}{4\pi} M_1 . \quad (8.17)$$

Any inaccuracy in the determination of M_1 thus immediately affects the value of the condensate [45, 46]¹.

Recently, the calculations of $M^2(m)$ have been extended to second order. Using functional integral techniques and mass perturbation theory, Adam [175] found the analytical expression

$$M_2 = 4\pi^2 \langle 0 | \bar{\psi} \psi | 0 \rangle^2 (A \cos 2\theta + B) , \quad (8.18)$$

where $A = -0.6599$ and $B = 1.7277$ are numerical constants, given in terms of integral expressions containing the ‘pion’ propagator for $m = 0$. Inserting the values for A , B and the condensate, and setting $\theta = 0$, (8.18) becomes

$$M_2 = 3.3874 . \quad (8.19)$$

This result has been confirmed independently by Fields et al. [176], who derived the same integral expressions by summing up all relevant Feynman diagrams in the bosonized theory using near-light-cone coordinates.

To first order in m , the mass-squared of the ‘t Hooft model ‘pion’ has already been calculated by ‘t Hooft [158] by solving the associated light-cone bound-state equation. He derived this equation by projecting the covariant Bethe-Salpeter equation onto three-dimensional hypersurfaces of equal light-cone time, x^+ . Soon afterwards, the equation was rederived using light-cone Hamiltonian techniques [177]. The result for the ‘pion’ mass-squared is

$$M^2(m) = 2 \frac{\pi}{\sqrt{3}} m + O(m^2) . \quad (8.20)$$

¹ At this point it should be mentioned that within light-cone quantization there have been many attempts to calculate the condensate of the massless model alternatively by solving for its vacuum structure [171, 100, 101, 105, 172, 106, 173, 174].

As the light-cone bound state equations of the two models at hand differ only by an additive contribution due to the anomaly (see [99, 178] and below), it is not too surprising that 't Hooft's and Bergknoff's results coincide. For the 't Hooft model, all masses are expressed in units of the basic scale $\mu_0^2 = g^2 N_C / 2\pi^2$. It is obvious from (8.20) that M^2 vanishes in the chiral limit, $m \rightarrow 0$. As explained in [161, 162], this is not in contradiction with Coleman's theorem [179] as the 'pion' is not a Goldstone boson and decouples from the S -matrix.

The condensate of the 't Hooft model has first been calculated by Zhitnitsky [162] using basically the identity (8.4),

$$\langle 0 | \bar{\psi} \psi | 0 \rangle / N_C = -\frac{1}{4\pi} M_1 = -0.28868. \quad (8.21)$$

As expected, the condensate is proportional to N_C and can again be written in terms of M_1 . The result has been confirmed analytically [180, 47] and numerically [181]. Via (8.21), the numerical value [181] for the condensate leads to a numerical estimate for M_1 ,

$$M_1 = 3.64 \pm 0.05. \quad (8.22)$$

Though the numerical calculation did not use light-cone methods, it seems to favor the 't Hooft-Bergknoff value (8.14), which is *the* standard value for the 't Hooft model. Higher order corrections to (8.20) have been discussed in [182] without explicit calculation of the expansion coefficients.

In recent years, both models have been serving as a testing ground for new techniques developed in order to solve bound state problems using light-cone (or, equivalently, light-front) quantization. These new methods are 'discretized light-cone quantization' (DLCQ) [63, 64, 65], where one works in a finite volume leading to an equally-spaced momentum grid (cf. Chapters 4 and 7), and the 'light-front Tamm-Dancoff approximation' [128], which (drastically) truncates the Fock space of the theory (cf. Section 7.4), thus limiting the number of constituents a bound state can have. The latter approach can be viewed as a generalization of the techniques of 't Hooft and Bergknoff. Both methods aim at a numerical solution of relativistic bound state problems. DLCQ has been applied to the massive Schwinger model [183] and to QCD in 1+1 dimensions (for arbitrary N_C) [178, 184]. There are analogous LFTD calculations for both models, QCD₁₊₁ [185, 186] and the Schwinger model [187, 188]. These latter works are closer in spirit to ours than the DLCQ approaches. We will compare to all this recent work in more detail later on.

The purpose of this chapter is to obtain the 'pion' mass squared of both the Schwinger and 't Hooft model to high accuracy including the third order in m . In addition, we want to calculate the light-cone wave function of the 'pion' with high precision. We will use the light-front techniques of 't Hooft and Bergknoff

²Some authors (including 't Hooft) use a different convention for the coupling, which amounts to the replacement $g^2 \rightarrow 2g^2$.

and extensions thereof. Our starting point is a Tamm-Dancoff truncation to the two-particle (valence) sector. The low order calculations will be done analytically. To go beyond, computer algebraic and numerical methods will be applied and their convergence tested. The final goal is to shed some light on the advantages and limitations of this particular approach to bound-state equations.

The remainder of this chapter is organized as follows. In Section 8.3 we review 't Hooft's ansatz for the wave function yielding the lowest order solution of the bound-state equation. We compare the exact endpoint analysis with a variational procedure which is introduced at this point. In Section 8.4 we extend 't Hooft's ansatz by adding more variational parameters in such a way that an analytic solution can still be obtained. To this end we employ computer algebraic methods which allow to treat up to five variational parameters. This is sufficient to achieve rather good convergence. In Section 8.5, these results are compared with the outcome of purely numerical calculations. We conclude in Section 8.6 with some discussion of the presented as well as related work.

8.3 't Hooft's Ansatz

Our starting point is the bound state equation of the 't Hooft and Schwinger model in the two-particle sector, which, in a unified way, can be written as [99, 178]

$$\begin{aligned} M^2\phi(x) &= (m^2 - 1)\frac{\phi(x)}{x(1-x)} - \mathcal{P} \int_0^1 dy \frac{\phi(y)}{(x-y)^2} \\ &+ \alpha \int_0^1 dx \phi(x) . \end{aligned} \quad (8.23)$$

This expression corresponds to the first line of (6.18) with only the matrix element $W_2 \equiv \langle q\bar{q}|W|q\bar{q}\rangle$ retained. We will refer to (8.23) as the 't Hooft-Bergknoff equation in what follows. $\phi(x)$ denotes the valence part of the 'pion' wave function, x and y the momentum fraction of one of the two fermions in the meson. The symbol $\mathcal{P}$ indicates that the integral is defined as a principal value [127, 62]. It regularizes the Coulomb singularity $1/(x-y)^2$ in the matrix element W_2 .

The parameter α measures the strength of the anomaly. In the 't Hooft model, $\alpha = 0$ (no anomaly), and in the Schwinger model $\alpha = 1$ (representing the usual chiral anomaly). The scale parameters, as already mentioned, are given by $\mu_0^2 = g^2 N_C / 2\pi$ and $\mu_0^2 = e^2 / \pi$, respectively. M denotes the mass of the lowest lying bound state (the 'pion'). Our objective is to obtain (approximate, but accurate) solutions for M and ϕ .

Upon multiplying (8.23) with $\phi(x)$ and integrating over x we obtain for the eigenvalue

$$M^2(m) = (m^2 - 1)\frac{I_1}{I_0} - \frac{I_2}{I_0} + \alpha \frac{I_3^2}{I_0} , \quad (8.24)$$

where we have defined the integrals

$$I_0 = \int_0^1 dx \phi^2(x) , \quad (8.25)$$

$$I_1 = \int_0^1 dx \frac{\phi^2(x)}{x(1-x)} , \quad (8.26)$$

$$I_2 = \mathcal{P} \int_0^1 dx dy \frac{\phi(x)\phi(y)}{(x-y)^2} , \quad (8.27)$$

$$I_3 = \int_0^1 dx \phi(x) . \quad (8.28)$$

I_0 is the norm of the wave function, I_1 and I_2 are matrix elements of the mass and interaction term in the light-cone Hamiltonian [129, 187] in the state $|\phi\rangle$. The integral I_3 is the wave function at the origin. An independent formula for the ‘pion’ mass-squared is obtained by integrating (8.23) over x . This results in the simple expression

$$M^2(m) = m^2 \frac{I_4}{I_3} + \alpha , \quad (8.29)$$

with the additional integral,

$$I_4 = \int_0^1 dx \frac{\phi(x)}{x(1-x)} . \quad (8.30)$$

Of course, for the exact wave functions, the right-hand sides of (8.24) and (8.29) have to coincide. As the wave functions cannot be obtained exactly, we will later use the agreement between both values for the mass-squared as a measure for the accuracy of our wave functions. To determine the latter we will use a particular class of variational ansätze.

In his original work on the subject, [127, 158], 't Hooft used the following ansatz for the wave function

$$\phi(x) = x^\beta (1-x)^\beta . \quad (8.31)$$

This ansatz is symmetric in $x \leftrightarrow 1-x$ (charge conjugation odd), and β is supposed to lie between zero and one so that the endpoint behavior is non-analytic. As a non-trivial boundary condition one has the exact solution of the massless case,

$$M^2 = \alpha , \quad \text{and} \quad \phi(x) = 1 , \quad \text{i.e.} \quad \beta = 0 . \quad (8.32)$$

The main effect of having a non-vanishing fermion mass is the vanishing of the wave functions at the endpoints implying a non-zero β . This suggests the following series expansion for β ,

$$\beta(m) = \beta_1 m + \beta_2 m^2 + \beta_3 m^3 + O(m^4) , \quad (8.33)$$

and for the ‘pion’ mass squared,

$$M^2 = \alpha + M_1 m + M_2 m^2 + M_3 m^3 + O(m^4) , \quad (8.34)$$

in accordance with (8.12).

8.3.1 Exact Endpoint Behavior

The exponent β in (8.31) can actually be determined exactly by studying the small- x behavior of the bound state equation (8.23). To this end we evaluate the principal value integral for $x \rightarrow 0$,

$$\begin{aligned} \mathcal{P} \int_0^1 dy \frac{y^\beta (1-y)^\beta}{(x-y)^2} &= x^{\beta-1} \left[\mathcal{P} \int_0^\infty dz \frac{z^\beta}{(1-z)^2} + O(x) \right] \\ &= -x^{\beta-1} \left[\pi\beta \cot \pi\beta + O(x) \right]. \end{aligned} \quad (8.35)$$

Plugging this into (8.23) and demanding the coefficient of $x^{\beta-1}$ to vanish yields the transcendental equation [127]

$$m^2 - 1 + \pi\beta \cot \pi\beta = 0, \quad (8.36)$$

which is independent of the anomaly α . Using this expression we can determine β either numerically for arbitrary m or analytically for small m by expanding

$$\beta = \frac{\sqrt{3}}{\pi} m \left(1 - \frac{1}{10} m^2 \right) + O(m^4). \quad (8.37)$$

Note that the second order coefficient, β_2 , is vanishing. Furthermore, for the exact β one has $\beta_1/\beta_3 = -1/10$, which we will use as a check for our numerical results later on.

Our task is now to determine the coefficients M_i , $i = 1, 2, 3$ in (8.34). The ansatz (8.31) leads to the following results for the 'pion' mass squared (8.24),

$$\begin{aligned} M^2(m, \beta) &= (m^2 - 1) \left(\frac{1}{\beta} + 4 \right) \\ &+ \left(\frac{1}{4} + \beta \right) \frac{B^2(\beta, \beta)}{B(2\beta, 2\beta)} \left[1 + \alpha \frac{\beta}{(2\beta + 1)^2} \right]. \end{aligned} \quad (8.38)$$

In the above, $B(z_1, z_2)$ denotes the Beta function. The relevant formulae for double integrals like I_2 (8.27) can be found in [189] and [190], Appendix C. For $\alpha = 0$, (8.38) has also been obtained in [186]. Let us emphasize that this result, which expresses the 'pion' mass squared as a function of the (exactly known) endpoint exponent β , is only approximate as 't Hooft's ansatz (8.31) for the wave function does *not* represent an exact solution of the bound-state equation. It is only the endpoint behavior that is known exactly; in the intermediate- x region 't Hooft's ansatz is only an approximation that presumably becomes rather bad for large masses (non-relativistic limit) where the wave function is strongly peaked at $x = 1/2$. The accuracy of the result will be discussed extensively later on.

In order to find M^2 to order m^3 we need to expand (8.38) to order β^3 , as β itself is of order m . The result is

$$\begin{aligned}
M^2(m, \beta) &= \frac{m^2}{\beta} + 4m^2 + \alpha + \frac{\pi^2}{3}\beta \\
&+ 4\left[\pi^2/3 - 3\zeta(3) + \alpha(\pi^2/12 - 1)\right]\beta^2 \\
&+ \left[\frac{3}{5}\pi^4 - 48\zeta(3) + 4\alpha(4 - 3\zeta(3))\right]\beta^3 \\
&+ O(\beta^4).
\end{aligned} \tag{8.39}$$

Inserting β from (8.37) one finds

$$\begin{aligned}
M^2(m) &= \alpha + 2\frac{\pi}{\sqrt{3}}m \\
&+ \left[4\left(2 - \frac{9}{\pi^2}\zeta(3)\right) + \alpha\left(1 - \frac{12}{\pi^2}\right)\right]m^2 \\
&+ \frac{3\sqrt{3}}{\pi^3}\left[\left(\frac{3}{5}\pi^4 - 48\zeta(3)\right) + 4\alpha(4 - 3\zeta(3))\right]m^3 \\
&+ O(m^4).
\end{aligned} \tag{8.40}$$

Let us give the explicit results for the 't Hooft and Schwinger model, *i.e.* for $\alpha = 0$ and $\alpha = 1$, respectively. For $\alpha = 0$, we have

$$M_1 = 2\frac{\pi}{\sqrt{3}} = 3.627599, \tag{8.41}$$

$$M_2 = 4\left(2 - \zeta(3)\frac{9}{\pi^2}\right) = 3.615422, \tag{8.42}$$

$$M_3 = \frac{3\sqrt{3}}{\pi^3}\left(\frac{3}{5}\pi^4 - 48\zeta(3)\right) = 0.125139, \tag{8.43}$$

and for $\alpha = 1$,

$$M_1 = 2\frac{\pi}{\sqrt{3}} = 3.627599, \tag{8.44}$$

$$M_2 = 4\left(2 - \zeta(3)\frac{9}{\pi^2}\right) + 1 - \frac{12}{\pi^2} = 3.399568, \tag{8.45}$$

$$\begin{aligned}
M_3 &= \frac{3\sqrt{3}}{\pi^3}\left[\left(\frac{3}{5}\pi^4 - 48\zeta(3)\right) + 4\left(4 - 3\zeta(3)\right)\right] \\
&= 0.389137.
\end{aligned} \tag{8.46}$$

We note that the anomaly does *not* contribute to the first order term which therefore is the same in both the Schwinger and 't Hooft model. This confirms the observation made in Section 8.2, the coincidence of (8.14) and (8.20). As

M_1 is proportional to the condensate, *cf.* (8.17) and (8.21), the latter is also independent of the anomaly [99].

For the Schwinger model ($\alpha = 1$), the second order result can be compared with the analytical calculation of Adam, (8.19). Astonishingly, the relative difference is smaller than for the first order, namely 0.36 %. This is accidental, as we shall see below.

As a cross check, we determine the 'pion' mass squared using the alternative formula (8.29). Plugging in the ansatz (8.31), we find the fairly simple expression

$$\tilde{M}^2(m, \beta) = 4m^2 + \alpha + \frac{2m^2}{\beta} . \quad (8.47)$$

Inserting the expansion (8.37) this becomes

$$\tilde{M}^2 = \alpha + \tilde{M}_1 m + \tilde{M}_2 m^2 + \tilde{M}_3 m^3 + O(m^4) , \quad (8.48)$$

with the coefficients $\tilde{M}_i$ explicitly given by

$$\tilde{M}_1 = 2\pi/\sqrt{3} = 3.627599 , \quad (8.49)$$

$$\tilde{M}_2 = 4 , \quad (8.50)$$

$$\tilde{M}_3 = \tilde{M}_1/10 = 0.03627599 . \quad (8.51)$$

In (8.48) the anomaly contributes only in zeroth order (as β is independent of α), so that the $\tilde{M}_i$ are the same for $\alpha = 0$ and $\alpha = 1$. Comparing with (8.41) and (8.44), we see that the first order coefficients coincide for the two alternative mass formulae. In addition, one finds that $\tilde{M}_2$ roughly coincides with the M_2 of both the 't Hooft and Schwinger model to within 10-20 %. The value for $\tilde{M}_3$ is smaller by approximately an order of magnitude.

We conclude that 't Hooft's ansatz works well to lowest non-trivial order in m but needs improvement if one wants to go further. As a preparative step for such a development we now introduce a variational method that can easily be extended to accurately calculate higher orders in m .

8.3.2 Variational Approach

At variance with the above we are now going to regard β as a variational parameter to be determined by minimizing the function $M^2(m, \beta)$ of (8.39). To this end we insert the expansion (8.33) into (8.39) and obtain

$$\begin{aligned} M^2(m) &= \alpha + \left(\frac{\pi^2}{3} \beta_1 + \frac{1}{\beta_1} \right) m \\ &+ \left[4 + \beta_2 \left(\frac{\pi^2}{3} - \frac{1}{\beta_1^2} \right) \right] m^2 \end{aligned}$$

$$\begin{aligned}
& + 4\left(\frac{\pi^2}{3} - 3\zeta(3) + \alpha\left(\frac{\pi^2}{12} - 1\right)\right)\beta_1^2 m^2 \\
& + \left[8\beta_1\beta_2\left(\frac{\pi^2}{3} - 3\zeta(3)\right) + \beta_1^3\left(\frac{3}{5}\pi^4 - 48\zeta(3)\right) \right. \\
& + \frac{\beta_2^2}{\beta_1^3} + \beta_3\left(\frac{\pi^2}{3} - \frac{1}{\beta_1^2}\right) \\
& + \left.\alpha\left(2\beta_1\beta_2\left(\frac{\pi^2}{3} - 4\right) + 4\beta_1^3\left(4 - 3\zeta(3)\right)\right)\right] m^3 \\
& + O(m^4) .
\end{aligned} \tag{8.52}$$

Note that to order m only the leading coefficient β_1 contributes. We will furthermore shortly see that the dependence of M_2 on β_2 and of M_3 on β_3 is only apparent.

Solving the minimization equation, $\partial M^2/\partial\beta = 0$, for the coefficients β_i , leads to

$$\beta_1 = \sqrt{3}/\pi = 0.55133 , \tag{8.53}$$

$$\beta_2 = 0.11690 + 0.065612\alpha , \tag{8.54}$$

$$\beta_3 = 0.0049077 - 0.050811\alpha + 0.019521\alpha^2 . \tag{8.55}$$


Comparing with (8.37) we note that the coefficient β_1 is exact! Plugging it into (8.52) we verify the statement above that M_2 is independent of β_2 and M_3 independent of β_3 . Therefore, the expressions (8.52) and (8.40) coincide up to and including order m^2 . M_1 and M_2 are thus the same, irrespective of whether one uses the exact endpoint exponent of (8.37) or its variational estimate. The estimates (8.53-8.55) for the coefficients β_i differ in at least three respects from the exact values of (8.37): (i) β_2 and β_3 depend on α , (ii) $\beta_2 \neq 0$, (iii) $\beta_3 \neq -\beta_1/10$. However, all these shortcomings affect at most the third order coefficient M_3 , and thus become negligible for small m . Only if one wants to have reliable numbers for M_3 , (which we will be going to produce), these effects have to be taken into account. The present values of β_i lead to

$$\begin{aligned}
M_3(\alpha = 0) &= 0.043597 , \\
M_3(\alpha = 1) &= 0.190372 ,
\end{aligned} \tag{8.56}$$

which should be compared with (8.43) and (8.46), respectively. The differences are large so that the agreement is not particularly good. These discrepancies, however, are not disturbing at this point, as we are calculating a third order effect with just one variational parameter. We will do better later on by enlarging the number of these parameters until we see satisfactory convergence of the results.

The variational estimate for β can also be plugged into the alternative mass formula, $\tilde{M}^2(m, \beta)$, (8.47), which by itself does not constitute a variational problem. We do not give the analytical results for the $\tilde{M}_i$ here but simply refer

Figure 8.1: Comparison of the two alternative mass formulae, (8.38) for $M^2(\beta)$ and (8.47) for $\tilde{M}^2(\beta)$, and the second order bosonization results [175, 176] for $m = 0.1$. The vertical line marks the minimum of the curve $M^2(\beta)$ yielding the variational estimate for β using 't Hooft's ansatz. For this value of β one finds $M^2(m = 0.1) = 1.3969$ and $\tilde{M}^2(m = 0.1) = 1.3913$. The bosonization result (short-dashed horizontal line) is $M_{\text{Bos}}^2(m = 0.1) = 1.390 \pm 1 \cdot 10^{-3}$, where the error is basically an estimate of the unknown third order coefficient. Thus, within 't Hooft's ansatz, the alternative mass formula yields a somewhat better result at $m = 0.1$. In the next sections we will refine our methods so that the variational estimates for M^2 and $\tilde{M}^2$ will converge towards each other.



to Fig. 8.1 for a qualitative comparison of the two alternative formulae, and to Section 8.4, Tables 8.3 and 8.7, for the actual numbers.

As stated repeatedly, 't Hooft's ansatz (8.31) used so far has to be extended if one wants to accurately determine the second and third order coefficients of M^2 . This is our next issue.

8.4 Extension of 't Hooft's Ansatz

For the numerical calculations of [186, 190, 188] the following set of trial functions has been used:

$$\phi(x) = \sum_{k=0}^N u_k [x(1-x)]^{\beta+k}. \quad (8.57)$$

Obviously, the term with $k = 0$ and normalization $u_0 = 1$ corresponds to 't Hooft's original ansatz (8.31). The coefficients u_k are treated as additional variational parameters, so that, according to the variational principle, (8.57) *must* yield a better result than (8.31). The question is, how big the improvement will be. To this end we will calculate the coefficients M_i by adding more and more basis functions to 't Hooft's original ansatz (8.31), thus enlarging our space of variational parameters. We will follow the two different approaches mentioned above, namely, (i) use the exact β from (8.37), or, (ii) treat β as one of the variational parameters. For both approaches we will compare the calculated coefficients M_i with the $\tilde{M}_i$ obtained from the alternative mass formula (8.29). We will continue adding basis functions until we see our results converge. The maximum number N of basis functions in this section will be five, *i.e.* 't Hooft's original wave function plus four corrections.

A particular benefit of the ansatz (8.57) is the fact that the integrals I_1 to I_4 can still be evaluated analytically though the formulae become rather lengthy. For this reason we will make heavy use of the program package MAPLE in what follows. As a result we have analytical expressions for all the quantities we calculate. As these expressions cover pages and pages without being very instructive, we do not display them but rather the evaluated numbers. In this sense, our treatment might be called 'semi-analytical'. A major advantage, however, of using computer algebraic manipulations is the high calculational accuracy which is only limited by the maximum number of digits the machine can handle.

8.4.1 Purely Variational Approach

In this subsection, β will always be treated as a variational parameter and thus be obtained by minimizing the 'pion' mass squared, M^2 . In the ansatz (8.57) we use up to four additional basis functions, so that our maximum N is four. To avoid an inflation of indices we rename the coefficients u_k , $k = 1, \dots, 4$ by a , b , c and d , respectively. Their expansion coefficients, defined via

$$\begin{aligned} a &= a_1 m + a_2 m^2, \\ b &= b_1 m + b_2 m^2, \\ c &= c_1 m + c_2 m^2, \\ d &= d_1 m + d_2 m^2, \end{aligned} \tag{8.58}$$

are determined (recursively) by minimizing M^2 with respect to them. Let us start with the 't Hooft model ($\alpha = 0$).

't Hooft Model

In order to save space we do not display all the values for the coefficients in (8.58). The best values obtained are given in the last section when we discuss the quality

Table 8.1: *Expansion coefficients of the end point exponent β for the 't Hooft model obtained by successively varying with respect to β (first line) , β and a (second line), etc. The first column, β_1 , coincides with the result from the exact endpoint analysis, which in addition yields $\beta_2 = 0$, $\beta_3 = -\beta_1/10$.*

<i>Ansatz</i>	$\beta_1 = \sqrt{3}/\pi$	β_2	β_3
't Hooft	0.55132889	0.11689763	0.00490773
a	0.55132889	0.00976951	-0.04010634
b	0.55132889	0.00256081	-0.04889172
c	0.55132889	0.00102086	-0.05209341
d	0.55132889	0.00050738	-0.05343097

of the wave function. Here, we rather display the results for the coefficients β_i of β (see Table 8.1), as these can be compared with the exact values of (8.37).

As an important finding we note that β_1 remains unchanged at the exact value $\sqrt{3}/\pi$ (a 'variational invariant'). β_2 tends to zero, as it should. The non-vanishing of β_2 only affects the coefficient M_3 , as M_1 and M_2 do not depend on β_2 . The convergence of β_3 towards $-\beta_1/10$ seems somewhat slow; but as all M_i , $i = 1, 2, 3$, are independent of β_3 this has no observable effect.

In Table 8.2 we list the expansion coefficients of the mass squared, M^2 . One notes that the convergence of the results for M_2 and M_3 is rather good. For M_2 we finally have a relative accuracy of $8 \cdot 10^{-7}$, and for M_3 of $4 \cdot 10^{-5}$. Furthermore, the coefficients are getting smaller if one adds more basis functions, in accordance with the variational principle. As the coefficient M_1 is entirely determined by the 'variational invariant' β_1 it remains unchanged at $2\pi/\sqrt{3}$. This is another important result: M_1 stays fixed at the standard 't Hooft-Bergknoff value (8.14).

If we use the alternative mass formula (8.29) and evaluate it using the wave function calculated above we find the values for $\tilde{M}^2$ listed in Table 8.3.

We note that the values of the $\tilde{M}_i$ are somewhat more sensitive to the values of the expansion coefficients β_i , as $\tilde{M}_2$ does depend on β_2 , and $\tilde{M}_3$ on β_2 and β_3 . To check the quality of our wave functions one should compare the results for M_i and $\tilde{M}_i$ which are listed in Tables 8.2 and 8.3. Again, it is gratifying to note that we are improving our results step by step in the variational procedure. The values of $\tilde{M}_i$ converge towards those of M_i . The relative accuracy is approximately 10^{-3} . This error is entirely due to the difference between the calculated and the real wave function. It is clear that the accuracy in the single eigenvalue M^2 (Table 8.2) is higher than the one for the wave function, where in principle an infinite number of points has to be calculated. This represents an indirect proof that we are indeed improving our wave functions, and not just the eigenvalues.

Table 8.2: Expansion coefficients of M^2 for the 't Hooft model obtained by successively enlarging the space of variational parameters. M_1 is the standard 't Hooft-Bergknoff result. Note the good convergence towards the bottom of the table.

<i>Ansatz</i>	$M_1 = 2\pi/\sqrt{3}$	M_2	M_3
't Hooft	3.62759873	3.61542218	0.043597197
a	3.62759873	3.58136872	0.061736701
b	3.62759873	3.58107780	0.061805257
c	3.62759873	3.58105821	0.061795547
d	3.62759873	3.58105532	0.061793082

Table 8.3: Expansion coefficients of the alternative mass squared $\tilde{M}^2$ for the 't Hooft model obtained by successively enlarging the space of variational parameters. Again, the fixed value for $\tilde{M}_1$ is the standard 't Hooft-Bergknoff result.

<i>Ansatz</i>	$\tilde{M}_1 = 2\pi/\sqrt{3}$	$\tilde{M}_2$	$\tilde{M}_3$
't Hooft	3.62759873	3.23084437	0.130791594
a	3.62759873	3.54922830	0.066592425
b	3.62759873	3.57265307	0.059652468
c	3.62759873	3.57769969	0.060282959
d	3.62759873	3.57938609	0.060854772

Table 8.4: *Expansion coefficients of the endpoint exponent β for the Schwinger model obtained by successively enlarging the space of variational parameters. The first column, β_1 , coincides with the result from the exact endpoint analysis, which in addition yields $\beta_2 = 0$, $\beta_3 = -\beta_1/10$.*

<i>Ansatz</i>	β_1	β_2	β_3
't Hooft	0.55132889	0.18250945	-0.026382222
a	0.55132889	0.01177681	-0.040372413
b	0.55132889	0.00317437	-0.048081756
c	0.55132889	0.00127507	-0.051602713
d	0.55132889	0.00063609	-0.053130398

Table 8.5: *Expansion coefficients of M^2 for the Schwinger model obtained by successively enlarging the space of variational parameters. Again, the fixed value for M_1 is the standard 't Hooft-Bergknoff result. Note the good convergence towards the bottom of the table.*

<i>Ansatz</i>	$M_1 = 2\pi/\sqrt{3}$	M_2	M_3
't Hooft	3.62759873	3.39956798	0.19037224
a	3.62759873	3.30906326	0.34776772
b	3.62759873	3.30864244	0.34820193
c	3.62759873	3.30861240	0.34820513
d	3.62759873	3.30860791	0.34820389

Schwinger Model

For the Schwinger model ($\alpha = 1$), we perform exactly the same calculations. The expansion coefficients of β are listed in Table 8.4. Although the coefficients β_2 , β_3 are somewhat different from those in Table 8.1 (which they must be as they lose their dependence on α only in the strict limit of *exact* evaluation) we note the same tendency: β_2 converges to zero and β_3 towards $-\beta_1/10$.

The best values for the 'pion' mass squared are again provided by the variational results listed in Table 8.5. The numerical accuracy is practically the same as for the analogous Table 8.2. The comparison with the alternatively calculated coefficients $\tilde{M}^2$ is given in Table 8.6. Again, everything is completely analogous to the case of the 't Hooft model ($\alpha = 0$).

Table 8.6: *Expansion coefficients of the alternative mass squared $\tilde{M}^2$ for the Schwinger model obtained by successively enlarging the space of variational parameters. Again, the fixed value for $\tilde{M}_1$ is the standard 't Hooft-Bergknoff result.*

<i>Ansatz</i>	$\tilde{M}_1 = 2\pi/\sqrt{3}$	$\tilde{M}_2$	$\tilde{M}_3$
't Hooft	3.62759873	2.79913596	0.57111672
a	3.62759873	3.27031909	0.36868190
b	3.62759873	3.29819920	0.34869835
c	3.62759873	3.30441759	0.34753576
d	3.62759873	3.30651525	0.34762562

8.4.2 Variational Approach using the exact β

In this subsection we calculate M^2 and $\tilde{M}^2$ for both values of α using the exact value (8.37) for β . Thus only a , b , c , d are treated as variational parameters. This is the procedure employed numerically in [188]. From the discussion of the preceding subsection, in particular the values for the β_i displayed in Table 8.1, which differ minimally from the exact values, we expect that the results will be very close to those from the purely variational approach.

't Hooft Model

This is exactly what happens as can be seen from Tables 8.7 and 8.8. Comparing Table 8.7 with Table 8.2 one finds that the coefficients M_1 and M_2 of both tables coincide as these only depend on β_1 which is the same in both approaches. Only for M_3 there are slight differences, due to the dependence on β_2 . For the $\tilde{M}_i$, $i = 2, 3$, the discrepancies are somewhat bigger as these coefficients do depend on β_2 and β_3 (Table 8.3 *vs.* Table 8.8). Still the consistency is quite obvious.

Upon comparing the second columns of Tables 8.2 and 8.3, respectively Tables 8.7 and 8.8, one notes a funny coincidence. In the first case M_2 is slightly bigger than $\tilde{M}_2$, and vice versa in the second case. If one denotes the purely variational results with a superscript ' v ', and the results obtained with the exact β with a superscript ' e ', one finds that M_2 , which is the same in both approaches, is given by the arithmetic mean

$$M_2 = \frac{1}{2} \left(\tilde{M}_2^e + \tilde{M}_2^v \right) . \quad (8.59)$$

We have checked this analytically for 't Hooft's ansatz. For the higher orders this is difficult to do but the numerical evidence is beyond doubt.

Comparing the third columns of Tables 8.2 and 8.7, one finds that $M_3^v < M_3^e$. We thus see a slight tendency that the results of the purely variational procedure

Table 8.7: Expansion coefficients of M^2 ('t Hooft model) obtained by using the exact endpoint exponent β and successively enlarging the space of variational parameters.. Again, the fixed value for M_1 is the standard 't Hooft-Bergknoff result.

<i>Ansatz</i>	$M_1 = 2\pi/\sqrt{3}$	M_2	M_3
't Hooft	3.62759873	3.61542219	0.12513878
a	3.62759873	3.58136873	0.06230622
b	3.62759873	3.58107781	0.06184441
c	3.62759873	3.58105821	0.06180178
d	3.62759873	3.58105533	0.06179462

Table 8.8: Expansion coefficients of the alternatively defined mass squared $\tilde{M}^2$ ('t Hooft model) obtained by using the exact endpoint exponent β and successively enlarging the space of variational parameters. Again, the fixed value for $\tilde{M}_1$ is the standard 't Hooft-Bergknoff result.

<i>Ansatz</i>	$\tilde{M}_1 = 2\pi/\sqrt{3}$	$\tilde{M}_2$	$\tilde{M}_3$
't Hooft	3.62759873	4.0	0.36275987
a	3.62759873	3.61350915	0.08659575
b	3.62759873	3.58950254	0.07106398
c	3.62759873	3.58441672	0.06603497
d	3.62759873	3.58272458	0.06406179

Table 8.9: *Expansion coefficients of M^2 (Schwinger model) obtained by using the exact endpoint exponent β and successively enlarging the space of variational parameters. Again, the fixed value for M_1 is the standard 't Hooft-Bergknoff result.*

<i>Ansatz</i>	$M_1 = 2\pi/\sqrt{3}$	M_2	M_3
't Hooft	3.62759873	3.39956798	0.38913656
a	3.62759873	3.30906326	0.34859533
b	3.62759873	3.30864244	0.34826210
c	3.62759873	3.30861239	0.34821485
d	3.62759873	3.30860791	0.34820630

Table 8.10: *Expansion coefficients of the alternatively defined mass squared $\tilde{M}^2$ (Schwinger model) obtained by using the exact endpoint exponent β and successively enlarging the space of variational parameters. Again, the fixed value for $\tilde{M}_1$ is the standard 't Hooft-Bergknoff result.*

<i>Ansatz</i>	$\tilde{M}_1 = 2\pi/\sqrt{3}$	$\tilde{M}_2$	$\tilde{M}_3$
't Hooft	3.627598730	4.0	0.3627598730
a	3.627598730	3.347807427	0.3637551073
b	3.627598730	3.319085690	0.3566261465
c	3.627598730	3.312807215	0.3523062162
d	3.627598730	3.310700593	0.3504584408

are better than those obtained using the exact β . The same observation has been made by Mo and Perry [187], in particular for large values of the fermion mass m (where the third order coefficient M_3 becomes important). We interpret this fact as a hint that for larger m the behavior of the wave function in the intermediate region becomes more relevant compared to the endpoint behavior.

Schwinger Model

For the Schwinger model, the analogous results are listed in Tables 8.9 and 8.10. Exactly the same remarks as above apply including the size of the errors and the relation between M_2 and $\tilde{M}_2^e$, $\tilde{M}_2^v$. In the next section we will verify our results by purely numerical methods.

Table 8.11: Numerical results using lowest order (two-particle) light-front Tamm-Dancoff approximation for $M^2 - 1$ as a function of the fermion mass m , for $\alpha = 1$ (Schwinger model).

Ansatz	$m = 0.0001$	$m = 0.0005$	$m = 0.001$	$m = 0.005$	$m = 0.01$	$m = 0.05$	$m = 0.1$
't Hooft	0.000362794	0.00181465	0.00363100	0.0182230	0.0366163	0.189927	0.397130
a	0.000362858	0.00181465	0.00363086	0.0182207	0.0366072	0.189695	0.396181
b	—	0.00181461	0.00363100	0.0182210	0.0366071	0.189694	0.396176
c	—	0.00181459	—	—	0.0366071	0.189694	0.396176
d	—	—	—	—	0.0366073	0.189694	0.396176

8.5 Comparison With Numerical Results

For the numerical calculations the ansatz (8.57) has been used. The value for β has been determined numerically from (8.36). Again, up to four additional basis functions have been included (for larger masses even five). Let us begin with the Schwinger model ($\alpha = 1$). Here we can use the code developed in [188]. In Table 8.11 we list $M^2 - 1$ as a function of the fermion mass m , calculated within two-particle light-front Tamm-Dancoff approximation. The notations a,b,c,d are as in the preceding section.

One main difference in comparison with the computer algebraic treatment of Section 8.4 is the numerical inaccuracy for small m . This is a general disease of numerical treatments, and also shared e.g. by the lattice [168, 169] or DLCQ approach [183]. In these approaches, however, the numerical errors are typically much larger than ours (see the next section for an explicit comparison). Due to the small- m instability, our numerical calculation does not converge within an arbitrary number of digits. As soon as a calculated value for $M^2 - 1$ becomes bigger as the preceding one (a numerical violation of the variational principle), we terminate the procedure and pick the smaller value as our final result. The difference between these last two values can be used as an estimate of the numerical inaccuracy. The errors will be further discussed below and in the next section when we compare our results with related work (see Tables 8.15, 8.16).

It should also be pointed out that the numerical analysis is conceptually very different from the computer algebraic treatment. As one cannot perform a Taylor expansion numerically, the expansion coefficients of M^2 have to be obtained by fitting polynomials to $M^2(m)$. Clearly, this is an additional source of errors, and one expects the results to be less accurate than those of the preceding sections. A cubic fit to the optimum values for $M^2 - 1$ in Table 8.11 yields the following expansion coefficients of M^2 ,

$$M_1 = 3.62609, \quad (8.60)$$

$$M_2 = 3.33607, \quad (8.61)$$

$$M_3 = 0.22396, \quad (8.62)$$

which should be compared with the last line of Table 8.5. To estimate the accuracy of these values we also show the results from a quartic fit,

$$M_1 = 3.62755, \quad (8.63)$$

$$M_2 = 3.31029, \quad (8.64)$$

$$M_3 = 0.32984. \quad (8.65)$$

Comparing (8.60-8.62) and (8.63-8.65) one finds that the stability of the fits is not too impressive in view of the accuracy we would like to achieve. In particular the third order coefficient is numerically difficult to determine. We estimate the relative accuracy as being 10^{-3} for M_1 , 10^{-2} for M_2 and 10^{-1} for M_3 . This is also confirmed by comparing with Table 8.5.

For the Schwinger model it is possible to check the influence of higher particle sectors on the 'pion' mass squared by using the machinery developed in [188] for the wave functions of the higher Fock components. These are the amplitudes of finding not only two, but four, six, ... (anti-)fermions in the 'pion'. In Table 8.12, we list the best values for $M^2 - 1$ including up to six-body wave functions. We also show the 2-, 4-, and 6-body content of the total wave function. It is known that the 'pion' is entirely 2-particle in the chiral limit [187]. For small mass, one therefore expects only small contributions from the higher Fock sectors. This is confirmed by the numerical results. Astonishingly, this feature persists up to values of at least $m = 0.5$ for the fermion mass. This fact is in agreement with the observation of Mo and Perry that the four-particle component of the wave function is less than 0.4 % for *all* values of the fermion mass [187]. A similar result has been found in the DLCQ calculations of [183]. In addition we note that there seems to be some interesting kind of hierarchy between the relative strengths of the contributions from different particle sectors. If we denote the $2k$ -particle amplitude in the wave function by f_{2k} , we find that $|f_2|^2 \gg |f_4|^2 \gg |f_6|^2$, the individual proportions being several orders of magnitude (see Table 8.12). Here, we explicitly see the magic of light-front field theory at work: high Fock components in bound states tend to be largely suppressed, at variance with the situation encountered in field theory quantized the standard way.

From the point of view of the variational principle, the results shown in Table 8.12 are a bit better (i.e. smaller) than those of Table 8.11 (apart from the value for $m = 0.0005$), but the improvement is rather small. A cubic fit yields

$$M_1 = 3.62667, \quad (8.66)$$

$$M_2 = 3.23696, \quad (8.67)$$

$$M_3 = 0.36235, \quad (8.68)$$

and a quartic fit

$$M_1 = 3.62747, \quad (8.69)$$

Table 8.12: *Numerical Results for $M^2 - 1$ as a function of the fermion mass m , for $\alpha = 1$ (Schwinger model) including 2-, 4- and 6-body wave functions. The contributions of the different Fock sectors to the total wave function squared are given in percent.*

	$m = 0.0001$	$m = 0.0005$	$m = 0.001$	$m = 0.005$	$m = 0.01$	$m = 0.05$	$m = 0.1$
$M^2 - 1$	0.000362793	0.00181481	0.00363055	0.0182179	0.0365968	0.189468	0.395400
% 2-body	100.00	100.00	100.00	99.999990	99.999962	99.999213	99.997514
% 4-body	—	—	—	0.000010	0.000038	0.000782	0.002474
% 6-body	—	—	—	—	—	0.000005	0.000013

Table 8.13: *M^2 as a function of the fermion mass m , for $\alpha = 0$ ('t Hooft model), obtained with Code I.*

m	0.0001	0.001	0.01	0.1
M^2	0.000362795	0.00363109	0.0366342	0.398634

$$M_2 = 3.20864 , \quad (8.70)$$

$$M_3 = 0.60365 . \quad (8.71)$$

Compared with (8.60-8.62) and (8.63-8.65) we do not find any absolute improvement in M_1 that could be distinguished from the numerical inaccuracy. For the coefficient M_2 , which in bosonization schemes is 3.3874, we even get the wrong tendency: it becomes smaller upon including higher Fock states. The inaccuracy for M_3 is so large that this coefficient is only determined in its order of magnitude.

Altogether, we arrive at the very important conclusion that the inclusion of higher particle sectors in the light-front bound state equation of the Schwinger model does not diminish the discrepancy between the results obtained via the 't Hooft-Bergknoff method and those from bosonization techniques.

Let us move to the 't Hooft model. If we put $\alpha = 0$ in the Schwinger model code above (Code I) we find the 'pion' mass squared for the 't Hooft model. The best values within the 2-particle sector are listed in Table 8.13. Note that higher particle sectors are strictly suppressed in the large- N_C limit. Thus, there is no point in calculating these, unless one is interested in $1/N_C$ corrections to the 't Hooft model results [178, 184, 185].

In view of the restricted number of data points only a quadratic fit makes sense which yields

$$M_1 = 3.62754 , \quad (8.72)$$

$$M_2 = 3.58796 , \quad (8.73)$$

Table 8.14: M^2 as a function of the fermion mass m , for $\alpha = 0$ ('t Hooft model), obtained with Code II.

m	0.0001	0.001	0.01	0.1
M^2	0.000362795	0.00363118	0.0366341	0.398634

which is consistent with the results of Section 8.4 (see Table 8.2). A second code (Code II), which was independently developed for the 't Hooft model [185] yields the results displayed in Table 8.14. From a quadratic fit we obtain

$$M_1 = 3.62754 , \quad (8.74)$$

$$M_2 = 3.58806 . \quad (8.75)$$

Comparing with (8.72, 8.73) we see that both codes yield the same results within the numerical accuracy. This is reassuring, since, as already stated, the codes were developed independently from each other.

A cubic fit to a total of 30 data points produced with Code II yields

$$M_1 = 3.62758 , \quad (8.76)$$

$$M_2 = 3.58260 , \quad (8.77)$$

$$M_3 = 0.06450 , \quad (8.78)$$

in fair agreement with the variationally obtained values of Table 8.2. To check the stability of this fit we can compare with the extension to fourth order,

$$M_1 = 3.62756 , \quad (8.79)$$

$$M_2 = 3.58314 , \quad (8.80)$$

$$M_3 = 0.06280 , \quad (8.81)$$

$$M_4 = 0.001232 . \quad (8.82)$$

This leads to an estimate of roughly 10^{-6} , 10^{-4} and 10^{-2} for the (absolute) numerical error in the first, second and third order coefficient, respectively. Furthermore it is gratifying to note that the fourth order coefficient, M_4 , is numerically small.

As is obvious from the discussion above, the numerical errors for the 't Hooft model are much smaller than those for the Schwinger model. The basic reason for this is the Schwinger model anomaly. Note that in the bound state equation the anomaly factor α multiplies an integral over the wave function. To evaluate this, one needs the wave function as a whole. In the 't Hooft model, on the other hand, this term is absent and the eigenvalues are dominated entirely by the endpoint behavior of the wave function which is known exactly. This makes the numerical errors much smaller (see also Tables 8.15, 8.16).

8.6 Discussion

In the preceding sections we have calculated the expansion coefficients M_i in the series

$$M^2 = \alpha + M_1 m + M_2 m^2 + M_3 m^3 \quad (8.83)$$

for the ‘pion’ mass squared of both the ’t Hooft ($\alpha = 0$) and the Schwinger model ($\alpha = 1$). In order for the expansion (8.83) to make sense we have considered only small masses $m \ll 1$ *i.e.* $m \ll \mu_0$ in the original units. For the ’t Hooft model we have $\mu_0^2 = g^2 N_C / 2\pi$ with $N_C \rightarrow \infty$, $g^2 N_C$ fixed. Thus, we are working in the *weak* coupling phase where the limit $N_C \rightarrow \infty$ ($g \rightarrow 0$), is taken *before* the limit $m \rightarrow 0$, or, equivalently, such that one always has $m \gg g \sim 1/\sqrt{N_C} \rightarrow 0$ [162, 191]. For the Schwinger model, $\mu_0^2 = e^2/\pi$, so that small fermion mass corresponds to *strong* coupling.

We have used analytical, computer algebraic and numerical methods of different accuracy. Nonetheless, the overall picture is intrinsically consistent. Our main findings are the following:

(i) The first order coefficient, M_1 , in the expansion of the ‘pion’ mass squared is independent of the anomaly α , *i.e.* the same in both the ’t Hooft and Schwinger model. This confirms the results of ’t Hooft [127, 158] and Bergknoff [129].

(ii) The ’t Hooft-Bergknoff value, $M_1 = 2\pi/\sqrt{3}$, is a ‘variational invariant’: it does not get altered by extending the space of variational parameters. This has been checked analytically and numerically.

(iii) In the Schwinger model, the ’t Hooft-Bergknoff value does not change upon inclusion of higher particle sectors. Only the second and third order coefficients, M_2 and M_3 , are affected.

(iv) Thus, for the Schwinger model, there remains a few percent discrepancy in the coefficients M_1 and M_2 compared to bosonization results.

(v) The variational calculation yields the ‘pion’ wave function to a high accuracy. The endpoint behavior is reproduced exactly (in leading order in m). The behavior in the intermediate region, $0 < x < 1$, gets improved as can be seen from Fig. 8.3 below.

In Table 8.15 we summarize our optimum final results for the ’t Hooft model ($\alpha = 0$) and compare with results that have been obtained previously.

Upon inspection of Table 8.15 one finds a very good overall consistency of the results. All given values are in good agreement within error bars. The scale is set by the variational results which are the most accurate ones. Within error bars, they are matched by our numerical results as well as by previous analytical and numerical calculations.

As far as the latter are concerned, a few remarks are in order. The coefficients M_i displayed in the last line of Table 8.15 are obtained by performing polynomial fits to the values calculated numerically in [192] (which basically agree with those from ’t Hooft’s original calculation displayed in Fig. 5 of [127]). The averaged

Table 8.15: *The expansion coefficients M_i for the 't Hooft model ($\alpha = 0$). The errors of our results obtained within 2-particle Tamm-Dancoff (2PTD) approximation are estimated by comparing the last two lines in Table 8.2 (for the variational method) and the different polynomial fits (8.72 - 8.82) (for the numerical results). The numerical inaccuracies given in the last line were estimated by us using the polynomial fit method.*

	M_1	M_2	M_3
variational 2PTD	$2\pi/\sqrt{3} = 3.627599$	$3.581055 \pm 3 \cdot 10^{-6}$	$0.061793 \pm 3 \cdot 10^{-6}$
numerical 2PTD	$3.62758 \pm 2 \cdot 10^{-5}$	$3.5829 \pm 3 \cdot 10^{-4}$	$0.064 \pm 1 \cdot 10^{-3}$
't Hooft [158]	$2\pi/\sqrt{3} = 3.627599$	—	—
Burkardt [180]	$2\pi/\sqrt{3} = 3.627599$	3.5812	—
Li [181]	3.64 ± 0.06	—	—
Li et al. [192]	3.64 ± 0.03	3.60 ± 0.06	0.04 ± 0.04

result of the fits is shown including an estimate of the errors. It should also be stressed that the data in [192] are mainly obtained for $m > 0.5$ whereas the bulk of our values for the M_i are obtained for $m < 0.1$. Nevertheless, the agreement with our results is satisfactory within error bars.

From Table 8.15 it is obvious that all numerical results favor the first order 't Hooft-Bergknoff value of $2\pi/\sqrt{3}$. This agreement is particularly gratifying for the data of [192] which were not obtained via light-cone techniques but within ordinary quantization and even within a different gauge, namely axial gauge. The agreement is therefore highly non-trivial [193].

For the Schwinger model, ($\alpha = 1$), we summarize our findings in Table 8.16. For comparison we have also listed analytical results obtained via bosonization techniques and data which we extracted from polynomial fits to numerical light-front Tamm-Dancoff [187], DLCQ [183] and lattice results [194]. The contents of Table 8.16 are graphically displayed in Fig. 8.2, where we have chosen rescaled units as in [176].

As already stated, due to the anomaly the numerical errors for the Schwinger model are at least an order of magnitude larger than for the 't Hooft model. In addition, the 2% discrepancy between our light-cone results and the analytic bosonization results does not get resolved. However, we have shown that the discrepancy is not due to (i) inaccuracies in the wave function or (ii) neglect of contributions from the next higher Fock sectors (four and six particles). It should be mentioned that other numerical methods (DLCQ, lattice) are by far too inaccurate to distinguish between the 't Hooft-Bergknoff and the bosonization value. We will come back to these issues in a moment.

Figure 8.2: The rescaled ‘pion’ mass $\bar{M} = M/\sqrt{1+m^2}$ as a function of the fermion mass m . The short-dashed curve represents a ‘phenomenological’ parametrization for $\bar{M}$ with $M^2 = 1 + M_1 m + 4m^2$, which becomes exact for very small and very large m and thus smoothly interpolates between the strong and weak coupling regions. The long-dashed curve is our second-order and the solid curve our third-order result. As expected, mass perturbation theory breaks down as m becomes of order 1. The crosses are the lattice results [194], the diamonds the lattice results [195], which go down to comparatively small masses, however with large errors. The circles are light-front Tamm-Dancoff results [187], whereas the squares [183] and triangles [196], as quoted in [176], are DLCQ results. The values corresponding to the triangles are not included in Table 8.16. The 2% discrepancy in M_1 is invisible within the resolution of the figure.




Table 8.16: The expansion coefficients M_i for the Schwinger model ($\alpha = 1$). The errors for the variational 2-particle Tamm-Dancoff (2PTD) approximation are estimated by comparing the last two lines in Table 8.5. The errors of our numerical calculations are obtained from comparing polynomial fits of different order to the numerical results of Tables 8.11 and 8.12. Errors of other results are given where they could be estimated analogously.

	M_1	M_2	M_3
variational 2PTD	$2\pi/\sqrt{3} = 3.627599$	$3.308608 \pm 4 \cdot 10^{-6}$	$0.348204 \pm 1 \cdot 10^{-6}$
numerical 2PTD	$3.6268 \pm 8 \cdot 10^{-4}$	3.32 ± 0.02	0.28 ± 0.06
numerical 6PTD	$3.6267 \pm 4 \cdot 10^{-4}$	3.22 ± 0.02	0.5 ± 0.1
numerical 4PTD [187]	3.62 ± 0.07	3.2 ± 0.3	0.3 ± 0.2
DLCQ [183]	3.7 ± 0.2	3.5 ± 0.3	—
lattice [194]	3.5 ± 0.2	3.7	0.02
Adam [175]	$2e^\gamma = 3.562146$	3.3874	—
Fields et al. [176]	$2e^\gamma = 3.562146$	3.387399	—

Table 8.17: Lattice results for the first order coefficient M_1 and the (negative of the) condensate, $-\langle 0|\bar{\psi}\psi|0\rangle = M_1/4\pi$. The results are quoted in chronological order. They should be compared with the bosonization results, $M_1 = 3.562$, $-\langle 0|\bar{\psi}\psi|0\rangle = 0.283$ and the 't Hooft-Bergknoff values, $M_1 = 3.628$, $-\langle 0|\bar{\psi}\psi|0\rangle = 0.289$.

	[167]	[198]	[194]	[169]	[168]	[195]
M_1	3.42	3.644	3.48	2.97	3.33	3.77
$-\langle 0 \bar{\psi}\psi 0\rangle$	0.27	0.290	0.28	0.24	0.26	0.30

As is well known, [176, 187, 197], the DLCQ data are comparatively inaccurate for small fermion mass m (i.e. large coupling). This is due to the fact that the dominating feature of the light-cone wave function, its endpoint behavior, is not very accurately reproduced using an equally-spaced momentum grid. The poor convergence of DLCQ for small m has recently been overcome by incorporating the exact endpoint behavior [197].

The lattice data generally suffer from the same disease of having rather large error bars for small m . The chiral limit, $m = 0$, can only be reached via extrapolation in a very inaccurate manner unless one uses the Schwinger result $M^2(m = 0) = 1$ as a bias [194]. A representative collection of lattice results for M_1 is given in Table 8.17. For the convenience of the reader we also list the condensate $\langle 0|\bar{\psi}\psi|0\rangle = -M_1/4\pi$.

The values for the Schwinger model condensate in Table 8.17 should be com-

pared with our results. If we assume that formulae (8.17) and (8.18) are directly applicable to our calculation we get

$$\langle 0|\bar{\psi}\psi|0\rangle_1 \equiv -\frac{M_1}{4\pi} = -\frac{1}{2\sqrt{3}} = -0.28868, \quad (8.84)$$

$$\langle 0|\bar{\psi}\psi|0\rangle_2 \equiv -\frac{1}{2\pi}\sqrt{\frac{M_2}{A+B}} = -0.28015, \quad (8.85)$$

while the standard result is

$$\langle 0|\bar{\psi}\psi|0\rangle = -\frac{e^\gamma}{2\pi} = -0.28347. \quad (8.86)$$

As already mentioned in Section 8.2, the few percent discrepancies in M_1 and M_2 immediately affect the condensate. Note that the condensate value obtained from the second order coefficient is closer to the bosonization result, the relative error being 1.2 % compared to the 1.8 % at first order. Furthermore, the exact result (8.86) lies *between* our first and second order values, (8.84) and (8.85), so that their mean value, $\langle 0|\bar{\psi}\psi|0\rangle_{1/2} \equiv -0.28442$, is much closer to (8.86), the error being only 0.3%. A similar reduction of errors is actually at work if one considers the ‘pion’ mass-squared of the Schwinger model as a function of m . The first order overshoots the bosonization value while the second order contribution is too small. Adding both one gets closer to the exact result, at least if the mass m is not too tiny. For the value of Fig. 8.1, $m = 0.1$, one finds to second order $M^2(0.1) = 1.39585$, while, in this order, $M_{\text{Bos}}^2(0.1) = 1.39009$. The relative difference is only 0.4%. Possible resolutions of the discrepancies between (8.84), (8.85) and (8.86) will be discussed at the end of this section.

Apart from the ‘pion’ mass squared, the eigenvalue in the ’t Hooft-Bergknoff equation, we have also calculated the associated eigenfunction, the light-cone wave function. The latter has been obtained with high accuracy as can be seen from the good convergence of the alternatively defined mass squared coefficients, $\tilde{M}_i$ towards the variational estimates, M_i (see Section 8.4). If we denote $\phi_0(x) = x(1-x)$, our most accurate variational ansatz for the wave function can be written as

$$\phi[\beta, a, b, c, d] = \phi_0^\beta + a\phi_0^{\beta+1} + b\phi_0^{\beta+2} + c\phi_0^{\beta+3} + d\phi_0^{\beta+4}. \quad (8.87)$$

According to our mass perturbation theory, each of the variational parameters is expanded in powers of the fermion mass m , the coefficients being denoted $\beta_1, \beta_2, \beta_3, a_1, a_2$, etc. From ’t Hooft’s endpoint analysis, β is known exactly. It turns out to be independent of the anomaly (α) and thus is the same for the ’t Hooft and Schwinger model. In Table 8.18 we compare the exact expansion coefficients of β with their variational estimates. The best estimates for the other variational parameters are listed in Table 8.19 for both the ’t Hooft and Schwinger model.

Table 8.18: Comparison of the exact expansion coefficients of the endpoint exponent β with their best variational estimates for both the 't Hooft ($\alpha = 0$) and the Schwinger model ($\alpha = 1$).

	β_1	β_2	β_3
exact	$\sqrt{3}/\pi = 0.55132890$	0	$-\beta_1/10 = -0.05513289$
variational ($\alpha = 0$)	$\sqrt{3}/\pi = 0.55132890$	0.00051	-0.053
variational ($\alpha = 1$)	$\sqrt{3}/\pi = 0.55132890$	0.00063	-0.053

Table 8.19: Best estimates for the variational parameters in the light-cone wave function for both the 't Hooft ($\alpha = 0$) and Schwinger model ($\alpha = 1$).

	a_1	a_2	b_1	b_2	c_1	c_2	d_1	d_2
$\alpha = 0$	0.92	1.19	-2.2	-3.0	7.0	9.2	-9.9	-11.8
$\alpha = 1$	1.42	1.03	-2.7	-3.1	8.8	9.6	-12.5	-12.1


The errors in the coefficients are comparatively large and growing from the left to the right in Table 8.19. Good numerical convergence is evident for the coefficients of a . For the other coefficients our method does not provide enough iteration steps to make convergence explicit. However, the sensitivity of the mass expansion coefficients M_i and the wave functions on the parameters b, c, d is very weak (see Fig. 8.3).

Let us finally investigate possible sources for the few-percent discrepancies in the Schwinger model results like for the condensate values, (8.84) and (8.85) *vs.* (8.86). First of all, it has to be reemphasized that the light-cone calculations are conceptually rather different from the usual treatment based on bosonization within standard equal-time quantization. We are studying the lowest lying meson as a relativistic fermion-anti-fermion bound state. The fermionic degrees of freedom are *explicitly* taken into account as they define our Fock basis. In the bosonized theory, which is a sine-Gordon model with the Lagrangian [199, 164]

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} \mu_0^2 \phi^2 - c m \mu_0 \cos \sqrt{4\pi} \phi, \quad (8.88)$$

the ‘pion’ is the elementary particle described by the scalar field ϕ . One calculates its mass by perturbation theory in the ‘coupling’ $c m \mu_0$, $c = e^2/2\pi$. In the bosonized theory, this ‘coupling’ is actually dependent on the normal-ordering scale used to renormalize the theory. The standard choice is the most natural one, namely the Schwinger boson mass μ_0 , which explicitly appears in the Lagrangian (8.88). It is, however, not very hard to determine the general scale

Figure 8.3: The light-cone wave function of the 't Hooft model 'pion' for $m = 0.1$. The solid curve represents the result from 't Hooft's original ansatz, the dashed curve our best result (with maximum number of variational parameters). At the given resolution, however, the curves of all extensions of 't Hooft's ansatz (a, b, c, d) lie on top of each other.



dependence of quantities calculated within perturbation theory. All one needs is Coleman's re-normal-ordering prescription [199] which for the case at hand is

$$N_{\mu_0} \cos \sqrt{4\pi} \phi = \frac{\mu}{\mu_0} N_{\mu} \cos \sqrt{4\pi} \phi . \quad (8.89)$$

Here, N_{μ_0} and N_{μ} denote normal-ordering with respect to μ_0 and μ , respectively. From (8.89) one derives the following identity for the boson mass squared (setting $\mu_0 = 1$),

$$\begin{aligned} M^2 &= 1 + M_1 m + M_2 m^2 + M_3 m^3 + \dots \\ &= 1 + M'_1 \mu m + M'_2 \mu^2 m^2 + M'_3 \mu^3 m^3 + \dots . \end{aligned} \quad (8.90)$$

In this expression, the M_i denote the coefficients calculated with the standard normal-ordering scale μ_0 , where all tadpoles can be set to zero, while the M'_i are

calculated with normal-ordering scale μ , where one has non-vanishing tadpole contributions. In order that the ‘pion’ mass be scale independent one has to have that (upon reintroducing μ_0)

$$M'_n = \left(\frac{\mu_0}{\mu}\right)^n M_n . \quad (8.91)$$

We have checked this identity explicitly in the first two non-trivial orders of mass perturbation theory with the Lagrangian (8.88). It is also interesting to note that the scale dependence of the coefficients might equivalently be associated with a multiplicative rescaling of the fermion mass m by defining $m' = \mu m$ and using m' as the expansion parameter in (8.90).

Now, the upshot of all this is the following. As we are working entirely in the fermionic representation, we do not a priori know, to which normal-ordering scale of the *bosonized* theory our results correspond. If we assume that it is not the Schwinger mass μ_0 but a different scale μ , we can match the first order coefficients by choosing

$$\mu = \frac{M_1}{M'_1} \mu_0 = \frac{e^\gamma}{\pi/\sqrt{3}} \mu_0 , \quad (8.92)$$

where M'_1 now denotes our light-front result. Quite a similar point of view has been taken by Burkardt in his paper on light-cone quantization of the sine-Gordon model [200]. However, as is obvious from (8.91), the change of scale (8.92) propagates systematically through all coefficients. In particular, if M'_1 is larger than the corresponding bosonization result, the same has to be true for all M'_i . Our coefficient M'_2 , however, already fails this requirement. We thus conclude that a possible scale dependence alone cannot explain the discrepancy.

An additional source of error could still be contributions from higher Fock sectors. We have shown numerically that the first few low orders have very little impact on the expansion coefficients M_i . The contribution from each Fock sector might thus be exponentially small. The whole series, however, if it could be summed up, might add something sizeable. Let us assume, for example, that the contribution of higher Fock sectors to M_2 has the following form³

$$\delta M_2 = \sum_{n=2}^{\infty} c_{2n}(m) , \quad (8.93)$$

where $2n$ is the label of the $2n$ -particle sector. If we had $c_{2n}(m) = e^{-2nm}$, summation of the series would yield $\delta M_2 = 1/2m + O(1)$ and thus effectively a contribution to the first order in m . As we are not able to calculate the real high order behavior of the series in (8.93), we cannot, at the moment, make any definite conclusion about a possible relevance of such summation effects. All we

³We thank M. Burkardt for suggesting the following example.

can say is, that if such effects are present, they must be numerically small, i.e. at the few percent level.

As a first step towards getting some better control of higher particle sectors one would like to develop some tools to estimate their contributions analytically. This amounts to determine the coefficients $c_{2n}(m)$, at least for small n . So far, higher Fock states have only been included numerically.

It might also be interesting to calculate excited states. For the Schwinger model, this again requires the inclusion of higher Fock components as pointed out by Mo and Perry [187]. For the 't Hooft model, on the other hand, one can stay in the two-particle sector. One might then try to modify the variational procedure, basically by taking into account the growing number of nodes, with the aim to match the low energy spectrum with the high energy behavior derived by 't Hooft. Work in these directions is underway.

Chapter 9

The Pion Wave Function in the NJL Model

In this last chapter we will finally study ‘realistic’ light-cone bound-state equations and wave functions. To this end, we enter the world of 3+1 dimensions and investigate the pion wave function within a particularly suited model due to Nambu and Jona-Lasinio (NJL) [201].

The first derivation, analysis and solution of a light-cone bound-state equation appeared in ’t Hooft’s original paper on what is now called the ’t Hooft model [127]. We have discussed this model at length in the last section where we also rederived ’t Hooft’s solution. Interestingly, ’t Hooft did not use the light-cone formalism in the manner we presented it and which nowadays might be called standard. This amounts to deriving the canonical light-cone Hamiltonian and setting up the associated bound-state equation by projecting the Schrödinger equation on the different sectors of Fock space (cf. Chapter 6). Instead, he started from covariant equations, namely the Schwinger-Dyson equations for the quark propagator (or self-energy), and the Bethe-Salpeter equation for the bound state amplitude, which needs the quark self-energy as an input. The light-cone bound equation was then obtained by projecting the Bethe-Salpeter equation onto hypersurfaces of equal light-cone time thereby spoiling explicit covariance. In this way, one avoids to explicitly derive the light-cone Hamiltonian, which can be a tedious enterprise in view of possibly complicated constraints one may have to solve. This makes it worthwhile to have a closer look at this way of proceeding.

In a condensed notation, the Schwinger-Dyson equation for the propagator can be written as

$$S = S_0 + S_0 \Sigma S , \quad (9.1)$$

where S is the full propagator,

$$S(p) = \frac{1}{\not{p} - m_0 - \Sigma} , \quad (9.2)$$

S_0 the free propagator,

$$S_0(p) = \frac{1}{\not{p} - m_0} , \quad (9.3)$$

and Σ the quark self-energy. The Bethe-Salpeter equation for a quark-antiquark bound state (a meson), on the other hand, is

$$\chi_{\text{BS}} = S_1 S_2 K \chi_{\text{BS}} , \quad (9.4)$$

with S_1, S_2 the full propagators of quark and anti-quark, K the Bethe-Salpeter kernel, defined through the interaction one is considering, and χ_{BS} the Bethe-Salpeter amplitude. From the latter, one obtains the light-cone wave function via integration over the energy variable p^- [202, 203, 204, 205],

$$\phi_{\text{LC}}(\mathbf{p}) = \int \frac{dp^-}{2\pi} \chi_{\text{BS}}(p) , \quad \mathbf{p} = (p^+, \mathbf{p}_\perp) . \quad (9.5)$$

The program outlined above, deriving the light-cone bound-state equation from the covariant equations, when applied to QCD, is of course ambitious but should still be feasible in view of the wealth of results already obtained in studying the Schwinger-Dyson equations of QCD [206, 207]. In a first step, which is understood, say, as a feasibility study, a simpler approach will be pursued, namely a calculation within a particularly suited framework, the NJL model. In the language of the Schwinger-Dyson framework, this amounts to using a model gluon propagator that is constant in momentum space. Obviously, such a propagator does not lead to confinement, which however, as pointed out in the introduction, does not necessarily prohibit a reasonable description of hadron structure. The NJL model has been intensively studied within the last decade, and a wealth of results has been obtained (see the recent reviews [208, 209, 210, 211]). To the best of our knowledge, however, light-cone wave functions never have been calculated within the model.

9.1 The Condensate

In its simplest form, the NJL model has a chirally invariant four-fermion interaction, which can be imagined as the result of ‘integrating out’ the gluons in the QCD Lagrangian. Within the instanton model of QCD, this can indeed be done and leads to an NJL type interaction [16]. For two massless ($m_0 = 0$) quark flavors (u and d), the NJL Lagrangian is

$$\mathcal{L}_{\text{NJL}} = \bar{\psi} i \not{\partial} \psi + G [(\bar{\psi} \psi)^2 + (\bar{\psi} i \gamma_5 \boldsymbol{\tau} \psi)^2] . \quad (9.6)$$

In what follows we will mostly suppress flavor (and color) degrees of freedom to keep things as simple as possible. As the coupling G has negative mass dimension ($\sim \text{mass}^{-2}$), the model is not renormalizable and needs a cutoff as a parameter

that has to be fixed by phenomenology. Above a critical coupling, $G > G_c$, a fermion mass is dynamically generated so that chiral symmetry is spontaneously broken corresponding to a second order phase transition. The dynamical fermion mass is proportional to the fermionic condensate in the vacuum as was briefly mentioned in the introduction, Section 1.3.

These results are obtained via a self-consistent mean-field solution of the Schwinger-Dyson equation resulting in the gap equation,

$$\Sigma \equiv m = 2iG\text{tr}S(x=0) = \frac{iGm}{2\pi^4} \int \frac{d^4k}{k^2 - m^2} = -2G\langle 0|\bar{\psi}\psi|0\rangle_m. \quad (9.7)$$

This equation determines the dynamically generated mass m . The integral on the right-hand-side is of course divergent and has to be cut off (see below).

The mean-field solution has a very intuitive explanation. One essentially argues that the main effect of the interaction is to generate the mass of the quarks which become quasi-particles that interact only weakly. Neglecting this interaction entirely, one can view the process of mass generation like in Section 5.2.3 as the transition of quarks with mass $m_0 = 0$ to mass m resulting in a mass term $m\bar{\psi}\psi$ in the Lagrangian (9.6) or in the Hamiltonian. The condensate is obtained via the Feynman-Hellman theorem by differentiating the energy density of the quasi-particle Dirac sea,

$$\begin{aligned} \langle 0|\bar{\psi}\psi|0\rangle_m &= \frac{\partial}{\partial m}\mathcal{E}(m) = \frac{\partial}{\partial m} \int_{-\infty}^0 dk^+ \int \frac{d^2k_\perp}{16\pi^3} \frac{m^2 + k_\perp^2}{k^+} \\ &= -\frac{m}{8\pi^3} \int_0^\infty \frac{dk^+}{k^+} \int d^2k_\perp, \end{aligned} \quad (9.8)$$

Again, as it stands, the integral is divergent and requires regularization. In the most straightforward manner one might choose $m^2/\Lambda \leq k^+ \leq \Lambda$ and $|\mathbf{k}_\perp| \leq \Lambda$, so that the condensate becomes

$$\langle 0|\bar{\psi}\psi|0\rangle_m = -\frac{m}{8\pi^2} \int_{m^2/\Lambda}^\Lambda \frac{dk^+}{k^+} \int_0^{\Lambda^2} d(k_\perp^2) = -\frac{m}{8\pi^2} \Lambda^2 \ln \frac{\Lambda^2}{m^2}. \quad (9.9)$$

Plugging this result into the gap equation (9.7) one finds for the dynamical mass squared,

$$m^2(G) = \Lambda^2 \exp\left(-\frac{4\pi^2}{G\Lambda^2}\right). \quad (9.10)$$

The critical coupling is determined by the vanishing of this mass, $m(G_c) = 0$, and from (9.10) we find the surprising result

$$G_c = 0. \quad (9.11)$$

This result, however, is wrong since one knows from the conventional treatment of the model that the critical coupling is finite of the order π^2/Λ^2 , both for covariant and non-covariant cutoff [201]. In addition, it is quite generally clear that in the *free* theory ($G = 0$) chiral symmetry is not broken (as $m_0 = 0$) and, therefore, this should not happen for arbitrarily small coupling, either (cf. Chapter 7). The remedy is once more to use an information from the ordinary calculation of the condensate. We translate the non-covariant, but rotationally invariant, three-vector cutoff, $|\mathbf{k}| \leq \Lambda$, into light-cone coordinates [81], which leads to

$$0 \leq k_{\perp}^2 \leq 2\Lambda k^+ - m^2 - (k^+)^2, \quad \frac{m^2}{2\Lambda} \leq k^+ \leq 2\Lambda. \quad (9.12)$$

Note that the transverse cutoff becomes a polynomial in k^+ . The $k_{\perp}$ -integration thus has to be performed first. Renaming $2\Lambda \rightarrow \Lambda$ and introducing the longitudinal momentum fraction, $x \equiv k^+/\Lambda$, (9.12) becomes

$$0 \leq k_{\perp}^2 \leq \Lambda^2 x(1-x) - m^2, \quad m^2/\Lambda^2 \leq x \leq 1 - m^2/\Lambda^2. \quad (9.13)$$

Surprisingly, this is just the often used ‘invariant-mass cutoff’,

$$M_0^2 \equiv \frac{k_{\perp}^2 + m^2}{x(1-x)} \leq \Lambda^2, \quad (9.14)$$

with M_0^2 the invariant mass-squared (for a two-particle system) introduced in Chapter 6. Note that the cutoff is also invariant under the exchange $x \leftrightarrow 1-x$.

For the condensate (9.8) the cutoff (9.12) yields an analytic structure different from (9.9),

$$\langle 0 | \bar{\psi} \psi | 0 \rangle_m = -\frac{m}{8\pi^2} \left(2\Lambda^2 - m^2 \ln \frac{\Lambda^2}{m^2} \right), \quad (9.15)$$

where we have neglected sub-leading terms in the cutoff Λ . From (9.15), one infers the correct cutoff dependence of the critical coupling,

$$G_c = \frac{2\pi^2}{\Lambda^2}. \quad (9.16)$$

The moral of this calculation is that even in a non-renormalizable theory like the NJL model, the light-cone regularization prescription is a subtle issue. In order to get a physically sensible result the transverse cutoff has to be k^+ -dependent. Clearly, this dependence cannot be arbitrary but should be constrained from dimensional and symmetry considerations. For renormalizable theories, such arguments have been given by Perry and Wilson [212]. In the example above, it was rotational invariance that solved the problem.

The condensate (9.15) was originally obtained in [81], where, however, a slightly more complicated cut-off was used. In that work, the condensate was defined covariantly in terms of the fermion propagator at the origin, $S(x=0)$

like in (9.7). Performing the energy integration over k^- with the appropriate cutoff again leads to (9.15).

In the NJL model with its second-order phase transition of mean-field type, the usual analogy with magnetic systems can be made. Chiral symmetry corresponds to rotational symmetry, the vacuum energy density to the Gibbs free energy, and the mass m to an external magnetic field. The order parameter measuring the rotational symmetry breaking is the magnetization. It is obtained by differentiating the free energy with respect to the external field. This is the analogue of expression (9.8) as derived from the Feynman-Hellmann theorem.

9.2 The Light-Cone Bound State Equation

Once the physical fermion mass m is known, it can be plugged into the Bethe-Salpeter equation (9.4) which in ladder approximation reads

$$\phi_{\text{LC}}(\mathbf{k}) = \int \frac{dk^-}{2\pi} iS(k) iS(k-P) \int \frac{d^3\tilde{k}}{(2\pi)^3} \int \frac{d\tilde{k}^-}{2\pi} K(k, \tilde{k}) \chi_{\text{BS}}(\tilde{k}) , \quad (9.17)$$

with P denoting the bound-state four-momentum. On the left-hand-side, the projection onto $x^+ = 0$ (i.e. the k^- -integration) has already been carried out. On the right-hand-side, the two integrations over k^- and $\tilde{k}^-$ still have to be performed. Whether this can easily be done depends of course crucially on the kernel K , which in principle is a function both energy variables. For the NJL model, however, one has (after a Fierz transformation),

$$K(k, \tilde{k}) = 2\gamma_5 \otimes \gamma_5 - \gamma_\mu \gamma_5 \otimes \gamma^\mu \gamma_5 , \quad (9.18)$$

i.e. the kernel is momentum independent due to the four-point contact interaction! Thus, the $\tilde{k}^-$ -integration immediately yields ϕ_{LC} , and the k^- -integration can be performed via residue technique and is completely determined by the poles of the propagators, $S(k)$ and $S(k-P)$. As a result, one finds a non-vanishing result only if $0 \leq k^+ \leq P^+$, and *one* of the two particles is put on-shell, e.g. $k^2 = m^2$, as already observed by Gross [213].

The upshot of all this is nothing but the light-cone bound-state equation, which reads explicitly

$$\begin{aligned} \phi_{\text{LC}}(x, \mathbf{k}_\perp) &= -\frac{2G}{x(1-x)} \frac{(\hat{k} + m)\gamma_5(\hat{k} - \not{P} + m)}{M^2 - M_0^2} \\ &\times \int_0^1 dy \int \frac{d^2\tilde{k}_\perp}{8\pi^3} \text{tr} \left[\gamma_5 \phi_{\text{LC}}(y, \tilde{\mathbf{k}}_\perp) \right] \theta_\Lambda(y, \tilde{\mathbf{k}}_\perp) \\ &+ \frac{G}{x(1-x)} \frac{(\hat{k} + m)\gamma_\mu \gamma_5(\hat{k} - \not{P} + m)}{M^2 - M_0^2} \end{aligned}$$

$$\times \int_0^1 dy \int \frac{d^2 \tilde{k}_\perp}{8\pi^3} \text{tr} \left[\gamma^\mu \gamma_5 \phi_{\text{LC}}(y, \tilde{\mathbf{k}}_\perp) \right] \theta_\Lambda(y, \tilde{\mathbf{k}}_\perp) . \quad (9.19)$$

Here we have defined the longitudinal momentum fractions $x = k^+/P^+$, $y = \tilde{k}^+/P^+$, the on-shell momentum $\hat{k} = (\hat{k}^-, \mathbf{k}_\perp, k^+)$ with $\hat{k}^- = (k_\perp^2 + m^2)/k^+$, and the bound state mass squared, $M^2 = P^2$, which is the eigenvalue to be solved for. $\theta_\Lambda(x, \mathbf{k}_\perp)$ denotes the invariant mass cutoff (9.13).

Now, while (9.19) may appear somewhat complicated it is actually very simple; indeed, it is basically already the solution of the problem. The crucial observation is to note that the two integral expressions are mere normalization constants,

$$C_\Lambda \equiv \int_0^1 dy \int \frac{d^2 \tilde{k}_\perp}{8\pi^3} \text{tr} \left[\gamma_5 \phi_{\text{LC}}(y, \tilde{\mathbf{k}}_\perp) \right] \theta_\Lambda(y, \tilde{\mathbf{k}}_\perp) , \quad (9.20)$$

$$D_\Lambda P^\mu \equiv \int_0^1 dy \int \frac{d^2 \tilde{k}_\perp}{8\pi^3} \text{tr} \left[\gamma^\mu \gamma_5 \phi_{\text{LC}}(y, \tilde{\mathbf{k}}_\perp) \right] \theta_\Lambda(y, \tilde{\mathbf{k}}_\perp) . \quad (9.21)$$

Thus, up to normalization, the solution of the light-cone bound-state equation (9.19) is

$$\begin{aligned} \phi_{\text{LC}}(x, \mathbf{k}_\perp) &= -\frac{2GC_\Lambda}{x(1-x)} \frac{(\hat{k} + m)\gamma_5(\hat{k} - \not{P} + m)}{M^2 - M_0^2} \\ &+ \frac{GD_\Lambda}{x(1-x)} \frac{(\hat{k} + m)\not{P}\gamma_5(\hat{k} - \not{P} + m)}{M^2 - M_0^2} . \end{aligned} \quad (9.22)$$

As a first check of our bound state wave function (9.22) we look for a massless pion in the chiral limit. To this end we decompose the light-cone wave function according to [9],

$$\phi_{\text{LC}} = \phi_S + \phi_P \gamma_5 + \phi_A^\mu \gamma_\mu \gamma_5 + \phi_V^\mu \gamma_\mu + \phi_T^{\mu\nu} \sigma_{\mu\nu} . \quad (9.23)$$

Multiplying (9.22) with γ_5 , taking the trace and integrating over $\mathbf{k}$ we find

$$\begin{aligned} C_\Lambda &= -\frac{GC_\Lambda}{2\pi^3} \int_0^1 dx \int d^2 k_\perp \frac{M^2 x + M_0^2(1-x)}{x(1-x)(M^2 - M_0^2)} \theta_\Lambda(x, \mathbf{k}_\perp) \\ &+ \frac{GD_\Lambda}{2\pi^3} M^2 \int_0^1 dx \int d^2 k_\perp \theta_\Lambda(x, \mathbf{k}_\perp) . \end{aligned} \quad (9.24)$$

In the chiral limit one expects a solution for $M = 0$, the Goldstone pion. In this case one obtains

$$1 = \frac{G}{2\pi^3} \int_0^1 dx \int d^2 k_\perp \theta_\Lambda(x, \mathbf{k}_\perp) . \quad (9.25)$$

This is exactly the gap equation (9.7) upon performing the k^- -integration there (with an invariant mass cutoff (9.13) understood in both identities). Note once more the light-cone peculiarity that the (Fock) measure in (9.25) is entirely mass independent. All the mass dependence, therefore, has to come from the (invariant mass) cutoff. Otherwise one will get a wrong behavior of the dynamical mass m as a function of the coupling G , as was the case in (9.10).

With this in mind, we have found that the Goldstone pion is a solution of the light-cone bound-state equation exactly if the gap equation holds. This provides additional evidence for the self-consistency of the procedure. The same results are well known from the covariant treatment [201], and it is gratifying to get them also within the light-cone formalism. The deeper reason for the fact that the quark self-energy and the bound state amplitude satisfy essentially the same equation [214], is the chiral Ward identity relating the quark propagator and the pseudo-scalar vertex [215].

Our next task is to actually evaluate the solution (9.22) of the bound-state equation. ϕ_{LC} is a Dirac matrix and therefore is not yet a light-cone wave function as defined in Chapter 6. The relation between the two quantities has been given by Liu and Soper [204],

$$2P^+ \sqrt{x\bar{x}} \psi(x, \mathbf{k}_\perp, \lambda, \lambda') = \bar{u}(xP^+, \mathbf{k}_\perp, \lambda) \gamma^+ \phi_{\text{LC}}(\mathbf{k}) \gamma^+ v(\bar{x}P^+, -\mathbf{k}_\perp, \lambda'), \quad (9.26)$$

where we have denoted $\bar{x} \equiv 1 - x$ to save space and included a factor $(x\bar{x})^{1/2}$ to match with the conventions of Brodsky and Lepage [82, 216, 217]. A somewhat lengthy calculation, using the spinor identities of App. A, yields the result

$$\psi(x, \mathbf{k}_\perp, \lambda, \lambda') = \frac{2GP^+}{M^2 - M_0^2} \left(\frac{2C_\Lambda}{M} \bar{u}_\lambda M \gamma_5 v_{\lambda'} - D_\Lambda \bar{u}_\lambda \not{P} \gamma_5 v_{\lambda'} \right) \quad (9.27)$$

with the arguments of the spinors $\bar{u}$ and v suppressed. At this point we have to invoke another symmetry principle. Ji et al. have pointed out [93] that the spin structure $(\bar{u}\Gamma v)$ should be consistent with the one obtained from the instant form spinors via a subsequent application of a Melosh transformation [218] and a boost. Using this recipe, one obtains the following relation between the constants C_Λ and D_Λ ,

$$2C_\Lambda/M = -D_\Lambda \equiv N_\Lambda/2GP^+. \quad (9.28)$$

As a result, the spin structure in (9.27) coincides with the standard one used in a number of works [91, 219, 92, 220, 93]. The NJL wave function of the pion finally is

$$\psi(x, \mathbf{k}_\perp, \lambda, \lambda') = \frac{N(\Lambda)}{M^2 - M_0^2} \bar{u}_\lambda (M + \not{P}) \gamma_5 v_{\lambda'} \theta(\Lambda^2 - M_0^2). \quad (9.29)$$

$N(\Lambda)$ is a cutoff dependent normalization defined in (9.28), and the spin (or helicity) structure is given by

$$\begin{aligned} & \bar{u}(xP^+, \mathbf{k}_\perp, \lambda) (M + \not{P}) \gamma_5 v(\bar{x}P^+, -\mathbf{k}_\perp, \lambda') = \\ & = \frac{1}{\sqrt{x\bar{x}}} \left\{ \lambda \left[mM + m^2 - \mathbf{k}_\perp^2 + M^2 x\bar{x} \right] \delta_{\lambda, -\lambda'} - k_{-\lambda} (M + 2m) \delta_{\lambda\lambda'} \right\} \end{aligned} \quad (9.30)$$

where we have defined $k_\lambda \equiv k^1 + i\lambda k^2$. The first term with spins anti-parallel corresponds to $L_z = 0$, the second one (with spins parallel) to $L_z = \pm 1$. It has already been pointed out by Leutwyler that both spin alignments should contribute to the pion wave function [221, 222]. Note that the latter is a cutoff dependent quantity. This is necessary in order to make the wave function normalizable. In other words, it guarantees the boundary conditions (6.23) which state that the wave function drops off sufficiently fast in x and $k_\perp$.

As we are interested in analyzing the quality of a constituent picture, we approximate the pion by its valence state,

$$|\pi : \mathbf{P}\rangle = \sum_{\lambda, \lambda'} \int_0^1 \frac{dx}{\sqrt{x(1-x)}} \int \frac{d^2 k_\perp}{16\pi^3} \psi(x, \mathbf{k}_\perp, \lambda, \lambda') |q\bar{q} : x, \mathbf{k}_\perp, \lambda, \lambda'\rangle \quad (9.31)$$

which should be compared with the general expression (6.13). The normalization of this state is given by (6.10),

$$\langle \pi : \mathbf{P}' | \pi : \mathbf{P} \rangle = 16\pi^3 P^+ \delta^3(\mathbf{P} - \mathbf{P}') . \quad (9.32)$$

As usual we work in a frame in which the total transverse momentum vanishes, i.e. $\mathbf{P} = (P^+, \mathbf{P}_\perp = 0)$. Expression (9.32) yields the normalization (6.19) of the wave function,

$$\sum_{\lambda, \lambda'} \int_0^1 dx \int \frac{d^2 k_\perp}{16\pi^3} |\psi(x, \mathbf{k}_\perp, \lambda, \lambda')|^2 = 1 . \quad (9.33)$$

It is of course a critical assumption that the probability to find the pion in its valence state is one. In this way we enforce a constituent picture *by fiat*, and it is clear that such an assumption has to be explicitly checked. An alternative way of proceeding would be not to fix the normalization but to use $N(\Lambda)$ as an additional parameter that has to be fixed by phenomenology. We will, however, use (9.33) and check whether a valence (or constituent) picture makes sense.

9.3 Observables

With the light-cone wave function at hand, we are in the position to calculate observables. As a technical prelude, however, we first have to determine its normalization. To keep things simple we consider the chiral limit, $M = 0$. We write the pion wave function as a matrix in helicity space,

$$\psi(x, \mathbf{k}_\perp) = -N(\Lambda) \frac{\sqrt{x(1-x)}}{k_\perp^2 + m^2} \begin{pmatrix} -2m(k^1 + ik^2) & m^2 - k_\perp^2 \\ k_\perp^2 - m^2 & -2m(k^1 - ik^2) \end{pmatrix} \theta(\Lambda^2 - M_0^2) \quad (9.34)$$

The diagonal terms correspond to parallel spins, the off-diagonal ones to anti-parallel spins.

The normalization (9.33) determines N as a function of Λ . Using the explicit form of the invariant-mass cutoff, (9.13), one finds

$$1 = \frac{|N|^2}{8\pi^2} \int_{\epsilon^2}^{1-\epsilon^2} dx x(1-x) \int_0^{\Lambda^2(x)} dk_{\perp}^2, \quad (9.35)$$

where we have defined

$$\epsilon^2 \equiv m^2/\Lambda^2 \quad (9.36)$$

$$\Lambda^2(x) \equiv \Lambda^2 x(1-x) - m^2 = \Lambda^2 [x(1-x) - \epsilon^2]. \quad (9.37)$$

As an additional approximation we will neglect terms of order ϵ^2 , i.e. we assume that the cutoff is large compared to the constituent mass. From the standard values, $\Lambda \simeq 1$ GeV, $m \simeq 300$ MeV, we expect that this assumption should induce an error of the order of 10%. The advantage is that it allows for a simple analytic evaluation of all the integrals we will encounter. Furthermore, the leading order will be independent of the actual value of the constituent mass. From (9.35) one thus finds the normalization,

$$|N|^2 = \frac{240\pi^2}{\Lambda^2} [1 + 5\epsilon^2 + O(\epsilon^4)]. \quad (9.38)$$

Choosing N real we obtain to leading order,

$$N = 2\sqrt{15}\pi/\Lambda. \quad (9.39)$$

This fixes our normalization constant. We proceed by calculating the pion electromagnetic formfactor. It is defined by the matrix element of the electromagnetic current J_{em}^{μ} between pion states,

$$\langle \pi : \mathbf{P} | J_{\text{em}}^{\mu} | \pi : \mathbf{P}' \rangle = 2(P + P')^{\mu} F(Q^2), \quad Q^2 \equiv -(P - P')^2. \quad (9.40)$$

Considering $\mu = +$ in a frame where $\mathbf{P} = (P^+, \mathbf{0})$ and $\mathbf{P}' = (P^+, \mathbf{q}_{\perp})$ one is led to the the Drell-Yan formula [223, 82],

$$F(\mathbf{q}_{\perp}^2) = \sum_{\lambda\lambda'} \int_0^1 dx \int \frac{d^2 k_{\perp}}{16\pi^3} \psi^*(x, \mathbf{k}'_{\perp}, \lambda, \lambda') \psi(x, \mathbf{k}_{\perp}, \lambda, \lambda'). \quad (9.41)$$

The transverse momentum of the struck quark is $\mathbf{k}'_{\perp} = \mathbf{k}_{\perp} + x\mathbf{q}_{\perp}$. Note that the formula (9.41) with its overlap of two wave functions on the right-hand-side is rather similar to the non-relativistic result [82]. If one sets the momentum transfer $\mathbf{q}_{\perp}$ equal to zero, one is left with the normalization integral (9.33), so that the form factor is automatically normalized to one. Plugging the pion wave function (9.34) into (9.41), one finds to first order in $\mathbf{q}_{\perp}^2$,

$$F(\mathbf{q}_{\perp}^2) = 1 - \mathbf{q}_{\perp}^2 \frac{|N|^2}{80\pi^2} \left[1 - \frac{20}{3}\epsilon^2 + O(\epsilon^4) \right]. \quad (9.42)$$

Inserting the normalization (9.38), the formfactor becomes

$$F(\mathbf{q}_\perp^2) = 1 - \frac{3}{\Lambda^2} \mathbf{q}_\perp^2 \left[1 - \frac{5}{3} \epsilon^2 + O(\epsilon^4) \right]. \quad (9.43)$$

This leading order result already determines a particular observable, namely the charge radius of the pion, which is defined via the slope of the formfactor at zero momentum transfer,

$$F(\mathbf{q}_\perp^2) = 1 - \frac{\langle r_\pi^2 \rangle}{6} \mathbf{q}_\perp^2 + O(\mathbf{q}_\perp^4). \quad (9.44)$$

To leading order in ϵ^2 the pion charge radius thus becomes

$$\langle r_\pi^2 \rangle = 18/\Lambda^2, \quad (9.45)$$

and is thus entirely determined by the cutoff Λ . We might fix the latter by using the experimental result [224]

$$\langle r_\pi^2 \rangle = 0.44 \text{ fm}^2. \quad (9.46)$$

The cutoff thus becomes (to leading order in ϵ^2),

$$\Lambda^2 = \frac{18}{\langle r_\pi^2 \rangle} \simeq (1.3 \text{ GeV})^2. \quad (9.47)$$

Our preferred way, however, to fix the cutoff is to use the pion decay constant $f_\pi = 92.4 \text{ MeV}$ [225], which is known to a higher accuracy than the charge radius. Though this does not matter in view of our present approximations, it seems to be advantageous at least in principle. f_π is given by the ‘wave function at the origin’, ψ_0 , defined as

$$\psi_0 \equiv \int_0^1 dx \int \frac{d^2 k_\perp}{16\pi^3} \psi(x, \mathbf{k}_\perp, \uparrow, \downarrow) \theta(\Lambda^2 - M_0^2) = \frac{3N\Lambda^2}{2048\pi} + O(\epsilon^2). \quad (9.48)$$


Note that only the $L_z = 0$ component contributes. Using the normalization (9.38) this becomes

$$\psi_0 = \frac{3\sqrt{15}}{512} \Lambda \simeq 0.022 \Lambda. \quad (9.49)$$

The pion decay constant has been obtained from weak π decay [82] where only the pion valence wave functions contributes to the decay matrix elements. Therefore, the following result [82, 217] provides an *exact* constraint on the pion wave function,

$$\psi_0 = \frac{f_\pi}{2\sqrt{N_C}}. \quad (9.50)$$

Figure 9.1: The pion formfactor squared vs. momentum transfer $q^2 \equiv q_\perp^2$. The full line is the monopole fit of [224], $|F|^2 = n/(1 + q_\perp^2 \langle r_\pi^2 \rangle / 6)^2$ with $n = 0.991$, $\langle r_\pi^2 \rangle = 0.431 \text{ fm}^2$; the dashed line is the same fit with our values, $n = 1$, $\langle r_\pi^2 \rangle = 0.50 \text{ fm}^2$. The agreement is consistent with the expected accuracy of 10%.



Comparing with (9.49) we find that the cutoff is fixed through

$$\Lambda = \frac{256}{9\sqrt{5}} f_\pi \simeq 12.7 f_\pi \simeq 1.2 \text{ GeV} . \quad (9.51)$$

Having fixed the cutoff, we are in the position to actually predict the pion charge radius via (9.45). The result is

$$\langle r_\pi^2 \rangle = \frac{18}{\Lambda^2} = 0.50 \text{ fm}^2 = (0.71 \text{ fm})^2 . \quad (9.52)$$

which differs from the experimental value (9.46) by 13% and therefore is consistent with the expected accuracy.

This is also evident from Fig. 9.1 where we compare with the experimental results of [224]. The full and dashed line correspond to monopole fits of the form

$$F^2(\mathbf{q}_\perp^2) = \frac{n}{(1 + \mathbf{q}_\perp^2 \langle r_\pi^2 \rangle / 6)^2} , \quad (9.53)$$

with $n = 0.991$, $\langle r_\pi^2 \rangle = 0.431 \text{ fm}^2$ [224] and our result (9.52), $n = 1$, $\langle r_\pi^2 \rangle = 0.50 \text{ fm}^2$, respectively.

If we compare our result (9.51) for the cutoff with the value for the cutoff Λ_χ below which chiral effective theories make sense, $\Lambda_\chi = 4\pi f_\pi \simeq 12.6 f_\pi$ [226], we find that the agreement is surprisingly good. Clearly, this is somewhat accidental, as we do not expect an accuracy better than 10%. This can be seen explicitly again by alternatively using the pion charge radius to fix f_π . Plugging (9.47) into the general relation (9.51) between cutoff and f_π , and solving for f_π , we predict

$$f_\pi = \frac{9\sqrt{5}}{256} \Lambda \simeq 0.077 \Lambda \simeq 100 \text{ MeV} , \quad (9.54)$$

which indeed is accurate to within 10%.

We proceed by calculating the mean transverse momentum,

$$\langle \mathbf{k}_\perp^2 \rangle = \sum_{\lambda\lambda'} \int_0^1 dx \int \frac{d^2 k_\perp}{16\pi^3} \mathbf{k}_\perp^2 |\psi(x, \mathbf{k}_\perp, \lambda, \lambda')|^2 \theta(\Lambda^2 - M_0^2) . \quad (9.55)$$

To leading order in ϵ^2 this becomes

$$\langle \mathbf{k}_\perp^2 \rangle = \frac{|N|^2}{2240\pi^2} \Lambda^4 , \quad (9.56)$$

or, upon inserting the normalization (9.38),

$$\langle \mathbf{k}_\perp^2 \rangle = \frac{3}{28} \Lambda^2 \simeq (0.43 \text{ GeV})^2 . \quad (9.57)$$

The r.m.s. transverse momentum is thus of the order of 400 MeV, which is a reasonable value [221]. In particular, this value shows that the pion is a highly relativistic system as the momenta of its (valence) constituents exceed the pion mass.

The inverse of the r.m.s. momentum can be identified with the pion core radius, for which one finds

$$R_\pi \equiv \langle \mathbf{k}_\perp^2 \rangle^{-1/2} = \sqrt{28/3} \Lambda^{-1} \simeq 0.47 \text{ fm} . \quad (9.58)$$

As expected, it is smaller than the charge radius which we attribute to the fact that the charge distribution measured by the charge radius does not coincide with the distribution of baryon density. The core radius R_π is sometimes related to the decay constant f_π via the dimensionless quantity [227]

$$C \equiv f_\pi R_\pi . \quad (9.59)$$

In constituent quark models one typically gets

$$f_\pi = \frac{\sqrt{3}m}{2\pi} , \quad (9.60)$$

and if the quarks are confined by a harmonic oscillator potential, their mass is $m = 3/2R_\pi$, so that $C \simeq 0.4$. This implies the fairly large value $R_\pi \simeq 0.8$ fm. Using standard many-body techniques, Bernard et al. have calculated this quantity in a model treating the pion as a collective excitation of the QCD vacuum, and find $C \simeq 0.2$ [228]. This result is close to what we get from (9.51) and (9.58),

$$C = \frac{9\sqrt{5}}{256} \sqrt{\frac{28}{3}} \simeq 0.24 . \quad (9.61)$$

Thus, though we work within a constituent picture, we do get a reasonable value. We believe that this is due to the intrinsic consistency of the light-cone framework with the requirements of relativity.

9.4 The Pion Distribution Amplitude

We follow Brodsky and Lepage [82, 216] and define the pion distribution amplitude by integrating the wave function over $k_\perp$. Due to the cutoff dependence of the wave function, the integration extends effectively only up to the cutoff scale $\Lambda(x)$,

$$\phi_{\text{BL}}(x) \equiv \int \frac{d^2 k_\perp}{16\pi^3} \psi_\Lambda(x, \mathbf{k}_\perp, \lambda, \lambda') . \quad (9.62)$$

As usual, we evaluate this to leading order in ϵ so that the result becomes independent of the constituent mass,

$$\phi_{\text{BL}}(x) = \frac{\sqrt{15}}{4\pi} \Lambda [x(1-x)]^{3/2} = \frac{64\sqrt{3}}{9\pi} f_\pi [x(1-x)]^{3/2} . \quad (9.63)$$

The normalization of ϕ is obtained by integrating this over x , which simply yields the ‘wave function at the origin’, ψ_0 , defined in (9.48). We have used the subscript ‘BL’ (for Brodsky-Lepage), because in what follows we will use a distribution amplitude which is normalized to one, that is,

$$\phi(x) \equiv \frac{2\sqrt{3}}{f_\pi} \phi_{\text{BL}} = \frac{128}{3\pi} [x(1-x)]^{3/2} . \quad (9.64)$$

It has been shown by Brodsky and Lepage, that the distribution amplitude obeys an evolution equation in momentum scale Q^2 which can be solved exactly in the limit $Q^2 \rightarrow \infty$. The result is called the *asymptotic* distribution amplitude,

$$\phi_{\text{as}}(x) = 6x(1-x) . \quad (9.65)$$

The (non-asymptotic) pion distribution amplitude has been a rather controversial object. For a while people have tended to believe in a ‘double-humped’ shape of the amplitude (due to a factor $(1-2x)^2$), which was originally found by Chernyak and Zhitnitsky via QCD sum rules [229]. Recent CLEO data, however, [230] seem to support an amplitude that is not too different from the asymptotic one [231]. It turns out, that our result (9.64) is qualitatively consistent with these findings [232, 233, 234]. Belyaev and Johnson have recently reported two constraints which should be satisfied by the distribution amplitude [232],

$$\begin{aligned} \phi(x=0.3) &= 1 \pm 0.2 , \\ \phi(x=0.5) &= 1.25 \pm 0.25 . \end{aligned} \quad (9.66)$$


In Figure 9.2 we have displayed our distribution amplitude in comparison with the asymptotic one. The vertical lines represent the constraints (9.66). Obviously, these are not satisfied, as we have

$$\phi(x=0.3) = 1.31 , \quad (9.67)$$

$$\phi(x=0.5) = 1.70 . \quad (9.68)$$

As we already overshoot the asymptotic amplitude (for $0.3 < x < 0.7$) this does not come as a surprise. In order to improve on that we presumably have to increase our accuracy by taking into account finite pion and quark masses. Still, we get an amplitude which is at least qualitatively correct.

Figure 9.2: *The pion distribution amplitude. The full curve represents the asymptotic amplitude, the broken one the amplitude calculated within the NJL model. The vertical lines mark the constraints (9.66) of Belyaev and Johnson.*



It should also be mentioned that it is not obvious what the actual momentum scale is to which our result corresponds. Our transverse-momentum cutoff is x -dependent, $\Lambda^2(x) \simeq \Lambda^2 x(1-x)$, so, if we choose an average x of $\langle x \rangle \simeq 1/2$, a natural scale seems to be¹

$$\mu \equiv \left[\langle x \rangle (1 - \langle x \rangle) \right]^{1/2} \Lambda \simeq \Lambda/2 \simeq 600 \text{ MeV} . \quad (9.69)$$

This is somewhat lower than $\mu \simeq 1 \text{ GeV}$, which has been assumed by Belyaev and Johnson in their analysis of the distribution amplitude in terms of light-cone quark models [233].

¹We thank W. Schweiger for discussions on this point.

Conclusions

In this report we have discussed an alternative approach to relativistic (quantum) physics based on light-cone dynamics. It makes use of the fact that for relativistic systems the choice of the time parameter is not unique. Our particular choice of relativistic dynamics singles out Dirac's front form so that null-planes tangent to the light-cone become hypersurfaces of equal time. This apparently trivial change of coordinates has far-reaching consequences:

- The number of kinematical (i.e. interaction independent) Poincaré generators becomes maximal; there are seven of them instead of the usual six, among them the boosts.
- Lorentz boosts in z -direction become diagonal; the light-cone time and space coordinates, x^+ and x^- , respectively, do not get mixed but rather get rescaled.
- As a consequence, for many-particle systems one can introduce frame-independent relative coordinates, the longitudinal momentum fractions, x_i , and the relative transverse momenta, $\mathbf{k}_{\perp i}$.
- Because of a two-dimensional Galilei invariance, relative and center-of-mass motion separate. As a result, many formulae are reminiscent of non-relativistic physics and thus very intuitive.
- This is particularly true for light-cone wave functions which, because of the last two properties, are boost invariant and do not depend on the total momentum of the bound state. They are therefore ideal tools to study relativistic particle systems.
- The last statement even holds for relativistic quantum field theory where one combines the unique properties of light-cone quantization with a Fock space picture. The central feature making this a reasonable idea is the triviality of the light-cone vacuum which accordingly is an eigenstate of the fully interacting Hamiltonian. This implies that the Fock operators create the *physical* particles from the ground state.

As expected, however, the principle of ‘conservation of difficulties’ is at work so that there are problems to overcome. We have focused on the issue of vacuum properties. In instant-form quantum field theory, in particular in QCD, many non-perturbative phenomena are attributed to the non-triviality of the vacuum which shows up via the appearance of condensates. These are non-vanishing vacuum expectation values which support the idea that the instant-form vacuum is a complicated many-body state (like, e.g. the BCS ground state). Of further physical relevance is the fact that many of these condensates signal the spontaneous breakdown of a symmetry. The conceptual problem which arises at this point is how to reconcile the existence of condensates with the triviality of the light-cone vacuum.

We have shown that for scalar field theory in 1+1 dimensions, the field mode with longitudinal momentum k^+ is the carrier of non-trivial vacuum properties. Its c -number part coincides with the vacuum expectation value of the field. The reflection symmetry, mapping the field onto its negative, is thus seen to be spontaneously broken.

For fermionic fields with a bilinear condensate reflecting the spontaneous breakdown of chiral symmetry, we had to proceed differently. We calculated the bound state spectrum of model field theories and reconstructed ground state properties like the condensates from the particle spectrum. The paradigm identity was the Gell-Mann-Oakes-Renner relation which relates the masses of the bound states and the fermions to the condensates. Alternatively, one can try to derive gap equations like in the NJL model, which determine the mass of the dressed, effective fermions in terms of the condensate.

It is crucial for the latter approach to solve the associated bound-state problem, or, in other words, the light-cone Schrödinger equation. For a relativistic quantum field theory, this, in principle, amounts to solving an infinite system of coupled integral equations for the amplitudes to find an ever increasing number of constituents in the bound state. Experience, however, shows that the *light-cone* amplitudes to find more than the valence quanta in the bound state tend to become rather small. We have confirmed this fact for the theories we have studied. Note that the same is *not* true within ordinary, that is, instant-form quantization. The light-cone approach, however, provides some confidence that also in real QCD a constituent picture may emerge, so that hadrons are consistently described as light-cone bound states of a minimal number of two or three constituents.

A first step in this direction was performed using an effective field theory, the NJL model. We have seen that, though we made a number of approximations, in particular by *enforcing* a constituent picture, a number of pionic observables are predicted with reasonable accuracy.

At this point one has to ask whether the ideas just outlined can really be used in QCD. This is a difficult question. The reason why light-cone QCD has not been presented in this report is that, in our opinion, there are still technical prob-

lems to overcome. First of all, the quantization procedure itself is not entirely clear beyond perturbation theory. There are the usual gauge redundancies to be taken into account *together* with the peculiar first-order nature of the light-cone Lagrangian. We have reported on this elsewhere [235]. One generally uses the light-cone gauge $A^+ = 0$, a variant of an axial gauge which has its own well-known problems, among them residual gauge invariance and infrared divergences. However, it is only in this gauge that the canonical quantization is straightforward. The latter has been performed quite some time ago [236, 237, 217] and successfully been used as the basis of light-cone perturbation theory. Some impressive achievements of the latter method can be found in the recent review [238] which focuses on the light-cone formulation of gauge theories.

Still, even if one wants to utilize perturbation theory for bound-state calculations (in case this is justified) one is confronted with difficulties. One big problem is to find a systematic renormalization program in order to get rid of the ubiquitous infinities. As one does neither have explicit covariance nor rotational invariance there are very few guiding principles and restrictions. Accordingly, there is a rather large variety of divergences which can even become nonlocal, power counting gets more complicated, and the number of counter-terms tends to become large [137]. As a result, to our knowledge, the light-cone renormalization program has not been pushed beyond one loop so far.

We therefore consider it worthwhile to first pursue a less ambitious program, the first steps of which have been undertaken in the last chapter. One should make use of effective field theories, that are obtained from QCD in a well-defined manner [239, 240], and in general are not renormalizable. The cutoff is a parameter of the theory which sets the energy scale below which the theory makes sense. Thus, one only has to deal with the issue of regularization to render the appearing integrals finite. Still, for the case of the NJL model, we have seen that already this can be a nontrivial problem. An alternative candidate suited for quark bound-state studies is the effective quark theory derived within the QCD instanton model [16]. It is also of the NJL type, however, with a more complicated (and presumably more realistic) interaction.

Further improvements can be achieved by using more elaborate kernels in the covariant Bethe-Salpeter equation. These can be found in the literature on the Schwinger-Dyson approach to QCD [206, 207] and basically amount to choosing a model for the gluon propagator. The NJL model corresponds to an infrared-finite propagator which is constant in momentum space. Therefore, one does not have confinement. The latter can be implemented by using a gluon propagator which diverges for small momenta. Other constraints follow from multiplicative renormalizability. Clearly, the procedure of deriving the light-cone bound-state equation will no longer be as simple as in the last chapter but should still be feasible. Work in this direction is underway.

Acknowledgments

It is a pleasure to thank my former supervisor and present collaborator E. Werner for the continuous support and encouragement which he has provided over the years. Furthermore, I am grateful to U. Heinz for some valuable advice concerning this work. I thank my Regensburg colleagues for their help, friendship and company, in particular during mountain hiking: G. Fischer, T. Hirschmann, A. Marquardt, M. Meister, M. Oleszczuk, T. Pause, S. Simbürger, C. Skornia, C. Stern and B. Zellermann. Warm thanks go also to the non-scientific staff, G. Heyer and C. Leichtl, for raising my spirits with birthday cakes and the like.

I am indebted to my colleagues in the light-cone community for discussions and inspiration, the junior ones: M. Burkardt, K. Itakura, A. Kalloniatis, D. Robertson, B. van de Sande, U. Trittman, and S. Tsujimaru, as well as the senior ones: A. Bassetto, P. Grangé, J. Hiller, G. McCartor, H.-C. Pauli, R. Perry, and S. Pinsky. Special thanks are due to K. Harada for his collaboration on the 't Hooft and Schwinger model.

Appendix A

Light-Cone Spinors

In this appendix we collect some basic formulae representing our conventions for light-cone spinors in $d = 3+1$, together with some frequently occurring identities involving these. Our conventions for Dirac matrices are those of Bjorken and Drell [17].

Let $\psi(x)$ be a solution of the free Dirac equation

$$(i\cancel{\partial} - m)\psi(x) = 0 \quad . \quad (\text{A.1})$$

The Fock expansion of ψ reads

$$\psi(\mathbf{x}, 0) = \sum_{\lambda} \int_0^{\infty} \frac{dk^+}{k^+} \int \frac{d^2 k_{\perp}}{16\pi^3} \left[b(\mathbf{k}, \lambda) u(\mathbf{k}, \lambda) e^{-i\mathbf{k} \cdot \mathbf{x}} + d^{\dagger}(\mathbf{k}, \lambda) v(\mathbf{k}, \lambda) e^{i\mathbf{k} \cdot \mathbf{x}} \right] , \quad (\text{A.2})$$

where

$$\mathbf{k} \equiv (k^+, \mathbf{k}_{\perp}) , \quad \mathbf{x} \equiv (x^-, \mathbf{x}_{\perp}) , \quad \mathbf{k} \cdot \mathbf{x} \equiv \frac{1}{2} k^+ x^- - \mathbf{k}_{\perp} \cdot \mathbf{x}_{\perp} . \quad (\text{A.3})$$

The projections $\psi_{\pm}$ are obtained through the action of the projectors $\Lambda_{\pm}$, $\psi_{\pm} \equiv \Lambda_{\pm} \psi$, with (in a two-by-two block notation)

$$\Lambda_{\pm} \equiv \frac{1}{4} \gamma^{\mp} \gamma^{\pm} = \frac{1}{2} \begin{pmatrix} \mathbb{1} & \pm \sigma_3 \\ \pm \sigma_3 & \mathbb{1} \end{pmatrix} . \quad (\text{A.4})$$

As Λ_+ and Λ_- are projectors they obey

$$\Lambda_{\pm}^2 = \Lambda_{\pm} , \quad \Lambda_{\pm}^{\dagger} = \Lambda_{\pm} , \quad (\text{A.5})$$

and

$$\Lambda_+ + \Lambda_- = 1 , \quad \Lambda_+ \Lambda_- = \Lambda_- \Lambda_+ = 0 . \quad (\text{A.6})$$

The following identities are frequently used ($i = 1, 2$),

$$\Lambda_{\pm} \beta = \beta \Lambda_{\mp} , \quad (\text{A.7})$$

$$\Lambda_{\pm} \gamma^i = \gamma^i \Lambda_{\pm} , \quad (\text{A.8})$$

$$\Lambda_{\pm} \alpha^i = \alpha^i \Lambda_{\mp} . \quad (\text{A.9})$$

The spinors u and v in (A.2) satisfy the equations

$$(\not{k} - m) u(\mathbf{k}, \lambda) = 0, \quad (\text{A.10})$$

$$(\not{k} + m) v(\mathbf{k}, \lambda) = 0, \quad (\text{A.11})$$

and are explicitly given by [241, 216, 86]

$$u(\mathbf{k}, \lambda) = \frac{1}{\sqrt{k^+}} (k^+ + \beta m + \alpha_i k_i) X_\lambda, \quad (\text{A.12})$$

$$v(\mathbf{k}, \lambda) = \frac{1}{\sqrt{k^+}} (k^+ - \beta m + \alpha_i k_i) X_{-\lambda}. \quad (\text{A.13})$$

The X_λ are eigenspinors of the projector Λ_+ with eigenvalue one,

$$\Lambda_+ X_\lambda = X_\lambda, \quad X_\lambda^\dagger X_{\lambda'} = \delta_{\lambda\lambda'}, \quad (\text{A.14})$$

or explicitly,

$$X_1 \equiv X_\uparrow = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad X_{-1} \equiv X_\downarrow = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}. \quad (\text{A.15})$$

If we introduce the two-spinors χ_λ , the usual eigenspinors of σ_3 ,

$$\chi_1 \equiv \chi_\uparrow = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \chi_{-1} \equiv \chi_\downarrow = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad (\text{A.16})$$

we can write for X_λ ,

$$X_\lambda = \frac{1}{\sqrt{2}} \begin{bmatrix} \chi_\lambda \\ \lambda \chi_\lambda \end{bmatrix}. \quad (\text{A.17})$$

Using the identities

$$\Lambda_\pm \begin{bmatrix} \chi_\lambda \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} \chi_\lambda \\ \pm \lambda \chi_\lambda \end{bmatrix}, \quad \Lambda_\pm \begin{bmatrix} 0 \\ -\lambda \chi_\lambda \end{bmatrix} = \frac{1}{2} \begin{bmatrix} \pm \chi_{-\lambda} \\ -\lambda \chi_{-\lambda} \end{bmatrix}, \quad (\text{A.18})$$

we can rewrite the elementary spinors (A.12, A.13) according to

$$u(\mathbf{k}, \lambda) = \sqrt{\frac{2}{k^+}} [m \Lambda_- + (k^+ + \alpha^i k_i) \Lambda_+] \begin{bmatrix} \chi_\lambda \\ 0 \end{bmatrix}, \quad (\text{A.19})$$

$$v(\mathbf{k}, \lambda) = \sqrt{\frac{2}{k^+}} [m \Lambda_- + (k^+ + \alpha^i k_i) \Lambda_+] \begin{bmatrix} 0 \\ -\lambda \chi_{-\lambda} \end{bmatrix}. \quad (\text{A.20})$$

The projections of the spinors u and v are

$$u_+(\mathbf{k}, \lambda) = \sqrt{k^+} X_\lambda \equiv u_+(k^+, \lambda) , \quad (\text{A.21})$$

$$v_+(\mathbf{k}, \lambda) = \sqrt{k^+} X_{-\lambda} \equiv v_+(k^+, \lambda) , \quad (\text{A.22})$$

$$u_-(\mathbf{k}, \lambda) = \frac{1}{\sqrt{k^+}} (m\beta + \alpha_i k_i) X_\lambda , \quad (\text{A.23})$$

$$v_-(\mathbf{k}, \lambda) = \frac{1}{\sqrt{k^+}} (-m\beta + \alpha_i k_i) X_{-\lambda} . \quad (\text{A.24})$$

One has the completeness relation in the ‘+’-sector,

$$\sum_\lambda u_+(\mathbf{k}, \lambda) u_+^\dagger(\mathbf{k}, \lambda) = \sum_\lambda v_+(\mathbf{k}, \lambda) v_+^\dagger(\mathbf{k}, \lambda) = k^+ \Lambda_+ . \quad (\text{A.25})$$

The following provides a (very incomplete) list of matrix elements (for further matrix elements, see [216]),

$$u_+^\dagger(k^+, \lambda) u_+(k^+, \lambda') = v_+^\dagger(k^+, \lambda) v_+(k^+, \lambda') = k^+ \delta_{\lambda\lambda'} , \quad (\text{A.26})$$

$$u_\pm^\dagger(\mathbf{k}, \lambda) \beta u_\mp(\mathbf{k}, \lambda') = m \delta_{\lambda\lambda'} \mp \lambda k_{-\lambda} \delta_{\lambda, -\lambda'} , \quad (\text{A.27})$$

$$v_\pm^\dagger(\mathbf{k}, \lambda) \beta v_\mp(\mathbf{k}, \lambda') = -m \delta_{\lambda\lambda'} \pm \lambda k_\lambda \delta_{-\lambda, \lambda'} , \quad (\text{A.28})$$

where we have defined

$$k_\lambda \equiv k^1 + i\lambda k^2 , \quad \text{so that} \quad k_\lambda k_{-\lambda} = k_\perp^2 . \quad (\text{A.29})$$

Introducing

$$\gamma_5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix} , \quad (\text{A.30})$$

one finds that X_λ is also an eigenspinor of γ_5 ,

$$\gamma_5 X_\lambda = \lambda X_\lambda . \quad (\text{A.31})$$

Furthermore, one has

$$k^i \gamma^i X_\lambda = \lambda k_\lambda X_{-\lambda} . \quad (\text{A.32})$$

The last two identities lead to the matrix elements

$$u_+^\dagger(k^+, \lambda) \gamma_5 v_+(p^+, \lambda') = \sqrt{k^+ p^+} \lambda \delta_{\lambda, -\lambda'} , \quad (\text{A.33})$$

$$u_+^\dagger(k^+, \lambda) k^i \gamma^i \gamma_5 v_+(p^+, \lambda') = \sqrt{k^+ p^+} k_{-\lambda} \delta_{\lambda\lambda'} , \quad (\text{A.34})$$

which have been used in Chapter 9 to determine the spin structure of the pion wave function.

Bibliography

- [1] R. M. Barnett *et al.*, Phys. Rev. **D54**, 1 (1996).
- [2] M. Gell-Mann, Phys. Lett. **8**, 214 (1964).
- [3] G. Zweig, (1964), CERN report 8419/TH 412.
- [4] M. Gell-Mann and Y. Ne'eman, *The Eightfold Way* (Benjamin, New York, 1964).
- [5] D. Gross and F. Wilczek, Phys. Rev. Lett. **30**, 1343 (1973).
- [6] S. Weinberg, Phys. Rev. Lett. **31**, 494 (1973).
- [7] H. Fritzsch, M. Gell-Mann, and H. Leutwyler, Phys. Lett. **B47**, 365 (1973).
- [8] I. Montvay and G. Münster, *Quantum fields on a lattice* (Cambridge University Press, Cambridge, 1994).
- [9] W. Lucha, F. Schöberl, and D. Gromes, Phys. Rept. **200**, 127 (1991).
- [10] S. Mukherjee *et al.*, Phys. Rept. **231**, 201 (1993).
- [11] S. Chakrabarty and S. Deoghuria, Prog. Part. Nucl. Phys. **33**, 577 (1994).
- [12] M. Neubert, Phys. Rept. **245**, 259 (1994), hep-ph/9306320.
- [13] H. Dosch and S. Narison, Phys. Lett. **B417**, 173 (1998).
- [14] J. Goldstone, Nuovo Cim. **19**, 154 (1961).
- [15] B. R. Holstein, (1995), hep-ph/9510344.
- [16] D. Diakonov, Chiral symmetry breaking by instantons, hep-ph/9602375, in: *Selected Topics in Nonperturbative QCD*, A. Di Giacomo and D. Diakonov, eds., Proceedings International School of Physics “Enrico Fermi”, Course CXXX, Varenna, Italy, 1995, IOS Press, Amsterdam, 1996.
- [17] J. Bjorken and S. Drell, *Relativistic Quantum Mechanics* (McGraw-Hill, 1964).
- [18] F. Yndurain, *Relativistic Quantum Mechanics and Introduction to Field Theory* (Springer, Heidelberg, 1996).

- [19] T. Newton and E. Wigner, *Rev. Mod. Phys.* **21**, 400 (1949).
- [20] M. Green, J. Schwarz, and E. Witten, *Superstring Theory* (Cambridge University Press, 1987).
- [21] A. Polyakov, *Gauge Fields and Strings* (Harwood Academic, 1987).
- [22] A. Hanson, T. Regge, and C. Teitelboim, *Constrained Hamiltonian Systems* (Accademia Nazionale dei Lincei, Rome, 1976).
- [23] K. Sundermeyer, *Constrained Dynamics* (Springer, Heidelberg, 1982).
- [24] P. A. M. Dirac, *Can. J. Math.* **2**, 129 (1950).
- [25] P. Dirac, *Lectures on Quantum Mechanics* (Benjamin, New York, 1964).
- [26] P. Bergmann, *Helv. Phys. Acta Suppl.* **4**, 79 (1956).
- [27] J. Anderson and P. Bergmann, *Phys. Rev.* **83**, 1018 (1951).
- [28] D. Gitman and I. Tyutin, *Quantization of Fields with Constraints* (Springer, Heidelberg, 1990).
- [29] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton, 1992).
- [30] C. Isham, Conceptual and geometrical problems in quantum gravity, in: *Recent Aspects of Quantum Fields*, H. Mitter and H. Gausterer, Lecture Notes in Physics Vol. 396, p. 123, Springer, Heidelberg, 1991.
- [31] H. Leutwyler and J. Stern, *Ann. Phys. (N.Y.)* **112**, 94 (1978).
- [32] P. A. M. Dirac, *Rev. Mod. Phys.* **21**, 392 (1949).
- [33] S. Fubini, A. Hanson, and R. Jackiw, *Phys. Rev.* **D7**, 1732 (1973).
- [34] A. Peres, *J. Math. Phys.* **9**, 785 (1968).
- [35] C. Sommerfield, *Ann. Phys. (N.Y.)* **84**, 285 (1974).
- [36] D. Gromes, H. Rothe, and B. Stech, *Nucl. Phys.* **B75**, 313 (1974).
- [37] R. Neville and F. Rohrlich, *Nuovo Cim.* **1A**, 625 (1971).
- [38] F. Rohrlich, *Acta Physica Austriaca, Suppl.* **8**, 277 (1971).
- [39] F. Coester, *Prog. Part. Nucl. Phys.* **29**, 1 (1992).
- [40] L. Parker and G. Schmieg, *Am. J. Phys.* **38**, 218 (1970).
- [41] H. Leutwyler, *Nuovo Cim.* **37**, 556 (1965).
- [42] X. Jaen, A. Molina, and J. Llosa, in: *Lecture Notes in Physics*, 212, Springer, Heidelberg, 1984.

- [43] S. Sokolov and A. Shatnii, *Theor. Math. Phys.* **37**, 1029 (1979).
- [44] F. Coester and W. Polyzou, *Phys. Rev.* **D26**, 1348 (1982).
- [45] E. Prokhvatilov and V. Franke, *Sov. J. Nucl. Phys.* **47**, 559 (1988).
- [46] E. Prokhvatilov and V. Franke, *Sov. J. Nucl. Phys.* **49**, 688 (1989).
- [47] F. Lenz, M. Thies, K. Yazaki, and S. Levit, *Annals Phys.* **208**, 1 (1991).
- [48] K. Hornbostel, *Phys. Rev.* **D45**, 3781 (1992).
- [49] J. Schwinger, *Phys. Rev.* **82**, 914 (1951).
- [50] J. Schwinger, *Phys. Rev.* **91**, 713 (1953).
- [51] J. Schwinger, *Phil. Mag.* **44**, 1171 (1953).
- [52] S.-J. Chang, R. G. Root, and T.-M. Yan, *Phys. Rev.* **D7**, 1133 (1973).
- [53] R. Peierls, *Proc. Roy. Soc.* **A214**, 143 (1952).
- [54] B. DeWitt, The space-time approach to quantum field theory, in: *Relativity, Groups and Topology, II*, B.S. DeWitt and R. Stora, eds., Proceedings Les Houches 1983, Session XL, North-Holland, Amsterdam, 1984.
- [55] T. Heinzl and E. Werner, *Z. Phys.* **C62**, 521 (1994), hep-th/9311108.
- [56] T.-P. Cheng and L.-F. Li, *Gauge Theory of Elementary Particle Physics* (Clarendon Press, Oxford, 1984).
- [57] S. Schweber, *An Introduction to Relativistic Quantum Field Theory* (Harper and Row, New York, 1961).
- [58] G. Domokos, in: *Boulder Lectures, Vol. XIV*, 1971, A.O. Barut, W.E. Brittin, eds., Colorado University Press, Boulder 1972.
- [59] R. Courant and D. Hilbert, *Methods of Mathematical Physics* (Interscience, New York, 1962).
- [60] T. Myint-U and L. Debnath, *Partial Differential Equations for Scientists and Engineers* (North Holland, Amsterdam, 1987).
- [61] N. Bogolubov, A. Logunov, A. Oksak, and I. Todorov, *General Principles of Quantum Field Theory* (Kluwer Academic, 1990).
- [62] I. Gelfand and G. Shilov, *Generalized Functions* (Academic Press, New York, 1964).
- [63] T. Maskawa and K. Yamawaki, *Prog. Theor. Phys.* **56**, 270 (1976).
- [64] H.-C. Pauli and S. J. Brodsky, *Phys. Rev.* **D32**, 1993 (1985).

- [65] H.-C. Pauli and S. J. Brodsky, Phys. Rev. **D32**, 2001 (1985).
- [66] H. Leutwyler, J. Klauder, and L. Streit, Nuovo Cim. **66A**, 536 (1970).
- [67] R. Jackiw, in: Springer Tracts in Modern Physics, Vol. 62, G. Höhler, ed., Springer, Heidelberg, 1972.
- [68] L. Faddeev and R. Jackiw, Phys. Rev. Lett. **60**, 1692 (1988).
- [69] R. Jackiw, (1993), hep-th/9306075.
- [70] R. Streater and A. Wightman, *PCT, Spin and Statistics, and all that* (Benjamin, 1964).
- [71] N. Bogolubov, A. Logunov, and I. Todorov, *Introduction to Axiomatic Field Theory* (Benjamin, 1975).
- [72] C. Stern, *Normalordnungsprobleme in Lichtkegelquantenfeldtheorien*, diploma thesis, Regensburg, 1994, (in German).
- [73] R. Jackiw, Functional representations for quantized fields, in: *Conformal Field Theory, Anomalies, and Superstrings*, C.K. Chew et al., eds., Proceedings of the First Asia Pacific Workshop on High Energy Physics, Singapore, 1987, World Scientific, Singapore, 1988.
- [74] W. Vogel and D.-G. Welsch, *Lectures on Quantum Optics* (Akademie-Verlag, Berlin, 1994).
- [75] S. Schlieder and E. Seiler, Comm. Math. Phys. **25**, 62 (1972).
- [76] F. Rohrlich and L. Streit, Nuovo Cim. **7B**, 166 (1972).
- [77] S.-J. Chang and S.-K. Ma, Phys. Rev. **180**, 1506 (1969).
- [78] L. Susskind, Phys. Rev. **165**, 1535 (1968).
- [79] T. Heinzl, *QCD-Vakuum und Lichtkegel-Quantisierung*, diploma thesis, Regensburg, 1988, (in German).
- [80] C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- [81] C. Dietmaier, T. Heinzl, M. Schaden, and E. Werner, Z. Phys. **A334**, 220 (1989).
- [82] S. Brodsky and G. Lepage, Exclusive processes in quantum chromodynamics, in: *Perturbative Quantum Chromodynamics*, A.H. Mueller, ed., World Scientific, Singapore, 1989.
- [83] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Clarendon Press, Oxford, 1996).

- [84] I. Tamm, J. Phys. (Moscow) **9**, 449 (1945).
- [85] S. Dancoff, Phys. Rev. **78**, 382 (1950).
- [86] J. Namysłowski, Prog. Part. Nucl. Phys. **14**, 49 (1985).
- [87] M. Lavelle, E. Werner, and S. Glazek, Few Body Systems, Suppl. **2**, 519 (1987).
- [88] F. Antonuccio, S. J. Brodsky, and S. Dalley, Phys. Lett. **B412**, 104 (1997), hep-ph/9705413.
- [89] M. Burkardt, Phys. Rev. **D57**, 1136 (1998), hep-th/9704162.
- [90] M. Terent'ev, Sov. J. Nucl. Phys. **24**, 106 (1976).
- [91] Z. Dziembowski, Phys. Rev. **D37**, 778 (1988).
- [92] W. Jaus, Phys. Rev. **D41**, 3394 (1990).
- [93] C.-R. Ji, P. Chung, and S. Cotanch, Phys. Rev. **D45**, 4214 (1992).
- [94] S. Weinberg, Phys. Rev. **150**, 1313 (1966).
- [95] S. Fubini and G. Furlan, Physics **1**, 229 (1965).
- [96] R. Dashen and M. Gell-Mann, Phys. Rev. Lett. **17**, 340 (1966).
- [97] V. D. Alfaro, S. Fubini, G. Furlan, and C. Rossetti, *Currents in Hadron Physics* (North Holland, Amsterdam, 1973).
- [98] S. Coleman, J. Math. Phys. **7**, 787 (1966).
- [99] T. Heinzl, Phys. Lett. **B388**, 129 (1996), hep-th/9608043.
- [100] T. Heinzl, S. Krusche, and E. Werner, Phys. Lett. **B256**, 55 (1991).
- [101] T. Heinzl, S. Krusche, and E. Werner, Phys. Lett. **B275**, 410 (1992).
- [102] A. C. Kalloniatis and H.-C. Pauli, Z. Phys. **C60**, 255 (1993).
- [103] A. Kalloniatis and H.-C. Pauli, Z. Phys. **C63**, 161 (1994).
- [104] A. C. Kalloniatis, H.-C. Pauli, and S. Pinsky, Phys. Rev. **D50**, 6633 (1994), hep-th/9403038.
- [105] G. McCartor and D. G. Robertson, Z. Phys. **C62**, 349 (1994), hep-th/9311065.
- [106] G. McCartor, D. G. Robertson, and S. S. Pinsky, Phys. Rev. **D56**, 1035 (1997), hep-th/9612083.
- [107] T. Heinzl, S. Krusche, S. Simbürger, and E. Werner, Z. Phys. **C56**, 415 (1992).

- [108] D. G. Robertson, Phys. Rev. **D47**, 2549 (1993).
- [109] C. M. Bender, S. S. Pinsky, and B. V. de Sande, Phys. Rev. **D48**, 816 (1993), hep-th/9212009.
- [110] S. S. Pinsky and B. van de Sande, Phys. Rev. **D49**, 2001 (1994), hep-ph/9310330.
- [111] S. S. Pinsky, B. van de Sande, and J. R. Hiller, Phys. Rev. **D51**, 726 (1995), hep-th/9409019.
- [112] T. Heinzl, C. Stern, E. Werner, and B. Zeller mann, Z. Phys. **C72**, 353 (1996), hep-th/9512179.
- [113] H. Weyl, Z. Phys. **46**, 1 (1927).
- [114] J. Moyal, Proc. Cambridge Phil. Soc. **45**, 99 (1949).
- [115] M. Hillery *et al.*, Phys. Rept. **106**, 121 (1984).
- [116] C. Grosche, (1993), hep-th/9302097.
- [117] T. Heinzl, S. Krusche, and E. Werner, (1991), Regensburg preprint, TPR-91-23.
- [118] T. Heinzl, S. Krusche, and E. Werner, Phys. Lett. **B272**, 54 (1991).
- [119] T. Heinzl, S. Krusche, E. Werner, and B. Zeller mann, (1992), Regensburg preprint, TPR-92-17.
- [120] T. Lee, *Particle Physics and Introduction to Field Theory* (Harwood Academic, Chur, 1981).
- [121] R. Wittman, in: Nuclear and Particle Physics on the Light-Cone, Proceedings, LAMPF Workshop, Los Alamos, 1988, M.B. Johnson, L.S. Kisslinger, eds., World Scientific, Singapore, 1989.
- [122] M. Abramowitz and I. Stegun, editors, *Handbook of Mathematical Functions* (Dover, New York, 1970).
- [123] D. Amit, *Field Theory, the Renormalization Group, and Critical Phenomena* (World Scientific, Singapore, 1984).
- [124] X.-M. Xu and H. J. Weber, Phys. Rev. **D52**, 4633 (1995), hep-ph/9507450.
- [125] S. Simbürger, *Methoden zur Berechnung des effektiven Potentials in der Lichtkegelquantenfeldtheorie*, diploma thesis, Regensburg, 1992, (in German).
- [126] T. Heinzl, *Spontane Symmetriebrechung in Lichtkegel-Quantenfeldtheorien*, PhD thesis, Regensburg, 1991, (in German).

- [127] G. 't Hooft, Nucl. Phys. **B75**, 461 (1974).
- [128] A. Harindranath, R. Perry, and K. Wilson, Phys. Rev. Lett **65**, 2959 (1990).
- [129] H. Bergknoff, Nucl. Phys. **B122**, 215 (1977).
- [130] R. J. Perry and A. Harindranath, Phys. Rev. **D43**, 4051 (1991).
- [131] A. Harindranath and R. Perry, Phys. Rev. **D43**, 492 (1991).
- [132] A. Harindranath, R. J. Perry, and J. Shigemitsu, Phys. Rev. **D46**, 4580 (1992).
- [133] D. Mustaki and S. Pinsky, Phys. Rev. **D45**, 3775 (1992).
- [134] S. Glazek, A. Harindranath, S. Pinsky, J. Shigemitsu, and K. Wilson, Phys. Rev. **D47**, 1599 (1993).
- [135] S. Glazek and K. Wilson, Phys. Rev. **D48**, 5863 (1993).
- [136] R. J. Perry, Ann. Phys. **232**, 116 (1994), hep-th/9402015.
- [137] K. G. Wilson *et al.*, Phys. Rev. **D49**, 6720 (1994), hep-th/9401153.
- [138] T. Sugihara and M. Yahiro, Phys. Rev. **D53**, 7239 (1996).
- [139] K. Harada and A. Okazaki, Phys. Rev. **D55**, 6198 (1997), hep-th/9610020.
- [140] C.-R. Ji, in: *Theory of Hadrons and Light-Front QCD*, S. Glazek, ed., World Scientific, Singapore 1995.
- [141] M. M. Brisudova, R. J. Perry, and K. G. Wilson, Phys. Rev. Lett. **78**, 1227 (1997), hep-ph/9607280.
- [142] B. D. Jones, R. J. Perry, and S. Glazek, Phys. Rev. **D55**, 6561 (1997).
- [143] B. D. Jones and R. J. Perry, Phys. Rev. **D55**, 7715 (1997), hep-th/9612163.
- [144] M. Schmidt, *Zur Beschreibung gebundener, relativistischer Systeme mittels Integralgleichungen in Standard- und Lichtfront-Koordinaten*, PhD thesis, Erlangen, 1995, (in German).
- [145] B. Simon and R. B. Griffiths, Comm. Math. Phys **33**, 145 (1973).
- [146] P. Grangé, P. Ullrich, and E. Werner, (1997), hep-th/9709165, to appear in Phys. Rev. **D**.
- [147] S. Coleman and R. Jackiw, Ann. Phys. **67**, 552 (1971).
- [148] B. Chibisov and A. R. Zhitnitsky, Phys. Lett. **B362**, 105 (1995), hep-ph/9502258.
- [149] M. Burkardt, Phys. Rev. **D53**, 933 (1996), hep-ph/9509226.

- [150] M. Gell-Mann, R. J. Oakes, and B. Renner, Phys. Rev. **175**, 2195 (1968).
- [151] J. Gasser and H. Leutwyler, Ann. Phys. **158**, 142 (1984).
- [152] R. Pascual and R. Tarrach, *QCD: Renormalization for the Practitioner* (Springer, Heidelberg, 1984).
- [153] A. Casher, S. Noskowicz, and L. Susskind, Nucl. Phys. **B32**, 75 (1971).
- [154] A. Casher and L. Susskind, Phys. Rev. **D9**, 436 (1974).
- [155] J. Jersak and J. Stern, Nucl. Phys. **B7**, 413 (1968).
- [156] K. Harada, T. Heinzl, and C. Stern, Phys. Rev. **D57**, 2460 (1998), hep-th/9705159.
- [157] J. Schwinger, Phys. Rev. **128**, 2425 (1962).
- [158] G. 't Hooft, Lecture given at the International School of Subnuclear Physics, Erice, Italy, 1975; in: New Phenomena in Subnuclear Physics, Part A, A. Zichichi, ed., Plenum, New York, 1977.
- [159] C. Adam, Ann. Phys. (N.Y.) **259**, 1 (1997), hep-th/9704064.
- [160] E. Abdalla and M. C. B. Abdalla, Phys. Rept. **265**, 253 (1996), hep-th/9503002.
- [161] E. Witten, Nucl. Phys. **B145**, 110 (1978).
- [162] A. Zhitnitsky, Sov. J. Nucl. Phys. **43**, 999 (1986).
- [163] S. Coleman, R. Jackiw, and L. Susskind, Ann. Phys. **93**, 267 (1975).
- [164] S. Coleman, Ann. Phys. **101**, 239 (1976).
- [165] J. H. Lowenstein and J. A. Swieca, Ann. Phys. **68**, 172 (1971).
- [166] J. Kogut and L. Susskind, Phys. Rev. **D11**, 3594 (1975).
- [167] T. Banks, L. Susskind, and J. Kogut, Phys. Rev. **D13**, 1043 (1976).
- [168] C. J. Hamer, J. Kogut, D. P. Crewther, and M. M. Mazzolini, Nucl. Phys. **B208**, 413 (1982).
- [169] E. Marinari, G. Parisi, and C. Rebbi, Nucl. Phys. **B190**, 734 (1981).
- [170] A. V. Smilga, Phys. Lett. **B278**, 371 (1992).
- [171] G. McCartor, Z. Phys. **C52**, 611 (1991).
- [172] G. McCartor, Z. Phys. **C64**, 349 (1994), hep-th/9406094.
- [173] A. C. Kalloniatis and D. G. Robertson, Phys. Lett. **B381**, 209 (1996), hep-th/9602006.

- [174] J. Gamboa, I. Schmidt, and L. Vergara, (1996), hep-th/9610173.
- [175] C. Adam, Phys. Lett. **B382**, 111 (1996), hep-th/9602175.
- [176] J. P. Vary, T. J. Fields, and H. J. Pirner, Phys. Rev. **D53**, 7231 (1996).
- [177] M. Marinov, A. Perelomov, and M. Terent'ev, JETP Lett. **20**, 225 (1974).
- [178] K. Hornbostel, S. J. Brodsky, and H.-C. Pauli, Phys. Rev. **D41**, 3814 (1990).
- [179] S. Coleman, Comm. Math. Phys. **31**, 259 (1973).
- [180] M. Burkardt, *Multiquarksysteme in der 1+1 dimensionalen QCD*, PhD thesis, Erlangen, 1989, (in German).
- [181] M. Li, Phys. Rev. **D34**, 3888 (1986).
- [182] R. Brower, W. Spence, and J. Weis, Phys. Rev. **D19**, 3024 (1979).
- [183] T. Eller, H.-C. Pauli, and S. Brodsky, Phys. Rev. **D35**, 1493 (1987).
- [184] M. Burkardt, Nucl. Phys. **A504**, 762 (1989).
- [185] T. Sugihara, M. Matsuzaki, and M. Yahiro, Phys. Rev. **D50**, 5274 (1994), hep-th/9402092.
- [186] K. Aoki and T. Ichihara, Phys. Rev. **D52**, 6435 (1995), hep-th/9506058.
- [187] Y. Mo and R. J. Perry, J. Comput. Phys. **108**, 159 (1993).
- [188] K. Harada, A. Okazaki, and M. Taniguchi, Phys. Rev. **D52**, 2429 (1995), hep-th/9502102.
- [189] W. A. Bardeen, R. B. Pearson, and E. Rabinovici, Phys. Rev. **D21**, 1037 (1980).
- [190] K. Harada, T. Sugihara, M. Taniguchi, and M. Yahiro, Phys. Rev. **D49**, 4226 (1994), hep-th/9309128.
- [191] A. R. Zhitnitsky, Phys. Rev. **D53**, 5821 (1996), hep-ph/9510366.
- [192] M. Li, M. Birse, and L. Wilets, J. Phys. **G13**, 915 (1987).
- [193] I. Bars and M. B. Green, Phys. Rev. **D17**, 537 (1978).
- [194] D. P. Crewther and C. J. Hamer, Nucl. Phys. **B170**, 353 (1980).
- [195] S. R. Carson and R. D. Kenway, Ann. Phys. **166**, 364 (1986).
- [196] S. Elser, PhD thesis, Heidelberg, 1994.
- [197] B. van de Sande, Phys. Rev. **D54**, 6347 (1996), hep-ph/9605409.

- [198] A. Carroll, J. Kogut, D. K. Sinclair, and L. Susskind, Phys. Rev. **D13**, 2270 (1976).
- [199] S. Coleman, Phys. Rev. **D11**, 2088 (1975).
- [200] M. Burkardt, Phys. Rev. **D47**, 4628 (1993).
- [201] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
- [202] C. Michael and F. Payne, Z. Phys. **C12**, 145 (1982).
- [203] S. Brodsky, C.-R. Ji, and M. Sawicki, Phys. Rev. **D32**, 1530 (1985).
- [204] H. H. Liu and D. E. Soper, Phys. Rev. **D48**, 1841 (1993).
- [205] A. Mitra *et al.*, Prog. Part. Nucl. Phys. **22**, 43 (1989).
- [206] C. D. Roberts and A. G. Williams, Prog. Part. Nucl. Phys. **33**, 477 (1994), hep-ph/9403224.
- [207] M. R. Frank and C. D. Roberts, Phys. Rev. **C53**, 390 (1996), hep-ph/9508225.
- [208] U. Vogl and W. Weise, Prog. Part. Nucl. Phys. **27**, 195 (1991).
- [209] S. P. Klevansky, Rev. Mod. Phys. **64**, 649 (1992).
- [210] T. Hatsuda and T. Kunihiro, Phys. Rept. **247**, 221 (1994), hep-ph/9401310.
- [211] J. Bijnens, Phys. Rept. **265**, 369 (1996).
- [212] R. J. Perry and K. G. Wilson, Nucl. Phys. **B403**, 587 (1993).
- [213] F. Gross, in: Nuclear and Particle Physics on the Light-Cone, Proceedings, LAMPF Workshop, Los Alamos, 1988, M.B. Johnson, L.S. Kisslinger, eds., World Scientific, Singapore, 1989.
- [214] Ç. Şavklı and F. Tabakin, (1997), hep-ph/9702251.
- [215] C. Dietmaier, *Dynamische Brechung der chiralen Symmetrie im NJL-Modell in Lichtkegel-Quantisierung*, diploma thesis, Regensburg, 1988, (in German).
- [216] G. Lepage and S. Brodsky, Phys. Rev. **D22**, 2157 (1980).
- [217] G. Lepage, S. Brodsky, T. Huang, and P. Mackenzie, Proceedings of the Banff Summer Institute 1981.
- [218] H. Melosh, Phys.Rev. **D9**, 1095 (1974).
- [219] P. Chung, F. Coester, and W. Polyzou, Phys. Lett. **B205**, 545 (1988).
- [220] C.-R. Ji and S. Cotanch, Phys. Rev. **D41**, 2319 (1990).

- [221] H. Leutwyler, Phys. Lett. **B48**, 45 (1974).
- [222] H. Leutwyler, Nucl. Phys. **B76**, 413 (1974).
- [223] S. Drell and T.-M. Yan, Phys. Rev. Lett. **24**, 181 (1970).
- [224] S. R. Amendolia *et al.*, Nucl. Phys. **B277**, 168 (1986).
- [225] B. R. Holstein, Phys. Lett. **B244**, 83 (1990).
- [226] A. Manohar and H. Georgi, Nucl. Phys. **B234**, 189 (1984).
- [227] W. Weise, in: International Review of Nuclear Physics, Vol.1, Quarks and Nuclei, W. Weise, ed., World Scientific, Singapore, 1984.
- [228] V. Bernard, R. Brockmann, and W. Weise, Regensburg Preprint, TPR 85-02, 1985.
- [229] V. Chernyak and A. Zhitnitsky, Phys. Rept. **112**, 173 (1984).
- [230] CLEO Collaboration, V. Savinov, (1995), hep-ex/9507005.
- [231] P. Kroll and M. Raulfs, Phys. Lett. **B387**, 848 (1996).
- [232] V. Belyaev, hep-ph/9703244.
- [233] V. Belyaev and M. Johnson, hep-ph/9707329.
- [234] V. Y. Petrov and P. Pobylitsa, hep-ph/9712203.
- [235] T. Heinzl, (1995), hep-th/9604018.
- [236] E. Tomboulis, Phys. Rev. **D8**, 2736 (1973).
- [237] A. Casher, Phys. Rev. **D14**, 452 (1976).
- [238] S. Brodsky, H.-C. Pauli, and S. Pinsky, (1997), hep-ph/9705477.
- [239] D. Kaplan, nucl-th/9506035.
- [240] A. Manohar, Effective field theories, hep-ph/9606222, in: *Perturbative and Nonperturbative Aspects of Quantum Field Theory*, H. Latal and W. Schweiger, eds., Lecture Notes in Physics, Vol. 479, p. 311, Springer, Heidelberg, 1997.
- [241] H. Leutwyler, in: Springer Tracts in Modern Physics, Vol. 50, G. Höhler, ed., Springer, Heidelberg, 1969.